



US00PP24613P3

(12) **United States Plant Patent**
Weber(10) **Patent No.:** US PP24,613 P3
(45) **Date of Patent:** Jul. 8, 2014(54) **STRAWBERRY PLANT NAMED 'HERRIOT'**(50) Latin Name: *Fragaria×ananassa*
Varietal Denomination: Herriot(71) Applicant: **Cornell University**, Ithaca, NY (US)(72) Inventor: **Courtney A. Weber**, Geneva, NY (US)(73) Assignee: **Cornell University**, Ithaca, NY (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 111 days.

(21) Appl. No.: **13/573,626**(22) Filed: **Sep. 28, 2012**(65) **Prior Publication Data**

US 2014/0096296 P1 Apr. 3, 2014

(51) **Int. Cl.**
A01H 5/00 (2006.01)(52) **U.S. Cl.**USPC **Plt./209**(58) **Field of Classification Search**USPC Plt./209
See application file for complete search history.*Primary Examiner* — Annette Para(74) *Attorney, Agent, or Firm* — MacMillan, Sobanski & Todd, LLC(57) **ABSTRACT**

This invention relates to a new and distinct June-bearing (short day responsive) strawberry plant named 'Herriot' primarily adapted to the growing conditions of the west central New York and other regions of similar climate. The new plant is primarily characterized by strong vigor, longer than broad fruit, conical fruit shape, uniformity in shape between primary and secondary fruit, reflexed calyx, bright red fruit color, glossy fruit skin and highly pigmented petioles.

5 Drawing Sheets**1**Genus and species: *Fragaria×ananassa*.

Variety denomination: 'Herriot'.

BACKGROUND AND SUMMARY OF THE INVENTION**I. Field & Utility Summary**

The present invention to a new and distinct June-bearing (short day responsive) strawberry plant named 'Herriot.' Strawberry plant 'Herriot' is primarily adapted to the growing conditions of the west central New York and other regions of similar climate. The new plant is primarily characterized by strong vigor, longer than broad fruit, conical fruit shape, uniformity in shape between primary and secondary fruit, reflexed calyx, bright red fruit color, glossy fruit skin and highly pigmented petioles.

Test plantings in various locations at in New York, Illinois, Minnesota, Michigan, and Ontario, Canada have shown this plant to be widely adapted to differing soil and climactic conditions. It has shown cold hardiness typical to strawberries in a matted row system when covered with straw in the winter, which is standard procedure for this growing region.

The new plant is primarily adapted to the climate and growing conditions of west central NY, the upper mid-western U.S., the mid-Atlantic states, and southeastern Canada. This region provides the necessary winter temperatures required for it to produce a strong vigorous plant and to produce fruit in the summer harvest season from May to July, depending on location.

II. Cultivation Summary

'Herriot' originated from hand-pollinated hybridization in 1997 in Geneva, N.Y.

III. Comparisons

NYUS299, the proprietary female parent (unpatented) is June-bearing with smaller average fruit size that is medium red, firm fleshed, and conic shaped that ripens in a similar season as 'Herriot'.

2

'MNUS 210', the proprietary male plant, commonly known as 'Winona' in the industry (U.S. Plant Pat. No. 10,191) is June-bearing with similar fruit size that is bright red, conic shaped, moderately firm fleshed and ripens later than 'Herriot'.

The varieties which are believed to most closely resemble 'Herriot' are 'Honeoye' (not patented), 'L'Amour' (U.S. Plant Pat. No. 16,480) and 'Jewel' (U.S. Plant Pat. No. 5,897).

In comparison to the similar plant 'Honeoye', 'Herriot' differs by the following characteristics. The plant of 'Herriot' is shorter and more upright in growth habit with greater vigor. The fruit of 'Herriot' is larger than that of 'Honeoye' with a more reflexed calyx. The fruit skin and flesh color of 'Herriot' is lighter red than that of 'Honeoye'. The yield of 'Herriot' is larger than that of 'Honeoye'. The fruit of 'Herriot' is firmer with glossier skin than that of 'Honeoye'. The fruit of 'Herriot' has lower soluble solids (% Brix) and lower titratable acids than that of 'Honeoye' with lower juice pH. The harvest season of 'Herriot' is later than that of 'Honeoye' and slightly less compact.

In comparison to the similar plant, 'L'Amour', 'Herriot' differs by the following combination of characteristics. The plant of 'Herriot' is shorter and smaller with more upright leaves. The leaflets are less wide and similar in length giving them a greater length to width ratio and an ovate shape. The stipules of 'Herriot' are shorter and less wide than those of 'L'Amour'. The fruit of 'Herriot' is smaller in size, with lighter red flesh color than 'L'Amour'. The fruit of 'Herriot' has lower average soluble solids, lower titratable acids and higher juice pH than 'L'Amour'.

The flowering trusses of 'Herriot' are shorter than those of 'L'Amour'. The flowers of 'Herriot' are smaller in diameter than the flowers of 'L'Amour' with a greater petal length to width ratio than 'L'Amour'. The harvest season for 'Herriot' is more compact than that of 'L'Amour'.

In comparison to the similar plant, 'Jewel', 'Herriot' differs by the following combination of characteristics. The plant of

'Herriot' is more vigorous but with shorter stature than 'Jewel'. The foliage is smaller in size with less interveinal leaf blistering. Double petiole bracts are occasionally present in 'Herriot' and absent in 'Jewel'. The leaves are more sharply serrated in 'Herriot' than 'Jewel'. The stipules of 'Herriot' are shorter and less broad than those of 'Jewel'.

The fruit of 'Herriot' is darker red in color than 'Jewel' with more recessed seeds and a more reflexed calyx. The fruiting trusses of 'Herriot' are shorter in overall length than those of 'Jewel'. The fruit of 'Herriot' has lower soluble solids and lower titratable acidity than that of 'Jewel'.

Strawberry plant 'Herriot' has these distinguishing characteristics:

1. short plants with an upright growth habit and strong vigor;
2. predominantly trifoliate leaves with 4 and 5 leaflet leaves common with high leaf gloss and ovate leaflets with greater length than width;
3. longer than broad fruit, conical fruit shape, uniform shape between primary and secondary fruit, bright red fruit color, light red flesh color with ring of white flesh in the center and a small light red cavity, high fruit gloss and moderate flesh firmness;
4. reflexed calyx generally equal or smaller in diameter than the fruit width;
5. short pedicels and peduncles with few bracts;
6. mid-season production with high yields; and
7. compact harvest season.

IV. Breeding History

Strawberry plant 'Herriot' originated from a hand-pollinated hybridization made in 1997 in Geneva, N.Y. between NYUS299×'MNUS 210'. Both parents of 'Herriot' are hybrids of the strawberry genus of the species *Fragaria x ananassa*. Thus 'Herriot' is of the species *Fragaria x ananassa*. The seeds resulting from this controlled hybridization were germinated in a greenhouse during the winter of 1997-98. Resulting seedlings were planted in the spring of 1998 in a field in Geneva, N.Y. The seedlings fruited in the summer of 1999 and one, designated NY99-21 ('Herriot'), was selected for its early season, large, attractive, bright red fruit, strong strawberry flavor, and firm flesh.

V. Asexual Reproduction

During 1999, the original plant selection was propagated asexually by stolons (runners) at Geneva, N.Y. and a test planting of ten plants was established. Subsequently, larger test plantings have been established with asexually multiplied plants at Geneva, N.Y. The new plant was then tested over the next several years in the fruiting fields at Geneva, N.Y. Further propagation was done using tissue culture to produce disease free material for use in further testing. Tissue culture plants were used as mother plants for propagation from stolons.

VI. Stability

Asexual propagation has demonstrated that the combination of traits disclosed herein as characterizing strawberry plant 'Herriot' are fixed and remain true to type through successive generations of asexual reproduction. All propagules of strawberry plant 'Herriot' have been observed to be true to type in that during all asexual multiplication, the vegetative and fruit characteristics of the original plant have

been maintained. All plants planted from dormant crowns or rooted runner tips have fruited after one season of growth in the field.

BRIEF DESCRIPTION OF THE FIGURES

The accompanying color photographs show typical specimens of the new plant at various stages of development as nearly true as it is possible to make in color reproductions. The depicted plant and plant parts were from the first harvest season, approximately 13 months after planting.

FIG. 1 shows typical external fruit characteristics of 'Herriot' (A) including calyx structure and fruit shape compared to 'Jewel' (B) on Jun. 6, 2012.

FIG. 2 shows typical internal fruit characteristics of 'Herriot' (A) compared to 'Jewel' (B) on Jun. 6, 2012.

FIG. 3 shows petiole pigmentation of 'Herriot' (A-three on the left) compared to 'Jewel' (B-three on the right).

FIG. 4 shows leaf margin serrations for 'Herriot' (A), 'Jewel' (B) and 'L'Amour' (C).

DETAILED BOTANICAL DESCRIPTION OF THE INVENTION

The following description of 'Herriot' unless otherwise noted, is based on observations taken during the 2012 growing season in Geneva, N.Y. These measurements and ratings were taken from plants planted in May 2011. The age of the planting is approximately 13 months and in its first harvest season. Yield observations and fruit quality characteristics are averaged from data collected during the 2003-04 and 2008-09 production seasons. The characteristics of the new plant may vary in detail, depending upon variations in environmental factors (temperature, rainfall, humidity and light intensity). 'Herriot' has not been observed under all possible environmental conditions. Color terminology where noted follows The Royal Horticultural Society Colour Chart, London.

Comparative Fruit Characteristics

'Herriot' fruit, fruit production and fruit quality characteristics. Fruit characteristics are taken from the first harvest season.

TABLE 1

Cultivar (kg per ha)	Fruit characteristics: Yield				
	2003	2004	2008	2009	Average
'Herriot'	8,740	24,790	4,910	20,010	14,610
'Honeoye'	11,730	17,160	3,990	7,970	6,360
'L'Amour'	7,090	14,490	—	—	10,790
'Jewel'	11,160	15,440	11,270	5,760	10,910

Total fruit yield of 'Herriot' and similar varieties from 2 separate trials, one harvested in 2003-04 and one in 2008-09 in Geneva, NY. Fruit was harvested in June in each year. The plants of 'Herriot' and the other varieties were grown in a nursery in South Deerfield, Mass. and planted in May prior to the first harvest year for both trials.

TABLE 2

Cultivar (g per berry)	Fruit Characteristics: Weight				
	2003	2004	2008	2009	Average
'Herriot'	10.8	10.6	10.7	11.2	10.8
'Honeoye'	13.2	10.0	9.5	9.2	10.5

TABLE 2-continued

Fruit Characteristics: Weight					
Cultivar (g per berry)	2003	2004	2008	2009	Average
'L'Amour'	11.9	11.7	—	—	11.8
'Jewel'	10.9	10.6	11.1	11.1	10.9

Average fruit weight of 'Herriot' and similar varieties from 2 separate trials, one harvested in 2003-04 and one in 2008-09 in Geneva, NY. Fruit was harvested in June in each year. The plants of 'Herriot' and the other varieties were grown in a nursery in South Deerfield, Mass. and planted in May prior to the first harvest year for both trials.

TABLE 3

Fruit Characteristics: Color				
	Character RHS Exterior Color Mature Fruit			
	'Herriot' red-purple 45A	'L'Amour' red 46B	'Jewel' red 44A	'Honeoye' red 46A
Fruit Length mean (cm)	3.70	4.22	3.46	3.85
Fruit Width mean (cm) ¹	3.40	3.62	3.61	3.55
Fruit Length/ Width Ratio	1.10	1.20	1.05	1.08
No. Sepals/ Berry	12.3	11.0	12.4	12.7

Comparison of secondary fruit characteristics of 'Herriot', with standards from Geneva, NY Jun. 6, 2012.

¹ Width is measured across the widest part of the berry, typically across the shoulders.

TABLE 4

Fruit Characteristics: Quality				
Character	'Herriot'	'L'Amour'	'Jewel'	'Honeoye'
Soluble solids ^{1,2}	9.03	10.37	11.77	10.93
Titratable acidity ^{2,3}	9.29	9.64	10.93	10.89
Juice pH ²	3.38	3.50	3.38	3.42

Comparison of 2010 fruit quality characteristics and soluble solids (% Brix), titratable acidity and juice pH of 'Herriot', with standards from Geneva, NY.

¹ soluble solids is estimated from % Brix with % Brix being an indirect measurement of the sugar content in the fruit

² Averaged from 3 samples on 3 dates during the harvest season.

³ g per L equivalents of citric acid

TABLE 5

Fruit Characteristics: Details	
Characteristic	Description
Ratio of length/width	Longer than broad
Size	Medium to large
Aroma	Moderate to high
Predominant shape	Conical
Difference in shapes between primary and secondary fruit	Little to none
Band without achenes	Intermediate width
Color of mature fruit	Bright to dark red
Evenness of color	Very even
Glossiness	Strong
Insertion of achenes	Level to slightly recessed from surface

TABLE 5-continued

Fruit Characteristics: Details		
Characteristic	Description	
Attitude of the calyx segments	Somewhat reflexed	
Size of calyx in relation to fruit diameter	Generally equal to or smaller	
Adherence of calyx (when fully ripe)	Strong	
Firmness of skin	Moderately firm	
Color of flesh	Medium red with lighter ring (red 45A to 41B).	
Distribution of red color of the flesh	Marginal through central with white inner ring (white N155C)	
Hollow center	Moderately to strongly expressed	
Seed color	Medium yellow-green to dark red (yellow-green 153D to red 46B).	
Time of flowering (50% of plants at first flower)	Medium to early	
Time of ripening (50% of plants with first ripe fruit)	Medium to early	
Type of bearing	Fully short day responsive	

Plant characteristics:

'Herriot' plant characteristics. Plant characteristics are taken from a fully mature mid-season plant.

TABLE 6

Plant Characteristics: Comparisons			
Character	'Herriot'	'L'Amour'	'Jewel'
Plant Height mean (cm)	21.9	31.9	27.3
Vigor ¹	8.0	8.3	5.0
Canopy density ¹	7.5	7.7	7.0

Comparison of plant characteristics of 'Herriot', with standards from Geneva, NY Jun. 22, 2012.

¹ Rated on a scale of 1-9 (9 being more vigorous or dense) in mature replicated plots.

TABLE 7

Plant Characteristics: Details		
Characteristic	Description	
Size	Medium	
Habit	Upright	
Density	High	
Vigor	Strong	

Foliage Characteristics

'Herriot' foliage characteristics. Foliage characteristics are taken from a fully mature tri-foliate leaf during mid season.

TABLE 8

Foliage Characteristics: Comparisons			
Character	'Herriot'	'L'Amour'	'Jewel'
RHS Color (upper surface)	green 137A	green 137A	green 138A
RHS Color (lower surface)	yellow-green 148B	green 139C	green 137D
Terminal Leaflet Length mean (cm)	8.3	8.4	7.5
Terminal Leaflet Width mean (cm)	7.0	8.5	7.6
Terminal Leaflet ratio (L/W)	1.18	0.99	0.99
Petiole Length mean (cm)	17.7	22.8	19.9

US PP24,613 P3

7

8

TABLE 8-continued

Foliage Characteristics: Comparisons			
Character	'Herriot'	'L'Amour'	'Jewel'
Serrations/Leaf	26.1	22.3	18.3
Stipule Length mean (cm)	1.7	3.5	3.0
Stipule Width mean (cm)	0.4	0.8	0.8

Comparison of leaf characteristics of 'Herriot', with standards from Geneva, NY on Jun. 22, 2012.

TABLE 9

Foliage Characteristics: Details	
Characteristic	Description
Color of upper surface	Medium to dark green
Color of under side	Medium to light green with yellow undertone.
Shape in cross section	Slightly concave
Interveinal blistering	Light to moderate
Glossiness	Moderate
Number of leaflets/leaf	Generally three with four or five common
Terminal leaflet size	Medium
Terminal leaflet - length/width ratio	Longer than broad
Terminal leaflet - shape of base	Cuneate
Terminal leaflet - shape serrations	Acute
Petiole - pubescence density	Moderate to high.
Petiole - stilule color	Light to medium yellow-green (yellow-green N144C)
Petiole - anthocyanin coloration of stipule	Little to none.
Petiole - attitude of hairs	Strongly outward
Petiole - size of bract leaflets	Small
Petiole - frequency of bract leaflets	Approximately 10% of the petioles

Flower and Inflorescence Characteristics

'Herriot' inflorescence and flower characteristics. Inflorescence characteristics are taken from a fully mature plant during full bloom. Flower characteristics are taken from a secondary flower during mid season at full maturity:

TABLE 10

Flower and Inflorescence Characteristics: Comparisons			
Character	'Herriot'	'L'Amour'	'Jewel'
Fruiting Truss Length ¹ mean (cm)	14.8	32.3	20.9
Corolla Diameter mean (mm)	32	36	31
Calyx Diameter mean (mm)	32	33	34
Petal Length mean (mm)	13.8	15.4	14.0
Petal Width mean (mm)	11.8	13.6	13.3
Petal L/W Ratio	1.16	1.13	1.09
Petals/Flower (mean)	5.8	5.1	6.3
Sepals/Flower (mean)	12.3	11.0	12.4

Comparison of inflorescence and secondary flower characteristics of 'Herriot', with standards from Geneva, NY on May 10, 2012.
¹ as measured from the base of the primary peduncle where it attaches to the crown of the plant to the furthest berry.

TABLE 11

Inflorescence Characteristics: Details	
Characteristic	Description
Position relative to foliage	Above
Fruiting truss length	Short

TABLE 12

Flower Characteristics: Details	
Characteristic	Description
Color	White (white 155C).
Size	Medium to large
Size of calyx relative to corolla	Equal to smaller
Relative position of petals	Slight overlap when first fully open.
Petal length/width ratio	Longer than broad
Petal shape	Obovate

Pest reactions: 'Herriot' is moderately resistant to the two-spotted spider mite and susceptible to aphids and flower thrips. It is resistant to leaf spot, scorch and blight. It is slightly susceptible to powdery mildew.

I claim:

1. A new and distinct strawberry plant cultivar as herein described and illustrated.

* * * * *

45




Fig. 1A




Fig. 1B




Fig. 2A




Fig. 2B




Fig. 3




Fig. 4A




Fig. 4B




Fig. 4C