



US00PP24910P2

(12) **United States Plant Patent**  
**Dümmen**

(10) **Patent No.:** **US PP24,910 P2**  
(45) **Date of Patent:** **Sep. 23, 2014**

- (54) **MANDEVILLA PLANT NAMED ‘DUEMIAWIBU’**
- (50) Latin Name: *Mandevilla sanderi*  
Varietal Denomination: **Duemiwibu**
- (71) Applicant: **Tobias Dümmen**, Rheinberg (DE)
- (72) Inventor: **Tobias Dümmen**, Rheinberg (DE)
- (73) Assignee: **Dümmen Group B.V.**, De Lier (NL)
- (\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 96 days.
- (21) Appl. No.: **13/694,137**
- (22) Filed: **Oct. 31, 2012**
- (51) **Int. Cl.**  
*A01H 5/00* (2006.01)
- (52) **U.S. Cl.**  
USPC ..... **Plt./232**

(58) **Field of Classification Search**  
USPC ..... Plt./232  
See application file for complete search history.

(56) **References Cited**  
**U.S. PATENT DOCUMENTS**  
PP15,202 P2 \* 10/2004 Misato et al. .... Plt./232  
\* cited by examiner

*Primary Examiner* — Wendy C Haas  
(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**  
A new and distinct cultivar of *Mandevilla* plant named ‘Duemiawibu’, characterized by its upright to somewhat outwardly spreading plant habit; vigorous growth habit; freely branching habit; early and freely flowering habit; light pink-colored flowers; and good garden performance.

**1 Drawing Sheet**

**1**

Botanical designation: *Mandevilla sanderi*.  
Cultivar denomination: ‘DUEMIAWIBU’ U.S. Plant patent application Ser. No. 13/694,138.

CROSS-REFERENCED TO CLOSELY RELATED APPLICATIONS

Title: *Mandevilla* Plant Named ‘Duemiare’  
Applicant: Tobias Dümmen  
Filed: Concurrently with this application

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Mandevilla* plant, botanically known as *Mandevilla sanderi* and hereinafter referred to by the name ‘Duemiawibu’.

The new *Mandevilla* plant is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. The objective of the breeding program is to create new vigorous, freely branching and early flowering *Mandevilla* plants with numerous attractive flowers.

The new *Mandevilla* plant originated from a cross-pollination made by the Inventor in Rheinberg, Germany in July, 2007 of a proprietary selection of *Mandevilla sanderi* identified as code number F-19-303-11, not patented, as the female, or seed parent with a proprietary selection of *Mandevilla sanderi* identified as code number F-04-08-3, not patented, as the male, or pollen, parent. The new *Mandevilla* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rheinberg, Germany in May, 2011.

Asexual reproduction of the new *Mandevilla* plant by cuttings in Rheinberg, Germany since July, 2011 has shown that

**2**

the unique features of this new *Mandevilla* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Mandevilla* have not been observed under all possible environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Duemiawibu’. These characteristics in combination distinguish ‘Duemiawibu’ as a new and distinct *Mandevilla* plant:

1. Upright to somewhat outwardly spreading plant habit.
2. Vigorous growth habit.
3. Freely branching habit.
4. Early and freely flowering habit.
5. Light pink-colored flowers.
6. Good garden performance.

Plants of the new *Mandevilla* can be compared to plants of the female parent selection. Plants of the new *Mandevilla* differ primarily from plants of the female parent selection in flower color as plants of the female parent selection have light red-colored flowers.

Plants of the new *Mandevilla* can be compared to plants of the male parent selection. Plants of the new *Mandevilla* differ primarily from plants of the male parent selection in flower color as plants of the male parent selection have white-colored flowers.

Plants of the new *Mandevilla* can be compared to plants of *Mandevilla sanderi* ‘Duemiare’, disclosed in U.S. Plant patent application Ser. No. 13/694,138. Plants of the new *Mandevilla* differ primarily from plants of ‘Duemiare’ in the following characteristics:

1. Plants of the new *Mandevilla* had larger leaves than plants of 'Duemiare'.
2. Plants of the new *Mandevilla* have larger flowers than plants of 'Duemiare'.
3. Plants of the new *Mandevilla* and 'Duemiare' differ in flower color as plants of 'Duemiare' have red-colored flowers.

Plants of the new *Mandevilla* can also be compared to plants of *Mandevilla hybrida* 'Sunmandecos', disclosed in U.S. Plant Pat. No. 15,202. In side-by-side comparisons conducted in Rheinberg, Germany, plants of the new *Mandevilla* differed primarily from plants of 'Sunmandecos' in the following characteristics:

1. Plants of the new *Mandevilla* were broader than plants of 'Sunmandecos'.
2. Plants of the new *Mandevilla* had smaller leaves than plants of 'Sunmandecos'.
3. Plants of the new *Mandevilla* and 'Sunmandecos' differed in flower color as 'Sunmandecos' had deep purplish pink-colored flowers.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Mandevilla* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the following detailed botanical description which accurately describe the actual colors of the new *Mandevilla* plant.

The photograph comprises a side perspective view of a typical flowering plant of 'Duemiawibu' grown in a container.

#### DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown in 10.5-cm containers during the summer in a glass-covered greenhouse in Rheinberg, Germany and under cultural practices typical of commercial *Mandevilla* production. During the production of the plants, day and night temperatures averaged 18° C. and light levels averaged 4,500 lux. Plants were pinched one time and were 20 weeks old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Mandevilla sanderi* 'Duemiawibu'.  
Parentage:

*Female, or seed, parent.*—Proprietary selection of *Mandevilla sanderi* identified as code number F-19-303-11, not patented.

*Male, or pollen, parent.*—Proprietary selection of *Mandevilla sanderi* identified as code number F-04-08-3, not patented.

Propagation:

*Type.*—By cuttings.

*Time to initiate roots, summer.*—About five days at temperatures about 20° C.

*Time to initiate roots, winter.*—About seven days at temperatures about 20° C.

*Time to produce a rooted young plant, summer.*—About three weeks at temperatures about 20° C.

*Time to produce a rooted young plant, winter.*—About four weeks at temperatures about 20° C.

*Root description.*—Fine, fibrous; white in color.

*Rooting habit.*—Freely branching; dense.

Plant description:

*Plant and growth habit.*—Upright and outwardly spreading plant habit; vigorous growth habit; moderate growth rate.

*Plant height.*—About 78 cm.

*Plant diameter.*—About 49 cm.

*Lateral branch description.*—Branching habit: Freely branching habit with about 15 to 20 lateral branches developing per plant; pinching enhances lateral branch development. Length: About 30 cm. Diameter: About 2 mm. Internode length: About 6.3 cm. Strength: Strong; flexible. Texture: Smooth, glabrous. Color: Close to 144A tinged with close to 200C.

Foliage description:

*Arrangement.*—Opposite, simple.

*Length.*—About 8.1 cm.

*Width.*—About 4.8 cm.

*Shape.*—Elliptical.

*Apex.*—Apiculate.

*Base.*—Rounded or truncate.

*Margin.*—Entire.

*Texture, upper and lower surfaces.*—Smooth, glabrous.

*Venation pattern.*—Pinnate, arcuate.

*Color.*—Developing leaves, upper surface: Close to 137A. Developing leaves, lower surface: Close to 137C. Fully expanded leaves, upper surface: Close to 137A; venation, close to 144C. Fully expanded leaves, lower surface: Close to 137C; venation, close to 144D.

*Petioles.*—Length: About 1.1 cm. Diameter: About 2.5 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper surface: Close to 143B. Color, lower surface: Close to 143D.

Flower description:

*Flower type and habit.*—Salverform flowers alternately arranged in terminal and axillary racemose clusters with about five to six flowers per cluster, freely flowering habit with about 40 to 50 flowers developing per plant; flowers face upright to somewhat outwardly.

*Natural flowering season.*—Early flowering habit and long flowering period, plants begin flowering about nine weeks after planting and flower continuously throughout the summer in Germany.

*Flower longevity on the plant.*—About three to five days; flowers persistent.

*Fragrance.*—None detected.

*Flowers.*—Appearance: Salverform with five-parted fused corolla; flowers roughly star-shaped. Diameter: About 8.8 cm. Depth (length): About 5.1 cm. Throat diameter: About 1.9 cm. Tube length: About 3 cm. Tube diameter at the base: About 4.5 mm.

*Flower buds.*—Height: About 4 cm. Diameter: About 6.8 mm. Shape: Elongated oblong, spindle-shaped. Color: Close to 149D and 65B.

*Corolla.*—Arrangement and appearance: Single whorl of five petals; petals not imbricate. Petal lobe length: About 3.5 cm. Petal lobe width: About 2.8 cm. Petal lobe shape: Roughly spatulate. Petal apex: Apiculate. Petal margin: Entire. Petal texture, upper and lower surfaces: Smooth, glabrous. Throat texture: Smooth, glabrous. Tube texture: Smooth, glabrous. Color: Petal lobe, when opening and fully opened, upper surface: Close to 62D; venation, close to 69A; color does not change with development. Petal lobe, when

opening and fully opened, lower surface: Close to 62C; venation, close to 69B. Throat: Close to 17A; venation, close to 14A. Tube: Close to 62C; venation, close to 144B.

*Sepals*.—Arrangement and appearance: Single whorl of 5  
five sepals; sepals not imbricate. Length: About 9.2  
mm. Width: About 2 mm. Shape: Ligulate. Apex:  
Acuminate. Margin: Entire. Texture, upper and lower  
surfaces: Smooth, glabrous. Color, upper and lower  
surfaces: Close to 145B.

*Peduncles*.—Length: About 6 cm. Diameter: About 2  
mm. Aspect: About 45° from stem axis. Texture:  
Smooth, glabrous. Strength: Strong, flexible. Color:  
Close to 154C.

*Pedicels*.—Length: About 1.9 cm. Diameter: About 2 15  
mm. Aspect: About 45° from stem axis. Texture:  
Smooth, glabrous. Strength: Strong, flexible. Color:  
Close to 154C.

*Reproductive organs*.—Stamens: Quantity per flower:  
Typically five. Anther length: About 0.5 mm. Anther 20

shape: Elongated oblong. Anther color: Close to 8B.  
Pollen amount: Moderate. Pollen color: Close to 3C.  
Pistils: Quantity per flower: Typically one. Pistil  
length: About 3.5 cm. Stigma shape: Star-shaped;  
five-lobed. Stigma color: Close to 138B. Style length:  
About 2.3 cm. Style color: Close to 1D. Ovary color:  
Close to 1D.

*Seeds and fruits*.—Seed and fruit production has not  
been observed on plants of the new *Mandevilla*.

10 Disease & pest resistance: Plants of the new *Mandevilla* have  
not been noted to be resistant to pathogens and pests com-  
mon to *Mandevilla* plants.

Garden performance: Plants of the new *Mandevilla* have been  
observed to have good garden performance and to tolerate  
rain, wind, full sunlight and temperatures from about 5° C.  
to about 40° C.

It is claimed:

1. A new and distinct *Mandevilla* plant named 'Duemi-  
awibu' as illustrated and described.

\* \* \* \* \*

