



(12) **United States Plant Patent**
Li

(10) **Patent No.:** **US PP24,955 P3**
(45) **Date of Patent:** **Oct. 7, 2014**

(54) **VERBENA PLANT NAMED ‘KLEVP12449’**

(50) Latin Name: *Verbena*×*hybrida*
Varietal Denomination: **KLEVP12449**

(71) Applicant: **Klemm+Sohn GmbH & Co. KG,**
Stuttgart (DE)

(72) Inventor: **Ruijun Li,** North Parramata (AU)

(73) Assignee: **Klemm+Sohn GmbH & Co. KG,**
Stuttgart (DE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 93 days.

(21) Appl. No.: **13/694,207**

(22) Filed: **Nov. 7, 2012**

(65) **Prior Publication Data**

US 2014/0130220 P1 May 8, 2014

(51) **Int. Cl.**

A01H 5/00 (2006.01)

(52) **U.S. Cl.**

USPC **Plt./308**

(58) **Field of Classification Search**

USPC **Plt./308**

See application file for complete search history.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — Jondle Plant Sciences
Division

(57) **ABSTRACT**

A new and distinct variety of *Verbena* plant named
‘KLEVP12449’, particularly characterized by a large flower
with a white eye, a freely branching growth habit, and
extremely good branching, is disclosed.

1 Drawing Sheet

1

Genus and species: *Verbena*×*hybrida*.
Variety denomination: ‘KLEVP12449’.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new and distinct variety
of *Verbena*, botanically known as *Verbena*×*hybrida*, and
hereinafter referred to by the variety name ‘KLEVP12449’.
The new variety is the result of a cross conducted in April of
2008 in Cobbity, New South Wales, Australia between female
parent *Verbena* plant, ‘1S112’ (unpatented), and male parent
Verbena plant, ‘V842’ (unpatented). A single plant selection
was subsequently chosen for further evaluation and asexual
propagation.

The new variety was first propagated in October 2009 in
Cobbity, New South Wales, Australia by vegetative cuttings
and has been asexually reproduced repeatedly by vegetative
cuttings for three to four generations. ‘KLEVP12449’ has
been found to retain its distinctive characteristics through
successive asexual propagations.

Plant Breeder’s Rights for this variety were applied for in
Canada on Feb. 6, 2012. ‘KLEVP12449’ has not been made
publicly available anywhere in the world more than one year
prior to filing of this application.

SUMMARY OF THE INVENTION

The following are the most outstanding and distinguishing
characteristics of this new variety when grown under normal
horticultural practices in a greenhouse in Sydney, Australia.

1. Large flower with white eye;
2. Freely branching growth habit; and
3. Extremely good branching.

DESCRIPTION OF THE PHOTOGRAPH

This new *Verbena* plant is illustrated by the accompanying
photograph which shows the foliage and inflorescences of the

2

plant. The colors shown are as true as can be reasonably
obtained by conventional photographic procedures. The pho-
tograph is of a plant about 3 months old, taken in the spring of
2012, grown from a rooted cutting in a glass greenhouse in
Stuttgart, Germany under normal horticultural practices.

DESCRIPTION OF THE NEW VARIETY

The following detailed descriptions set forth the distinctive
characteristics of ‘KLEVP12449’. The data which define
these characteristics were collected from asexual reproduc-
tions carried out in Stuttgart, Germany. The plant history was
taken in the spring of 2012 on 3-month old plants grown in 12
centimeter pots in a glass greenhouse with a pinch date at
week 10. The color readings were determined under natural
light. Color references are to The R.H.S. Colour Chart of The
Royal Horticultural Society of London (R.H.S.), Fifth Edi-
tion (2007).

DESCRIPTION OF THE NEW PLANT

Classification:

Family.—Verbenaceae.

Genus and species.—*Verbena*×*hybrida*.

Common name.—*Verbena*.

Denomination.—‘KLEVP12449’.

Parentage:

Female parent.—*Verbena* plant, ‘1S112’ (unpatented).

Male parent.—*Verbena* plant, ‘V842’ (unpatented).

Plant:

Form.—Semi-trailing.

Growth and branching habit.—Freely branching.

Height (from top of soil).—Approximately 10.0 cm.

Width (horizontal plant diameter).—Approximately
40.0 cm.

Time to produce a finished flowering plant.—Approxi-
mately 8 weeks.

Time to initiate and develop roots.—3 weeks.

Root description.—Fine; white, RHS 155B.

Stems:

Number of branches per plant.—Greater than 20.

Length.—Approximately 20.0 cm.

Diameter per branch (measured from the midpoint).—
2.0 mm.

Internode length.—3.0 cm.

Color.—RHS 139C.

Anthocyanin.—Absent.

Texture and appearance.—Short pubescence.

Leaves:

Arrangement.—Opposite.

Color.—Upper surface (both immature and mature leaves): RHS 139C. Lower surface (both immature and mature leaves): RHS 139D.

Length.—2.5 cm.

Width.—1.5 cm.

Shape.—Elliptic.

Apex.—Cuspidate.

Base.—Closed.

Margin.—Serrated.

Texture (both surfaces).—Leathery.

Venation pattern.—Palmate.

Venation color.—RHS 139C.

Petioles.—Length: 0.5 cm. Diameter: Less than 0.2 cm.

Color: RHS 139C. Texture: Rough.

Flower buds:

Shape.—Tall cylinder.

Length.—0.5 cm.

Diameter.—Less than 0.1 cm.

Color (at tight bud).—RHS 139C.

Inflorescence:

Blooming habit (flowering season): From early spring to late fall in central Europe

Lastingness of individual blooms on the plant.—2 weeks.

Fragrance.—Absent.

Inflorescence type.—Umbel.

Length of inflorescence.—4.0 cm.

Diameter of inflorescence.—7.0 cm.

Number of florets per inflorescence.—Greater than 25.

Peduncle.—Length: 3.5 cm. Diameter: 0.1 cm. Color: RHS 139C. Texture: Rough.

Florets:

Form.—Tall cylinder, opening at the end.

Length.—0.5 cm.

Diameter.—Less than 0.1 cm.

Floret depth.—1.0 cm.

Color.—Immature flower: Upper surface: RHS 83D with white eye of RHS N999D. Lower surface: RHS N82D with white eye of RHS N999D. Mature flower: Upper surface: RHS N82B with white eye of RHS N999D. Lower surface: RHS N82C with white eye of RHS N999D.

Flower throat color.—RHS 155D.

Corolla tube length.—2.0 cm.

Corolla tube color.—RHS 139C.

Corolla tube diameter.—<0.1 cm.

Petals.—Length of lobe: 0.5 cm. Diameter of lobe: 0.4 cm. Lobe shape: Reniform. Apex: Obtuse. Base: Closed. Margin: Entire.

Pedicels.—Not visible.

Calyx:

Arrangement.—Round.

Sepals.—Number per flower: 5. Shape: Funnel. Length: 2.0 cm. Diameter: Less than 0.1 cm. Color (Upper and lower surfaces): RHS 139C. Apex: Rounded. Base: Fused. Margin: Crenate. Texture: Smooth.

Reproductive organs:

Stamens.—Quantity: Greater than 5. Shape: Lanceolate. Color: RHS 139D. Filament length: Less than 0.1 cm. Filament diameter: Less than 0.1 cm.

Anther.—Shape: Fused. Color: RHS 139D. Length: Less than 0.1 cm. Diameter: Less than 0.1 cm.

Pollen.—Quantity: None observed.

Pistils.—Quantity: 1. Length: 0.2 cm. Diameter: Less than 0.1 cm. Stigma: Color: RHS 139D. Shape: Lanceolate. Length: Less than 0.1 cm. Diameter: Less than 0.1 cm. Style: Color: RHS 139D. Length: Less than 0.1 cm. Diameter: Less than 0.1 cm. Shape: Lanceolate.

Fruit and seed set: None observed.

Disease, insect resistance, and physiological stress: Resistant to powdery mildew.

COMPARISON WITH PARENTAL AND
COMMERCIAL VARIETIES

‘KLEVP12449’ differs from the female parent *Verbena* plant ‘1S112’ (unpatented), in that ‘KLEVP12449’ has larger flowers and is earlier flowering than ‘1S112’.

‘KLEVP12449’ differs from the male parent *Verbena* plant ‘V842’ (unpatented), in that ‘KLEVP12449’ has larger flowers with a white eye and better branching, whereas ‘V842’ has smaller flowers without an eye and weaker branching.

‘KLEVP12449’ differs from the commercial variety, ‘Vepita Lavender White’ (unpatented), in that ‘KLEVP12449’ has a more intense flower color, a better plant and branching habit, and is more vigorous than ‘Vepita Lavender White’.

I claim:

1. A new and distinct variety of *Verbena* plant named ‘KLEVP12449’ as shown and described herein.

* * * * *

