



US00PP24972P3

(12) **United States Plant Patent**
Mehlenbacher et al.(10) **Patent No.:** US PP24,972 P3
(45) **Date of Patent:** Oct. 21, 2014

- (54) **CORYLUS PLANT NAMED ‘YORK’**
- (50) Latin Name: *Corylus avellana*
Varietal Denomination: York
- (71) Applicant: State of Oregon acting by and through the State Board of Higher Education on behalf of Oregon State University, Corvallis, OR (US)
- (72) Inventors: **Shawn A. Mehlenbacher**, Corvallis, OR (US); **David C. Smith**, Corvallis, OR (US); **Rebecca L. McCluskey**, Corvallis, OR (US)
- (73) Assignee: **State Board of Higher Education on behalf of Oregon State University**, Corvallis, OR (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 50 days.
- (21) Appl. No.: **13/694,677**
- (22) Filed: **Dec. 24, 2012**

(65) **Prior Publication Data**

US 2014/0189913 P1 Jul. 3, 2014

- (51) **Int. Cl.**
A01H 5/00 (2006.01)
- (52) **U.S. Cl.**
USPC **Plt./152**
- (58) **Field of Classification Search**
USPC Plt./152
See application file for complete search history.

Primary Examiner — Anne Grunberg*(74) Attorney, Agent, or Firm* — Klarquist Sparkman, LLP**ABSTRACT**

A new and distinct cultivar of *Corylus* plant named ‘York’ characterized by globose plant habit and moderate vigor, green developing and fully expanded leaves during the spring and summer, resistance to eastern filbert blight caused by the fungus *Anisogramma anomala* (Peck) E. Müller, presence of random amplified polymorphic DNA markers 152-800 and AA12-850, expression of incompatibility alleles S₂ and S₂₁ in the styles, and DNA fingerprints at 10 of 23 microsatellite marker loci that differ from both parents OSU 479.027 and OSU 504.065, and from one parent at an additional 11 marker loci.

5 Drawing Sheets**1****ACKNOWLEDGMENT OF GOVERNMENT SUPPORT**

This invention was made with government support under Specific Cooperative Agreement No. 58-5358-4542 awarded by the United States Department of Agriculture. The government has certain rights in the invention.

Botanical denomination: *Corylus avellana*.

Variety designation: ‘York’.

BACKGROUND

The present Invention relates to a new and distinct cultivar of *Corylus* plant (hazelnut, filbert), botanically known as *Corylus avellana*, and hereinafter referred to by the name ‘York’. *Corylus avellana* is in the family Betulaceae.

The new *Corylus* resulted from a controlled cross of female parent OSU 479.027 (unpatented) and male parent OSU 504.065 (unpatented) made in 1997 by Shawn A. Mehlenbacher and David C. Smith. Hybrid seeds from the cross were harvested in August 1997, stratified, and seedlings grown in the greenhouse during the summer of 1998. From this cross, total of 183 seedling trees were planted in the field in Corvallis, Oreg., USA in October, 1998. ‘York’ was discovered and selected by the Inventors as a single plant within the progeny of the stated cross-pollination in a controlled environment in Corvallis, Oreg.

‘York’ was originally assigned the designation OSU 878.048, which indicates the row and tree location of the original seedling. OSU 479.027 is from a cross of OSU 231.111 (unpatented)×OSU 226.122 (unpatented). OSU 504.065 is from a cross of OSU 186.080 (unpatented)×VR 17-15 (unpatented). The pedigree of ‘York’ includes ‘Barce-

2

lona’ (unpatented) widely grown in Oregon, ‘Casina’ from Asturias, Spain (unpatented), ‘Tonda Gentile delle Langhe’ from Piemonte, northern Italy (unpatented), ‘Montebello’ from Sicily, Italy (unpatented), and ‘Tombul Ghiaghli’, a Turkish cultivar obtained from Greece (unpatented).

The new cultivar was asexually reproduced by rooted suckers annually for eight years (2003-2010) in Corvallis, Oreg. The new cultivar was also asexually propagated by whip grafting in Corvallis, Oreg. The unique features of this new *Corylus* are stable and reproduced true-to-type in successive generations of asexual reproduction.

SUMMARY

15 The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘York’. These characteristics in combination distinguish ‘York’ as a new and distinct cultivar:

1. Globose plant habit and moderate vigor.
2. Green developing and fully expanded leaves during the spring and summer.
3. Resistance to eastern filbert blight caused by the fungus *Anisogramma anomala* (Peck) E. Müller.
4. Presence of random amplified polymorphic DNA markers 152-800 and AA12-850 in DNA of ‘York’ amplified by the polymerase chain reaction. These two markers are linked to a dominant allele for resistance to eastern filbert blight from the cultivar Gasaway (unpatented).
5. Expression of incompatibility alleles S₂ and S₂₁ in the styles.
6. DNA fingerprints at 10 of 23 microsatellite marker loci differ from both parents OSU 479.027 and OSU 504.065, and from one parent at an additional 11 marker

loci. The microsatellite primers are shown in Table 1, and allele sizes are shown in Table 2. DNA fingerprints of standard cultivars 'Barcelona', 'Tonda Gentile delle Langhe' and 'Extra Ghiaglia', and 'Gasaway', the source of eastern filbert blight resistance, are also shown in attached Table 2.

In comparisons in two replicated trials conducted in Corvallis, Oreg., plants of the new *Corylus* differed from plants of the *Corylus avellana* cultivar 'Barcelona' (unpatented), and other cultivars and selections of *Corylus avellana* known to the Inventors primarily in nut size, nut shape, kernel percentage (ratio of kernel weight to nut weight), frequency of blank nuts (nuts lacking kernels), time of pollen shed, time of nut maturity, length of the husk or involucre, and plant size.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Foliage colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Corylus*.

FIG. 1 shows a tree of the new cultivar 'York' growing in a field in August, in Corvallis, Oreg.

FIG. 2 shows the tree of the new cultivar 'York' growing in a field in January, in Corvallis, Oreg.

FIG. 3 shows typical nuts, raw kernels, and blanched kernels of 'York' hazelnut compared to those of 'Jefferson' hazelnut.

FIG. 4 shows typical nuts, raw kernels, and blanched kernels of 'York' hazelnut compared to those of 'Lewis' hazelnut.

FIG. 5 shows the typical nuts, raw kernels, and blanched kernels of 'York' hazelnut compared to those of 'Barcelona' hazelnut and other hazelnut cultivars.

DETAILED BOTANICAL DESCRIPTION

The cultivar York has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype. The aforementioned photographs and following observations and measurements describe plants grown in Corvallis, Oreg. under commercial practice outdoors in the field during the fall, winter and spring. Plants used for the photographs and description were propagated by tie-off layerage and growing on their own roots, and about seven years old. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1966 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Corylus avellana* cultivar York.

Parentage:

Female, or seed, parent.—*Corylus avellana* selection OSU 479.027 (unpatented).

Male, or pollen, parent.—*Corylus avellana* selection OSU 504.065 (unpatented).

Propagation (type rooted suckers):

Time to initiate roots.—About 30 days at 20 degrees C.

Time to produce a rooted young plant.—About six months at 22 degrees C.

Root description.—Fine to thick; freely branching; creamy white in color.

Propagation (type whip grafting):

Time to budbreak on the scions.—About 14 days at 25° C.

Time to produce a grafted plant.—About six months at 25° C.

Plant description:

General appearance.—Perennial shrub.

Size.—Plant height is about 5 meters; plant diameter or spread is about 5 meters.

Form.—Spreading plant habit.

Growth and branching habit.—freely branching; about 15 lateral branches develop per plant. Pinching, i.e., removal of the terminal apices, enhances branching with lateral branches potentially forming at every node.

Vigor.—Moderate vigor growth habit.

Roots.—Fine to thick; freely branching; creamy white in color.

Lenticels.—11 circular within 1 square centimeter (counted on dormant scions).

Lateral branch description:

Length.—About 38 cm.

Diameter.—About 6 mm.

Internode length.—About 3.2 cm.

Texture.—Smooth, glabrous.

Strength.—Strong.

Color.—Immature — 152B, mature — 152B.

Foliage description:

Arrangement.—Alternate, simple

Length.—About 10.0 cm.

Width.—About 9.3 cm.

Shape.—Ablong to ovate.

Apex.—Obtuse to acute.

Base.—Cordate.

Margin.—Serrate.

Texture.—Upper and lower surfaces — slightly pubescent.

Venation pattern.—Pinnate.

Leaf bud shape.—Globular.

Time of leaf bud burst.—Midseason, 11 days after 'Barcelona'.

Color.—Developing foliage: upper surface 146B, lower surfaces: 146C. Fully expanded foliage, upper surface: Spring and summer, 146A; late summer and fall, 146A. Fully expanded foliage, lower surface: Spring and summer, 146C; late summer and fall, 146C. Venation, upper surface: Spring and summer, 146A; late summer and fall, 146A. Venation, lower surface: Spring and summer, 148D; late summer and fall, 148D. Leaf bud, 177C.

Petiole description:

Length.—About 2.7 cm.

Diameter.—About 1.8 mm.

Texture.—Upper and lower surfaces: pubescent.

Color.—Upper surface: Spring and summer, 139D; late summer and fall, 139D. lower surface: Spring and summer, 139D; late summer and fall, 139D.

Flower description:

Male inflorescences.—Catkins, color prior to elongation 194C.

Female inflorescence.—Style color 047B.

Stigma coloration.—047B.

Time of female flowering.—Midseason, 14 days after 'Barcelona'.

Time of pollen shed.—Midseason, around the same time as 'Daviana' (unpatented).

Involure description:

Involucre constriction.—Absent.

Involucre length.—25% longer than nuts.

Strength of serration of indentation.—Moderate.

Pubescence.—Little

Thickness of callus at base.—Moderate callus at base similar to 'Barcelona'.

Description of jointing of bracts.—Involucre slit to the base on one side. Nuts fall free of the involucre at maturity (95% fall free).

Nut description:

Length.—About 18.0 mm

Width.—About 19.7 mm.

Depth.—About 17.0 mm.

Nut shape.—Round.

Nut shape index $[(width+depth)/2 * length]$.—1.02.

Nut compression index (width/depth).—1.16.

Nut shell color.—164A.

Nut weight.—About 2.73 grams to 2.85 grams.

Predominant number of fruits per cluster.—Average 2 nuts per cluster.

Stripes on shell.—About 24, color slightly darker than 164A.

Fruit apex.—Slight (not prominent).

Size of the fruit pistil scar.—Very small (0.5 mm×1 mm).

Nut curvature of the basal scar.—Flat (plane).

Frequency of blank nuts.—9.7%.

Time of nut maturity.—About 3 days earlier than 'Barcelona'.

Husk length.—About 25% longer than the nuts.

Kernel weight.—About 1.23 grams to 1.32 grams.

Kernel percentage (kernel weight/nut weight).—About 46%. Kernel shape — globular.

Kernel cross section shape.—Circular.

Kernel base shape.—Flat.

Lateral grooves.—None.

Disease/pest resistance: Plants of the new *Corylus* are highly resistant to eastern filbert blight caused by the fungus *Anisogramma anomala* (Peck) E. Müller. Plants of the new *Corylus* are highly resistant to bud mites (*Phytoptus avellanae* Nal.), while plants of 'Tonda Gentile delle Langhe' are highly susceptible, and plants of 'Barcelona' are highly resistant.

Temperature tolerance: Tolerates temperatures from -10 to 38° C. in the field in Corvallis, Oreg.

TABLE 1

Primers and annealing temperatures for the 24 microsatellite marker loci used to fingerprint 'York' and other hazelnut cultivars.

Locus	Repeat motif	Size	T _a	n	He	Ho
A613	(TC) ₁₃ (CA) ₁₂	149-177	60	14	0.85	0.85
A614	(TC) ₁₇ (CA) ₁₀	125-156	60	14	0.85	0.85
	NNN(CA) ₆					
A616	(AC) ₁₁	136-162	60	13	0.85	0.85
A640	(CT) ₁₅	354-378	67	11	0.80	0.73
	(CA) ₁₃					
B107	(CT) ₁₄	112-151	55	14	0.85	0.80
B617	(GA) ₁₅	280-298	60	9	0.80	0.78
B619	(TC) ₂₁	146-180	60	14	0.88	0.88
B634	(AG) ₁₅	218-238	60	9	0.76	0.76

TABLE 1-continued

Primers and annealing temperatures for the 24 microsatellite marker loci used to fingerprint 'York' and other hazelnut cultivars.						
Locus	Primer sequence	Annealing temperature (°C)	n	He	Ho	Primers 5'-3'
B657	(AG) ₁₅	210-228	60	8	0.84	0.98
B671	(AG) ₆ NN	221-249	60	13	0.86	0.88
	(GA) ₁₇					
B709	(GA) ₂₁	219-233	60	8	0.74	0.76
B733	(TC) ₁₅	161-183	60	8	0.68	0.68
B741	(GT) ₅ (GA) ₁₂	176-194	60	10	0.77	0.78
B749	(TC) ₁₂	200-210	60	6	0.60	0.64
B751	(GA) ₁₅	141-153	60	7	0.80	0.80
B774	(AG) ₁₅	195-213	60	8	0.80	0.80
B776	(GA) ₁₇	134-148	60	7	0.71	0.60
B795	(TC) ₈ Ns(CT) ₇	296-332	60	12	0.76	0.74
	Ns(CT) ₁₀					
	Ns(TC) ₅					
C115	(TAA) ₅	167-226	60	14	0.80	0.80
	(GAA) ₁₂					
KG809	(AGG) ₆	333-345	55	5	0.66	0.64
KG811	(GA) ₁₇	240-278	58	12	0.83	0.82
KG827	(CT) ₁₃ AA	264-282	67	9	0.78	0.84
	(CA) ₇					
KG830	(CT) ₁₄	279-311	67	9	0.79	0.78
	GTATT					
	(CA) ₈					
Soman-G	(AAT) ₅				54	3
					0.60	0.98
Locus	Primer sequence	Annealing temperature (°C)	n	He	Ho	Primers 5'-3'
A613		0.85	0.00	11		Ned-CACACGCCTT
						GTCACTCTTT
						(SEQ ID NO: 1)
30 A614		0.84	0.00	6		Hex-TGGCAGAGCT
						TTGTCAGCTT
						(SEQ ID NO: 3)
A616		0.83	0.00	8		Fam-CACTCATACC
						GCAAACCTCCA
						(SEQ ID NO: 5)
35 A640		0.7	0.04	10		F-TGCCCTTGCA
						GTTAGTCATC
						AAATGTAGG
						(SEQ ID NO: 7)
B107		0.83	0.02	10		Ned-GTAGGTGCAC
						TTGATGTGCTT
						TAC (SEQ ID NO: 9)
40 B617		0.78	0.01	8		Fam-TCCGTGTTGA
						GTATGGACGA
						(SEQ ID NO: 11)
B619		0.7	0.00	3		Fam-AGTCGGCTCC
						CCTTTTCTC
						(SEQ ID NO: 13)
45 B634		0.73	0.00	4		Hex-CCTGCATCCA
						GGACTCATTA
						60 (SEQ ID NO: 15)
B657		0.82	-0.08	11		Ned-GAGAGTGCCTG
						CTTCCTCTGG
						(SEQ ID NO: 17)
50 B671		0.84	-0.01	9		Hex-TTGCCAGTGC
						ATACTCTGAT G
						(SEQ ID NO: 19)
B709		0.70	-0.01	5		Ned-CCAAGCACGA
						ATGAACCTCAA
						(SEQ ID NO: 21)
55 B733		0.63	0.00	7.2		Ned-CACCCCTTTCA
						CCACCTCAT
						(SEQ ID NO: 23)
B741		0.74	0.00	5		Fam-GTTCACAGGC
						TGTTGGTTT
						(SEQ ID NO: 25)
B749		0.51	-0.03	1		Hex-GGCTGACAAC
						ACAGCAGAAA
						(SEQ ID NO: 27)
B751		0.77	0.01	7.2		Fam-AGCTGGTTCTT
						CGACATTCC
						(SEQ ID NO: 29)
60						
65						

TABLE 1-continued

Primers and annealing temperatures for the 24 microsatellite marker loci used to fingerprint 'York' and other hazelnut cultivars.					
B774	0.77	0.01	5	Ned-GTTTGCAG CTCATGGCA (SEQ ID NO: 31)	5
B776	0.67	0.07	6	Fam-TGTATGACA CACGGAGAGA GAGA (SEQ ID NO: 33)	10
B795	0.74	0.01	NA	Fam-GACCCACAAA CAATAACCTA TCTC (SEQ ID NO: 35)	B774
C115	0.77	0.00	4	Fam-ATTTCCGCA GATAATACAGG (SEQ ID NO: 37)	15
KG809	0.60	0.01	4	Hex-AGGCATCAGT TCATCCAA (SEQ ID NO: 39)	KG809
KG811	0.81	0.01	2	Ned-AAGCGGCAC TCGCTCAC (SEQ ID NO: 41)	20
KG827	0.75	-0.04	9	Fam-AGAACTCCGA CTAATAATCC TAACCCTTGC (SEQ ID NO: 43)	KG811
KG830	0.76	0.00	9	Ned-TGGAGGAAGT TTGAATGGT AGTAGAGGA (SEQ ID NO: 45)	25
Soman-G	0.51	-0.27	NA	Hex-TGGCGTTGCA ACATATTCTC (SEQ ID NO: 47)	KG830

Locus	Primers 5'-3'	Reference
A613	R-CCCCTTCACAT GTTTGCTT (SEQ ID NO: 2)	Gurcan et al. 2010
A614	R-GCAGTGGAGGA TTGCTGACT (SEQ ID NO: 4)	Gurcan et al. 2010
A616	R-ATGGCTTTGCT TCGTTTG (SEQ ID NO: 6)	Gurcan et al. 2010
A640	Fam-CGCCATATAATT GGGATGCTTGT G (SEQ ID NO: 8)	Gurcan et al. 2010
B107	R-AACACCATTG AGTCTTCAAAG C (SEQ ID NO: 10)	Boccacci et al. 2005; Gokirmak et al. 2009
B617	R-TGTTTTGGTGG AGCGATG (SEQ ID NO: 12)	Gurcan et al. 2010
B619	R-GCGATCTGACCT CATTTTG (SEQ ID NO: 14)	Gurcan et al. 2010
B634	R-GTGCAGAGGTTG CACTCAA (SEQ ID NO: 16)	Gurcan et al. 2010
B657	R-AGCCTCACCTCC AACGAAC (SEQ ID NO: 18)	Gurcan et al. 2010
B671	R-ACCAGCTCTGGG CTTAACAC (SEQ ID NO: 20)	Gurcan et al. 2010
B709	R-GCGGGTTCTCGT TGTACACT (SEQ ID NO: 22)	Gurcan et al. 2010
B733	R-CATCCCCTGTTG GAGTTTC (SEQ ID NO: 24)	Gurcan et al. 2010
B741	R-CGTGTTGCTCAT GTGTTGTG (SEQ ID NO: 26)	Gurcan et al. 2010

TABLE 1-continued

Primers and annealing temperatures for the 24 microsatellite marker loci used to fingerprint 'York' and other hazelnut cultivars.					
B749	R-TCGGCTAGGGTT AGGGTTTT (SEQ ID NO: 28)	Gurcan et al. 2010			
B751	R-AAACTCAAATA AAACCCCTGCTC (SEQ ID NO: 30)	Gurcan et al. 2010			
B774	R-TGTGTGTGGTCT GTAGGCACT (SEQ ID NO: 32)	Gurcan et al. 2010			
B776	R-TGAGGGAAAGA GGTTTGATG (SEQ ID NO: 34)	Gurcan et al. 2010			
B795	R-TGGGCATCATCC AGGTCTA (SEQ ID NO: 36)	Gurcan et al. 2010			
C115	GTTCAGATCT GCCTCCATATAA T (SEQ ID NO: 38)	Bassil et al. 2005b, Gokirmak et al. 2009			
KG809	F-GGAAGGTGAGA GAAATCAAGT (SEQ ID NO: 40)	Gurcan and Mehlenbacher 2010			
KG811	F-GAACAACTGAA GACAGCAAAG (SEQ ID NO: 42)	Gurcan and Mehlenbacher 2010			
KG827	GAGGGAGCAAQ TCAAAGTTGAGA AGAAA (SEQ ID NO: 44)	Gurcan and Mehlenbacher 2010			
KG830	AAAGCAACTCAT AGCTGAAGTCCA ATCA (SEQ ID NO: 46)	Gurcan and Mehlenbacher 2010			
Soman-G	R-GCCATCTTCTAGA AAGTCGATACAG (SEQ ID NO: 48)	unpublished			

35 Primer fluorescent tags are FAM, HEX, and NED.
Ta: annealing temperature (° C.)
N: number of alleles
He: expected heterozygosity
Ho: observed heterozygosity
40 PIC: polymorphism information content
r: estimated null allele frequency
LG: linkage group

TABLE 2

Allele sizes in 'York', its parents, and four other hazelnut cultivars at 24 microsatellite loci.					
Tag	Locus	'York'	479.027	504.065	'Tonda Gentile delle Langhe'
NED	A613	157/177	177/177	157/159	151/157
HEX	A614	125/158	125/132	148/158	125/135
FAM	A616	142/150	142/150	150/150	148/150
FAM	A640	362/374	354/374	354/362	354/368
NED	B107	122/134	134/146	122/134	134/152
FAM	B617	286/290	286/290	286/286	286/296
FAM	B619	156/164	156/164	156/164	148/164
HEX	B634	226/234	226/226	232/234	226/226
NED	B657	220/222	222/226	220/224	218/226
HEX	B671	241/247	227/247	241/241	237/241
NED	B709	227/231	227/231	227/227	227/227
NED	B733	171/179	171/179	173/179	171/173
FAM	B741	177/186	177/186	177/184	177/184
HEX	B749	208/208	206/208	206/208	206/208
FAM	B751	151/153	151/153	143/151	149/153
NED	B774	203/209	209/211	203/207	203/211
FAM	B776	137/150	137/137	137/150	137/137
FAM	B795	330/330	296/330	330/330	312/330
FAM	C115	197/197	194/197	194/197	173/173

TABLE 2-continued

Allele sizes in 'York', its parents, and four other hazelnut cultivars at 24 microsatellite loci.					
HEX	KG809	336/345	336/339	339/345	336/339
NED	KG811	254/254	254/254	254/264	254/264
FAM	KG827	266/270	270/282	266/266	266/268
NED	KG830	295/295	295/295	295/295	291/295
HEX	SMNG	196/200	196/200	196/200	196/200

	Tag	Locus	'Barcelona'	'Extra Ghiaghli'	'Gasaway'
NED	A613	151/159	167/169	159/161	
HEX	A614	125/131	125/150	143/158	
FAM	A616	142/150	150/158	148/148	
FAM	A640	354/374	374/374	362/368	
NED	B107	112/134	116/116	122/128	
FAM	B617	286/290	294/296	292/296	
FAM	B619	156/170	164/174	170/174	
HEX	B634	226/226	226/226	220/232	
NED	B657	218/222	210/222	224/228	
HEX	B671	223/227	227/247	235/247	
NED	B709	225/233	225/227	227/227	
NED	B733	171/173	171/171	173/173	
FAM	B741	177/186	177/184	186/188	
HEX	B749	208/208	208/208	206/208	
FAM	B751	143/153	143/147	143/143	
NED	B774	203/207	195/203	203/209	
FAM	B776	135/137	135/137	146/150	
FAM	B795	330/330	296/310	314/316	
FAM	C115	173/194	182/194	215/218	
HEX	KG809	336/336	336/339	336/345	
NED	KG811	258/264	240/242	254/258	
FAM	KG827	280/282	276/282	270/280	
NED	KG830	291/295	291/295	291/305	
HEX	SMNG	196/200	196/200	196/196	

REFERENCES:

- 5 Bassil N. V., Botta R., Mehlenbacher S. A. 2005a. Microsatellite markers in hazelnut: Isolation, characterization and cross-species amplification. *J. Amer. Soc. Hort. Sci.* 130: 543-549.
- 10 Bassil N. V., Botta R., Mehlenbacher S. A. 2005b. Additional microsatellite markers of the European hazelnut. *Acta Hort.* 686:105-110.
- Boccacci P., Akkak A., Bassil N. V., Mehlenbacher S. A., Botta R. 2005. Characterization and evaluation of microsatellite loci in European hazelnut (*C. avellana*) and their transferability to other *Corylus* species. *Molec. Ecol. Notes* 5:934-937.
- 15 Boccacci P., Akkak, A. and Botta, R. 2006. DNA typing and genetic relations among European hazelnut (*Corylus avellana* L.) cultivars using microsatellite markers. *Genome* 49:598-611.
- Gökirmak T , Mehlenbacher S. A., Bassil N. V. 2009. Characterization of European hazelnut (*Corylus avellana*) cultivars using SSR markers. *Genetic Resources and Crop Evolution* 56:147-172.
- 20 Gürcan, K., S. A. Mehlenbacher and V. Erdogan. 2010a. Genetic diversity in hazelnut cultivars from Black Sea countries assessed using SSR markers. *Plant Breeding* (available on-line doi:10.1111/j.1439-0523.2009.01753.x).
- Gürcan, K., S. A. Mehlenbacher, N. V. Bassil, P. Boccacci, A. Akkak and R. Botta. 2010b. New microsatellite markers for *Corylus avellana* from enriched libraries. *Tree Genetics and Genomes* (available on-line as DOI 10.1007/s11295-010-0269-y).
- 25 Gürcan, K. and S. A. Mehlenbacher. 2010. Development of microsatellite marker loci for European hazelnut (*Corylus avellana* L.) from ISSR fragments. *Molecular Breeding* (available on-line).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 48

<210> SEQ ID NO 1

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 1

cacacgcctt gtcactcttt

20

<210> SEQ ID NO 2

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 2

cccctttcac atgtttgctt

20

<210> SEQ ID NO 3

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

- continued

<400> SEQUENCE: 3

tggcagagct ttgtcagctt

20

<210> SEQ ID NO 4

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 4

gcagtggagg attgctgact

20

<210> SEQ ID NO 5

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 5

cactcataacc gcaaactcca

20

<210> SEQ ID NO 6

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 6

atggctttg cttcgaaaa

20

<210> SEQ ID NO 7

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 7

tgcctctgca gtttagtcata aaatgttagg

29

<210> SEQ ID NO 8

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 8

cgcctataaa ttggatgtt tttttt

25

<210> SEQ ID NO 9

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 9

gttaggtgcac ttgatgtt ttac

24

- continued

```

<210> SEQ ID NO 10
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 10
aacaccatat tgagtctttc aaagc 25

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 11
tccgtgttga gtatggacga 20

<210> SEQ ID NO 12
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 12
tgtttttgtt ggagcgatg 19

<210> SEQ ID NO 13
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 13
agtcggctcc ccttttctc 19

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 14
gcgatctgac ctcattttg 20

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 15
cctgcatcca ggactcatta 20

<210> SEQ ID NO 16
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

```

- continued

<400> SEQUENCE: 16

gtgcagagg tgcactcaa

20

<210> SEQ ID NO 17

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 17

gagagtgcgt cttcctctgg

20

<210> SEQ ID NO 18

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 18

agcctcaccc ccaacgaac

19

<210> SEQ ID NO 19

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 19

ttgccagtgc atactctgat g

21

<210> SEQ ID NO 20

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 20

accagctctg ggcttaacac

20

<210> SEQ ID NO 21

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 21

ccaaggcacga atgaactcaa

20

<210> SEQ ID NO 22

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 22

gcgggttctc gttgtacact

20

- continued

```

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 23
caccctcttc accacccatc                                     20

<210> SEQ ID NO 24
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 24
catccccgt tggagtttcc                                     20

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 25
gttcacaggc tgggggtt                                     20

<210> SEQ ID NO 26
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 26
cggttgctc atgtgttgtg                                     20

<210> SEQ ID NO 27
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 27
ggctgacaac acagcagaaa                                     20

<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 28
tcggctaggg tttagggttt                                     20

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide

```

- continued

<400> SEQUENCE: 29

agctggttct tcgacattcc

20

<210> SEQ ID NO 30

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 30

aaaactcaaat aaaacccctg ctc

23

<210> SEQ ID NO 31

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 31

gttttgcgag ctcattgtca

20

<210> SEQ ID NO 32

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 32

tgtgtgtggc tctgttaggcac t

21

<210> SEQ ID NO 33

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 33

tgtatgtaca cacggagaga gaga

24

<210> SEQ ID NO 34

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 34

tgaggggaag aggtttgatg

20

<210> SEQ ID NO 35

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 35

gacccacaaa caataaaccta tctc

24

- continued

<210> SEQ ID NO 36
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<400> SEQUENCE: 36

tgggcatcat ccaggtcta

19

<210> SEQ ID NO 37
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<400> SEQUENCE: 37

atttccgca gataatacag g

21

<210> SEQ ID NO 38
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<400> SEQUENCE: 38

gtttccagat ctgcctccat ataat

25

<210> SEQ ID NO 39
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<400> SEQUENCE: 39

aggcatcagt tcatccaa

18

<210> SEQ ID NO 40
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<400> SEQUENCE: 40

ggaaggtgag agaaatcaag t

21

<210> SEQ ID NO 41
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<400> SEQUENCE: 41

aaggcggcac tcgctcac

18

<210> SEQ ID NO 42
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<400> SEQUENCE: 42

- continued

<400> SEQUENCE: 42

gaacacaactga agacagcaaa g

21

<210> SEQ ID NO 43

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 43

agaactccga ctaataatcc taacccttgc

30

<210> SEQ ID NO 44

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 44

gaggggagcaa gtcaaaggta agaagaaaa

28

<210> SEQ ID NO 45

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 45

tggaggaagt tttgaatggta agtagagga

29

<210> SEQ ID NO 46

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 46

aaagcaactc atagctgaag tccaatca

28

<210> SEQ ID NO 47

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 47

tggcggttgca acatatttc

20

<210> SEQ ID NO 48

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<400> SEQUENCE: 48

gccatcttta gaaaatcgat tacag

25

We claim:

1. A new and distinct cultivar of *Corylus* plant named 'York', as illustrated and described.

* * * * *



FIG. 1



FIG. 2




FIG. 3




FIG. 4




FIG. 5