



US00PP25150P2

(12) **United States Plant Patent**
Trees

(10) **Patent No.:** **US PP25,150 P2**
(45) **Date of Patent:** **Dec. 2, 2014**

(54) **MIMULUS PLANT NAMED ‘BOSBUREM’**

(50) Latin Name: *Mimulus aurantiacus*
Varietal Denomination: **Bosburem**

(71) Applicant: **Ball Horticultural Company**, West
Chicago, IL (US)

(72) Inventor: **Scott Trees**, Arroyo Grande, CA (US)

(73) Assignee: **Ball Horticultural Company**, West
Chicago, IL (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 47 days.

(21) Appl. No.: **13/986,610**

(22) Filed: **May 17, 2013**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./454**

(58) **Field of Classification Search**
USPC Plt./454
See application file for complete search history.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — Audrey Charles

(57) **ABSTRACT**

A new and distinct cultivar of *Mimulus* plant named ‘Bosbu-
rem’, characterized by its dark yellow-colored flowers, dark
green-colored foliage, low to moderate vigor, and compact-
upright growth habit, is disclosed.

1 Drawing Sheet

1

Latin name of genus and species of plant claimed: *Mimulus aurantiacus*.

Variety denomination: ‘Bosburem’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Mimulus* plant botanically known as *Mimulus aurantiacus* and hereinafter referred to by the cultivar name ‘Bosburem’.

The new cultivar originated in a controlled breeding program in Guadalupe, Calif. during May 2009. The objective of the breeding program was the development of *Mimulus* cultivars with large flowers, attractive flower coloration, and a compact-upright growth habit.

The new *Mimulus* cultivar was the result of a self-pollination of the proprietary *Mimulus aurantiacus* breeding selection coded nct-8518-e, not patented, characterized by its dark yellow-colored flowers, dark green-colored foliage, low to moderate vigor, and compact-upright growth habit. The new cultivar was discovered and selected as a single flowering plant within the progeny of the above stated self-pollination during July 2010 in a controlled environment in Guadalupe, Calif.

Asexual reproduction of the new cultivar by terminal stem cuttings since July 2010 in Guadalupe, Calif. has demonstrated that the new cultivar reproduces true to type with all of the characteristics, as herein described, firmly fixed and retained through successive generations of such asexual propagation.

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have been repeatedly observed and can be used to distinguish ‘Bosburem’ as a new and distinct cultivar of *Mimulus* plant:

1. Dark yellow-colored flowers;
2. Dark green-colored foliage;
3. Low to moderate vigor; and
4. Compact-upright growth habit.

2

Plants of the new cultivar differ from plants of the parent primarily in growth vigor and habit.

Of the many commercially available *Mimulus* cultivars, the most similar in comparison to the new cultivar is ‘Jelly Bean Yellow’, U.S. Plant Pat. No. 11,973. However, in side-by-side comparison, plants of the new cultivar differ from plants of ‘Jelly Bean Yellow’ in at least the following characteristics:

1. Plants of the new cultivar are more compact than plants of ‘Jelly Bean Yellow’; and
2. Plants of the new cultivar have a larger corolla than plants of ‘Jelly Bean Yellow’.

In addition, the new cultivar is similar to ‘Georgie Yellow’, U.S. Plant Pat. No. 22,945. However, in comparison, plants of the new cultivar differ from plants of ‘Georgie Yellow’ in at least the following characteristics:

1. Plants of the new cultivar are more compact than plants of ‘Georgie Yellow’;
2. Plants of the new cultivar have a larger corolla than plants of ‘Georgie Yellow’; and
3. Plants of the new cultivar have a darker fully-opened flower color than plants of ‘Georgie Yellow’.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it is reasonably possible to make the same in color illustrations of this type, typical flower and foliage characteristics of the new cultivar. Colors in the photographs differ slightly from the color values cited in the detailed description, which accurately describes the colors of ‘Bosburem’. The plants were grown in 4.5-inch pots for 13 weeks in a greenhouse at West Chicago, Ill. Plants were given one pinch one week after transplant.

FIG. 1 illustrates a side view of the overall growth and flowering habit of ‘Bosburem’.

FIG. 2 illustrates a close-up view of an individual flower of ‘Bosburem’.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible

that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2007 edition, except where general color terms of ordinary significance are used. The color values were determined in March 2013 under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe plants produced from cuttings from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown in West Chicago, Ill. in 4.5-inch pots for 13 weeks utilizing a soilless growth medium. Plants were given one pinch one week after transplant. Greenhouse temperatures were maintained at approximately 70° F. to 77° F. (21° C. to 25° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Greenhouse light levels of 2,500 footcandles to 6,000 footcandles were maintained during the day. Measurements and numerical values represent averages of typical plants.

Botanical classification: *Mimulus aurantiacus* cultivar Bosburem.

Parentage:

Female and male parent.—Proprietary *Mimulus aurantiacus* breeding selection coded nct-8518-e, not patented.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 10 to 14 days.

Time to produce a rooted cutting.—Approximately 21 to 25 days.

Root description.—Fibrous.

Rooting habit.—Freely branching.

Plant description:

Commercial crop time.—Approximately 10 to 12 weeks from a rooted cutting to finish in a 10 cm pot.

Growth habit and general appearance.—Low to moderate growth vigor, and compact-upright growth habit.

Size.—Height from soil level to top of plant plane: Approximately 24.0 cm. Width: Approximately 20.5 cm.

Branching habit.—Freely branching, pinching enhances basal branching. Quantity of main branches per plant: Approximately 4.

Branch.—Strength: Strong, somewhat flexible. Length: Approximately 19.5 cm. Diameter: Approximately 3.0 mm. Length of central internode: Approximately 2.1 cm. Texture: Glabrous, viscid, densely glandular pubescent. Color of young stems: 145B. Color of mature stems: 199B.

Foliage description:

General description.—Quantity of leaves per main branch: Approximately 20. Fragrance: Slight. Form: Simple. Arrangement: Opposite.

Leaves.—Aspect: Perpendicular to stem, slightly reflexed. Shape: Elliptic. Margin: Serrate. Apex: Rounded acute. Base: Attenuate. Venation pattern: Pinnate. Length of mature leaf: Approximately 4.9 cm. Width of mature leaf: Approximately 1.5 cm. Texture of upper surface: Glabrous, glossy. Texture of lower surface: Glabrous, viscid, densely glandular pubescent. Color of upper surface of young and

mature foliage: 137A with indistinguishable venation. Color of lower surface of young and mature foliage: Closest to 138B with midvein of 147C.

Petiole.—Length: Approximately 2.0 mm. Diameter: Approximately 3.0 mm. Texture: Glabrous, viscid, densely glandular pubescent. Color: 144D.

Flowering description:

Flowering habit.—‘Bosburem’ is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

Lastingness of individual flower on the plant.—Approximately 2 to 3 weeks.

Flower description:

General description.—Type: Single, salverform, zygomorphic, axillary, facing outward. Quantity per plant: Approximately 9. Fragrance: None detected.

Bud.—Rate of opening: Generally takes 2 to 3 days for bud to progress from first color to fully open flower. Quantity per plant: Approximately 2.

Bud just before opening.—Shape: Elongate. Length: Approximately 2.9 cm. Diameter: Approximately 9.0 mm. Texture: Glabrous, slightly viscid, glandular pubescent. Color: Sepals of 144B with folded midvein area of 137A, lobes of 15C.

Corolla.—Vertical axis: Approximately 4.2 cm. Horizontal axis: Approximately 4.0 cm.

Lobes.—Quantity: 5, fused to form a tube. Shape: Fan-shaped, upper two lobes reflexed. Margin: Irregularly sinuate and ruffled. Apex: Obtuse to truncate. Length from tube of upper two lobes: Approximately 2.3 cm. Length from tube of lower lobes: Approximately 1.6 cm. Width of upper two lobes: Approximately 2.1 cm. Width of lower lobes: Approximately 1.6 cm. Texture of upper and lower surfaces: Glabrous. Color of upper surface when first open: 13A. Color of lower surface when first and fully open: 13C. Color of upper surface when fully open: 13B.

Corolla tube.—Length: Approximately 3.8 cm. Diameter at distal end: Approximately 8.0 mm. Diameter at proximal end: Approximately 2.0 mm. Texture of inner and outer surfaces: Glabrous. Color of inner surface: 4D with two nectar glides at base of lower lobe of 13B. Color of outer surface: 13D.

Sepals.—Quantity per flower: 5, fused. Shape: Elliptic, folded at midveins. Margin: Entire. Apex: Acute. Length: Approximately 2.7 cm with one upper sepal of approximately 3.0 cm. Width: Approximately 4.0 mm. Texture of upper and lower surfaces: Glabrous, slightly viscid, glandular pubescent. Color of upper and lower surfaces: 144B with folded midvein area of 137A.

Peduncle.—Strength: Strong, flexible. Aspect: Acute angle to stem. Length: Approximately 4.0 mm. Diameter: Approximately 1.0 mm. Texture: Densely glandular pubescent. Color: 144B.

Reproductive organs.—Androecium: Stamen quantity: 4, didynamous, basifixed. Stamen length of longer pair: Approximately 3.2 cm. Stamen length of shorter pair: Approximately 2.8 cm. Filament length of fixed portion: Approximately 1.2 cm. Filament color: 1C. Anther shape: Bilobed, ovoid. Anther length: Approximately 2.0 mm. Anther color: 25B. Pollen amount: Moderate. Pollen color: 4B. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approxi-

mately 3.6 cm. Stigma shape: Flattened, bilobed.
Stigma length: Approximately 2.0 mm. Stigma color:
155D. Style length: Approximately 3.0 cm. Style
color: 155A. Ovary length: Approximately 4.0 mm.
Ovary color: 144A.
Seed and fruit production: Neither seed nor fruit production
has been observed.

Disease and pest resistance: Resistance to pathogens and
pests common to *Mimulus* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Mimulus* plant named
'Bosburem', substantially as herein shown and described.

* * * * *




FIG. 1



FIG. 2