

US00PP25552P2

(12) United States Plant Patent Dummen

(10) Patent No.:

US PP25,552 P2

(45) **Date of Patent:**

May 12, 2015

PETUNIA PLANT NAMED 'DUEPOTMOCA'

Latin Name: *Petunia×hybrida* Varietal Denomination: **Duepotmoca**

Applicant: **Tobias Dummen**, Rheinberg (DE)

Tobias Dummen, Rheinberg (DE) Inventor:

Assignee: **Dümmen Group B.V.**, DeLier (NL)

Subject to any disclaimer, the term of this Notice: patent is extended or adjusted under 35

U.S.C. 154(b) by 101 days.

Appl. No.: 13/987,110

Jul. 1, 2013 (22)Filed:

(51)Int. Cl. A01H 5/00

(2006.01)

(52)U.S. Cl.

Field of Classification Search (58)

See application file for complete search history.

Primary Examiner — Kent L Bell

(74) Attorney, Agent, or Firm — C. A. Whealy

ABSTRACT (57)

A new and distinct cultivar of *Petunia* plant named 'Duepotmoca', characterized by its compact, upright and mounding plant habit; moderately vigorous growth habit; freely branching habit; early and freely flowering habit; pink and creamcolored flowers with red purple-colored venation which gives a brownish appearance; and good garden performance.

1 Drawing Sheet

Botanical designation: *Petunia*×*hybrida*. Cultivar denomination: 'DUEPOTMOCA'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Petunia* plant, botanically known as *Petunia*×*hybrida* and hereinafter referred to by the name 'Duepotmoca'.

The new *Petunia* plant is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. 10 The objective of the breeding program is to create new compact Petunia plants with numerous unique and attractive flowers.

The new *Petunia* plant originated from a cross-pollination made by the Inventor in July, 2010 in Rheinberg, Germany of 15 a proprietary selection of *Petunia*×*hybrida* identified as code number T09-7350-001, not patented, as the female, or seed, parent with a proprietary selection of *Petunia*×*hybrida* identified as code number T08-3051-031, not patented, as the male, or pollen, parent. The new Petunia plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rheinberg, Germany in May, 2012.

Asexual reproduction of the new *Petunia* plant by terminal cuttings in a controlled greenhouse environment in Rheinberg, Germany since June, 2012 has shown that the unique features of this new *Petunia* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Petunia* have not been observed under all possible environmental conditions and cultural practices. The 35 phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Duepotmoca'.

These characteristics in combination distinguish 'Duepotmoca' as a new and distinct *Petunia* plant:

- 1. Compact, upright and mounding plant habit.
- 2. Moderately vigorous growth habit.
- 3. Freely branching habit.
- 4. Early and freely flowering habit.
- 5. Pink and cream-colored flowers with red purple-colored venation which gives a brownish appearance.
- 6. Good garden performance.

Plants of the new *Petunia* can be compared to plants of the female parent selection. Plants of the new Petunia differ primarily from plants of the female parent selection in flower color as plants of the female parent selection have yellowcolored flowers. In addition, plants of the new *Petunia* are more compact than plants of the female parent selection.

Plants of the new *Petunia* can be compared to plants of the male parent selection. Plants of the new *Petunia* differ primarily from plants of the male parent selection in flower color as plants of the male parent selection have white and red-colored flowers. In addition, plants of the new *Petunia* are more com-20 pact than plants of the male parent selection.

Plants of the new *Petunia* can be compared to plants of the Petunia×hybrida 'Bonnie Yellow', not patented. In side-byside comparisons conducted in Rheinberg, Germany, plants of the new Petunia differed primarily from plants of 'Bonnie Yellow' in the following characteristics:

- 1. Plants of the new *Petunia* were not as broad as plants of 'Bonnie Yellow'.
- 2. Plants of the new *Petunia* had thicker stems than plants of 'Bonnie Yellow'.
- 3. Plants of the new *Petunia* had smaller leaves than plants of 'Bonnie Yellow'.
- 4. Plants of the new *Petunia* and 'Bonnie Yellow' differed in flower color as plants of 'Bonnie Yellow' had light yellow-colored flowers.
- 5. Plants of the new *Petunia* had shorter peduncles than plants of 'Bonnie Yellow'.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new Petunia plant showing the colors as 3

10

50

60

65

true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Petunia* plant.

The photograph comprises a side perspective view of a typical flowering plant of 'Duepotmoca' grown in a container.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown during the summer in 12-cm containers in a glass-covered greenhouse in Rheinberg, Germany and under cultural practices typical of commercial *Petunia* production. During the production of the plants, day and night temperatures averaged 18° C. and light levels averaged 4,500 lux. Plants were pinched one time three weeks after planting and were 16 weeks old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Petunia*×*hybrida* 'Duepotmoca'. Parentage:

Female, or seed, parent.—Proprietary selection of Petunia×hybrida identified as code number T09-7350-001, not patented.

Male, or pollen, parent.—Proprietary selection of Petu-30 nia×hybrida identified as code number T08-3051-031, not patented.

Propagation:

Type.—By terminal cuttings.

Time to initiate roots, summer.—About five days at tem- 35 peratures about 20° C.

Time to initiate roots, winter.—About seven days at temperatures about 20° C.

Time to produce a rooted young plant, summer.—About three weeks at temperatures about 20° C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures about 20° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Compact and mounding plant habit; freely branching habit with about eight to ten lateral branches developing after pinching; moderately vigorous growth habit.

Plant height.—About 22.5 cm.

Plant diameter.—About 32 cm.

Lateral branch description:

Length.—About 15.5 cm.

Diameter.—About 3.8 mm.

Internode length.—About 1.6 cm.

Strength.—Moderately strong.

Aspect.—Initially upright to somewhat outwardly spreading.

Texture.—Pubescent.

Color.—Close to 144A to 144B.

Leaf description:

Arrangement.—Before flowering, alternate; after flowering, opposite; simple.

Length.—About 3.1 cm.

Width.—About 1.9 cm.

Shape.—Spatulate.

Apex.—Obtuse.

Base.—Attenuate.

Margin.—Entire.

Texture, upper and lower surfaces.—Pubescent; leathery.

Venation pattern.—Pinnate; arcuate.

Color.—Developing and fully expanded leaves, upper surface: Close to 144A; venation, close to 144A. Developing and fully expanded leaves, lower surface: Close to 144B; venation, close to 144B.

Petioles.—Length: About 4.8 mm. Diameter: About 3.7 mm. Texture, upper and lower surfaces: Pubescent. Color, upper surface: Close to 144C. Color, lower surface: Close to 144D.

Flower description:

Flower type and flowering habit.—Single salverform flowers arising from leaf axils; freely flowering habit with usually about four open flowers and flower buds per lateral branch and about 25 to 30 flowers developing per plant; flowers face mostly upright to outwardly.

Fragrance.—None detected.

Natural flowering season.—Plants flower continuously during the spring and summer in Germany; early flowering habit, plants typically beginning flowering about nine weeks after planting.

Flower longevity.—Individual flowers last about five to seven days on the plant; flowers persistent.

Flower diameter.—About 4.8 cm.

Flower depth (height).—About 3.5 cm.

Flower throat diameter.—About 1.5 cm.

Flower tube length.—About 2.5 cm.

Flower tube diameter.—About 3.8 mm.

Flower buds.—Length: About 1.1 cm. Diameter: About 3.9 mm. Shape: Ovoid. Color: Close to 144C.

Corolla.—Arrangement: Five petals fused at the base and opening into a flared trumpet. Petal lobe length (from throat): About 2.6 cm. Petal lobe width: About 2.7 cm. Petal shape: Roughly spatulate. Petal apex: Obtuse. Petal margin: Entire. Petal texture, upper and lower surfaces: Rippled, glabrous. Throat texture: Rippled, glabrous. Tube texture: Rippled, pubescent. Color: Petal lobe, when opening, upper surface: Close to 62D. Petal lobe, when opening, lower surface: Close to 155D. Petal lobe, fully opened, upper surface: Close to 62C and 158A; venation, close to 64D; color becoming closer to 62B with development. Petal lobe, fully opened, lower surface: Close to 155D; venation, close to 64B. Flower throat: Close to 144C; venation, close to 64A. Flower tube: Close to 144C; venation, close to 144B.

Calyx.—Arrangement: Five sepals fused at the base forming a star-shaped calyx. Sepal length: About 1.8 cm. Sepal width: About 5.1 mm. Sepal shape: Oblong. Sepal apex: Rounded. Sepal margin: Entire. Sepal texture, upper and lower surfaces: Smooth. Color, upper surface: Close to 144A. Color, lower surface: Close to 144B.

Peduncles.—Length: About 2.4 cm. Diameter: About 3 mm. Strength: Moderately strong. Texture: Smooth. Color: Close to 144B.

Reproductive organs.—Stamens: Quantity per flower: Five. Filament length: About 1.9 mm. Filament color: Close to 144D. Anther length: About 1 mm. Anther shape: Ovate. Anther color: Close to 20A. Pollen

5

Pathogen & pest resistance: Plants of the new *Petunia* have not been observed to be resistant to pathogens and pests common to *Petunia* plants.

amount: Abundant. Pollen color: Close to 155C. Pistils: Quantity per flower: One. Pistil length: About 2.2 cm. Style length: About 1.9 cm. Style color: Close to 145A. Stigma shape: Rounded. Stigma color: Close to 144A. Ovary color: Close to 145A. Seeds and fruits: Seed and fruit development have not been observed on plants of the new *Petunia*.

Garden performance: Plants of the new *Petunia* have been observed to have good garden performance and tolerate wind, rain and temperatures ranging from about 5° C. to about 40° C.

It is claimed:

1. A new and distinct *Petunia* plant named 'Duepotmoca' as illustrated and described.

* * * * *

6

