

(12) **United States Plant Patent**
Higaki

(10) **Patent No.:** **US PP26,004 P3**
(45) **Date of Patent:** **Oct. 20, 2015**

(54) **HYDRANGEA PLANT NAMED ‘BC7.3’**

(50) Latin Name: *Hydrangea macrophylla* (Thunb)
Varietal Denomination: **BC7.3**

(71) Applicant: **Harrison M. Higaki**, San Mateo, CA (US)

(72) Inventor: **Harrison M. Higaki**, San Mateo, CA (US)

(73) Assignee: **Bay City Flower Company, Inc.**, Half Moon Bay, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 61 days.

(21) Appl. No.: **13/986,045**

(22) Filed: **Mar. 25, 2013**

(65) **Prior Publication Data**
US 2014/0289914 P1 Sep. 25, 2014

(51) **Int. Cl.**
A01H 5/02 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./250**
CPC **A01H 5/02** (2013.01)

(58) **Field of Classification Search**
USPC Plt./250
CPC A01H 5/02; A01H 5/00; A01H 1/02
See application file for complete search history.

(56) **References Cited**
U.S. PATENT DOCUMENTS
PP23,801 P3 * 8/2013 Higaki Plt./250
* cited by examiner

Primary Examiner — June Hwu

(74) *Attorney, Agent, or Firm* — James R. Cypher; Charles R. Cypher

(57) **ABSTRACT**

A new and distinct cultivar of *Hydrangea macrophylla* (Thunb.) named ‘BC7.3’ originated as a controlled cross between varieties. The cultivar ‘BC7.3’ can be blue or red depending on the acidity of the soil and the presence of aluminum. The variety ‘BC7.3’ has relatively compact, attractive inflorescences with relatively small sepalous florets, distinct sepal pigmentation and good commercial characteristics. When grown in the presence of aluminum, the upperside of the sepals is primarily R.H.S. 94 A (violet-blue group) and 86 B (violet group), the underside of the sepals is R.H.S. 93 D (violet-blue group) and 86 C violet group).

3 Drawing Sheets

1

Botanical classification: *Hydrangea macrophylla* (Thunb.) ‘BC7.3’.

Variety denomination: ‘BC7.3’.

BACKGROUND OF THE INVENTION

This invention relates to a new and distinct cultivar of the Saxifragaceae family. The botanical name of the plant is *Hydrangea macrophylla* (Thunb.) ‘BC7.3’.

The new cultivar originated as a seedling from a controlled cross between a commercial variety known as ‘Venedig’ to the inventor which may be the subject of U.S. Plant Pat. No. 10,928 and registered as ‘Venice Raven’. ‘Venedig’ was the seed parent and the unpatented variety ‘LK49’ was the pollen parent. ‘LK49’ is relatively compact plant with wiry stems, relatively small leaves, relatively small sepalous florets, and inflorescences that are resistant to being damaged by conditions in commercial coolers.

The variety ‘BC7.3’ has compact, attractive inflorescences with relatively small sepalous florets, attractive sepal pigmentation and good commercial characteristics, including the characteristic that the sepals are highly resistant to osmotic pressure. The variety ‘BC7.3’ has pigmented sepals, and is preferably grown in soil conditions treated with aluminum to produce blue pigmentation. The color of the sepals changes as the plant ages. Below is a table comparing the new variety to similar varieties. Blooms approximately 1 month earlier than similar varieties grown under commercial conditions for sale as a decorative potted plant. The new variety is generally

2

taller than its parents and the stems need to be staked. ‘Venedig’ and the new variety have similarly sized inflorescences.

TABLE 1

	New Variety ‘BC7.3’	Patented Variety ‘BC6.1’ U.S. Plant Pat. No. 23,757
Leaf size	9 cm × 15 cm	12 cm wide × 15 cm long
Plant height	16" in 6" pot.	15" in 6" pot.
Stem strength	Stems are strong but benefit from being staked.	Stems are strong but benefit from being staked.
Sepal Pigmentation	Upper side of sepals is R.H.S. 94 A (violet-blue group) to 86 B (violet group); Under side of sepals is R.H.S. 93 D (violet group) to 86 C (violet group).	Upper side of sepals is R.H.S. 86 A (violet group); Under side of sepals is R.H.S. 88 D (violet group)
Sepalous Floret Size	50 mm	60 mm to 70 mm
	U.S. Plant Pat. No. 18,593 ‘True Blue’	Commerical variety ‘Venedig’ which may be U.S. Plant Pat. No. 10,928 ‘Venice Raven’
Leaf size	Unknown	9 cm wide × 15 cm long- observed controls grown alongside new variety
Plant height	Unknown	12" in 6" pot-observed controls grown alongside new variety.
Stem strength	Strong	Strong-observed controls grown alongside new variety

TABLE 1-continued

Sepal	Both sides of sepals are	Upper side of sepals is R.H.S.
Pigmentation	R.H.S. 100 D (blue group).	84 A (violet group). Under side of sepals is R.H.S. 85 A (violet group) observed controls grown alongside new variety.
Sepalous Floret Size	50 mm to 60 mm	70 mm - observed controls grown alongside new variety

The new cultivar ‘BC7.3’ has been successfully asexually reproduced under controlled environmental conditions at a nursery in Half Moon Bay, Calif. under the direction of the inventor with its distinguishing characteristics remaining stable.

Asexual reproduction was first accomplished when vegetative stem cuttings were taken from the initially selected plant. Examination of asexually reproduced, successive generations grown in Half Moon Bay, Calif. show that the combination of characteristics as herein disclosed for ‘BC7.3’ remains firmly fixed through.

DESCRIPTION OF THE DRAWINGS

The accompanying drawings consist of color photographs that show the typical plant form, including the inflorescence, foliage, and sepals.

FIG. 1 is a view of the entire plant showing its form, growth habit, dark green foliage, inflorescence, and the color of its sepals.

FIG. 2 is a view of the entire plant showing its form, growth habit, dark green foliage, inflorescence, and the color of its sepals.

FIG. 3 is a close-up view of the adaxial surface of a mature leaf.

FIG. 4 is a close-up view of the base of the stem.

FIG. 5 is a close-up view of the upper side of a panicle of the new variety.

FIG. 6 is a close-up view of the center of a panicle of the new variety.

DESCRIPTION OF THE NEW PLANT

The plants shown in the figures are approximately 51 weeks old. The plant started out as cuttings, taken from the stem of a grown plant. The cuttings were placed in a pot and the soil was periodically treated with aluminum to produce blue pigmentation. The plant was pinched early to promote lateral branches.

‘BC7.3’ has not been observed under all possible environmental conditions. The phenotype may vary significantly with variations in environment such as temperature, light intensity and day length. Color determinations were made with The Royal Horticultural Society (R.H.S.) Colour Chart, in association with the Flower Council of Holland, located in Lieden.

THE PLANT

Origin: Controlled cross. The new cultivar originated as a seedling from a controlled cross between the commercial variety known to the inventor as ‘Venedig’ and which may be ‘Venice Raven’, U.S. Plant Pat. No. 10,928, which was the seed parent and the unpatented variety ‘LK49’ which was the pollen parent.

Form: Upright, compact shrub. A typical plant with a mature inflorescence that is ready for sale is approximately 16" high and has a diameter of 16" when grown in a 6" pot with appropriate soil amendments. Plants typically have 3 to 4 stems with one inflorescence between each stem.

Growth: Upright, vigorous growth habit. Inflorescence is large. The plant branches easily with shoots forming at the base of the plant. Lateral branches are similar in appearance and form to the main stems.

Stems: Lenticels are present. Lenticels are R.H.S. 86 A (violet group) and are 1 to 2 mm long. The surface of young stems is glabrous. Stems become woody as they age. The color of typical young stems and young lateral branches is R.H.S. 144 A (green group). The older portions of the stems are R.H.S. 199 B (grey-brown group) Younger portions of the stems are 5 to 7 mm in diameter. Older portions of the stems are 5 to 7 mm in diameter.

Foliage: Abundant. Leaves are opposite on stem and lateral branches.

Shape of leaf.—Elliptic with acute base and apex. Margins are serrate.

Texture.—Glabrous; veins dominate on the underside of the leaf and are sunken on the upper leaf surface.

Color.—Mature leaves have an upper side that is R.H.S. 147 A (yellow-green group), and an under side that is R.H.S. 138 B (green group). Leaves are pinnately veined. The midvein and veins branching off the midvein are large and prominent on the underside of the leaves. Veins are R.H.S. 144 D (yellow-green group). Leaves are as wide as 9 cm and 15 cm long. Petioles are smooth and 2.0 cm long and 4 mm wide. Petioles are R.H.S. 144 D (yellow-green group).

BUDS

Form: Globose with 4 to 5 connate, elliptic, smooth petals. Most buds, whether they will mature into sepalous or non-sepalous florets, have 4 petals. Buds in the center of the inflorescence are non-sepalous. The majority of buds will develop into sepalous florets. They are approximately 1 mm by 1 mm when very young. Buds can be 3 mm in diameter and still unopened. Color of buds is R.H.S. 100 C (blue group).

Aspect: Unopened petals encasing buds are smooth. Arrangement: Borne on branched panicles.

INFLORESCENCE

Form: Paniculate. Terminal. As many as 100 individual flowers (florets) per inflorescence. Both sepalous florets and non-sepalous florets borne on same panicle, with many more non-sepalous florets than sepalous florets. Flowers do not produce a fragrance. The peduncle for the inflorescence is strong and upright. Very few non-sepalous florets developing early on cymes that are later hidden by sepalous florets. Florets, both sepalous and non-sepalous, have anthers and style. Inflorescences are long-lasting, changing color as they age.

Size of inflorescence: Compact and globose. Individual inflorescence size is dependent on the number of florets. A typical inflorescence can grow as large as 7" in diameter, and 4" high.

Shape: Clusters of numerous small florets; sepalous florets overlap one another. Sepals are persistent.

Appearance: Showy.

FLORETS

General: The non-sepalous florets at the center of the inflorescence open first. Sepalous and non-sepalous florets are perfect and complete.

Corolla.—Generally, for both sepalous and non-sepalous florets there are 4 petals which fall off as flower matures. Petals are typically 4 mm long and 3 mm wide. Lenticels that are very small are present on pedicels of both sepalous and non-sepalous florets. Pedicel of both sepalous and non-sepalous florets are primarily 95 C (violet-blue group) when inflorescence is ready for commercial sale. Non-sepalous florets typically lack petals and are 3 mm in diameter. Color of petals before they fall off are R.H.S. 100 C (blue group).

Stamens: 8 stamens, for both sepalous florets and non-sepalous florets. Pollen is R.H.S. 158 C (yellow-white group). Plant produces abundant pollen. Filament is approximately 3 mm long. Filament is R.H.S. 155 C (white group). Anther is 1 mm long and is regular and basally attached.

Stigma: Two to three style each, although most florets have two style, for both sepalous florets and non-sepalous florets. Each style has one stigma. Style is typically 1 mm long. For young inflorescences ready for commercial sale style is R.H.S. 155 D (white group) and stigma is R.H.S. 155 D (white group).

Ovary: Ovary is partially inferior, for both sepalous florets and non-sepalous florets.

Sepalous florets:

General.—Veins dominate on the underside of the sepals.

Number of sepals.—4 or 5 sepals per floret, usually 4.

Aspect of sepals.—Smooth and glaucescent.

Shape of sepals.—Reniform with acuminate apex.

Edges are entire, but with much crenation.

Size of sepals.—As the florets mature, the sepals enlarge and overlap each other more and more, until, often, there is no space between the sepals when the petals of the florets open. Sepals at maturity are typically 3.0 cm long and 4.0 cm wide. Flowers are typically 5 cm in diameter. The upper sides of the sepals are primarily R.H.S. 94 A (violet-blue group) and 86 B (violet group) and the under sides are primarily R.H.S. 93 D (violet-blue group) and R.H.S. 86 C (violet group). Blue pigmentation develops at the tips of the sepals and travels inward towards the base of the sepals. Color of the sepalous florets changes as the inflorescence ages. The upper sides of older sepals are primarily R.H.S. 77 A (purple group) and 93 D (violet group), becoming R.H.S. 138 A (green group).

Fruit: None.

Disease and pest resistance: Unknown.

Other distinguishing characteristics: The sepals of the new variety are highly resistant to osmotic pressure which means the blooms age well and can be sold as “vintage” varieties. Plants are typically held for another 10-12 weeks after the blooms reach maturity before they are sold. During the holding period, because the sepals are resistant to osmotic pressure, the inflorescences do not have to be cleaned as often to remove dessicated florets and sepals.

I claim:

1. A new and distinct *Hydrangea macrophylla* plant named ‘BC7.3’ substantially as herein shown and described.

* * * * *



Fig. 1



Fig. 2




Fig. 3




Fig. 4



FIG. 5



FIG. 6