



US00PP26234P2

(12) **United States Plant Patent**  
**Geibel**

(10) **Patent No.:** **US PP26,234 P2**  
(45) **Date of Patent:** **Dec. 15, 2015**

(54) **ANGELONIA PLANT NAMED ‘LOWHIBLU’**

(50) Latin Name: *Angelonia hybrida*  
Varietal Denomination: **Lowhiblu**

(71) Applicant: **Martin Geibel**, Dresden (DE)

(72) Inventor: **Martin Geibel**, Dresden (DE)

(73) Assignee: **Elsner pac Jungpflanzen Gbr**, Dresden (DE)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 28 days.

(21) Appl. No.: **13/999,884**

(22) Filed: **Apr. 1, 2014**

(51) **Int. Cl.**  
**A01H 5/02** (2006.01)

(52) **U.S. Cl.**

USPC ..... **Plt./404**

(58) **Field of Classification Search**

USPC ..... **Plt./404**

CPC ..... **A01H 5/02**

See application file for complete search history.

*Primary Examiner* — Kent L Bell

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Angelonia* plant named ‘Lowhiblu’ characterized by its compact, upright and outwardly slanting plant habit; freely branching habit; early and freely flowering habit; large white and violet blue-colored flowers; and good garden performance.

**2 Drawing Sheets**

**1**

Botanical designation: *Angelonia hybrida*.  
Cultivar denomination: ‘LOWHIBLU’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of *Angelonia* plant, botanically known as *Angelonia hybrida* and hereinafter referred to by the name ‘Anwedgim’.

The new *Angelonia* plant is a product of a planned breeding program conducted by the Inventor in Dresden, Germany. The objective of the breeding program is to create new compact and freely-flowering *Angelonia* plants with attractive flower coloration.

The new *Angelonia* plant is a naturally-occurring branch mutation of a proprietary selection of *Angelonia hybrida* identified as code number 7036, not patented. The new *Angelonia* plant was discovered and selected by the Inventor on a single flowering plant within a population of plants of the mutation parent in a controlled greenhouse environment in Dresden, Germany during the summer of 2008.

Asexual reproduction of the new *Angelonia* plant by cuttings in a controlled greenhouse environment in Dresden, Germany since the autumn of 2008 has shown that the unique features of this new *Angelonia* plant are stable and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

Plants of the new *Angelonia* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Lowhiblu’. These characteristics in combination distinguish ‘Lowhiblu’ as a new and distinct *Angelonia* plant:

1. Compact, upright and outwardly slanting plant habit.
2. Freely branching habit.

**2**

3. Early and freely flowering habit.
4. Large white and violet blue-colored flowers.
5. Good garden performance.

Plants of the new *Angelonia* differ from plants of the mutation parent primarily in flower color as plants of the mutation parent have solid light blue-colored flowers.

Plants of the new *Angelonia* can be compared to plants of *Angelonia hybrid* ‘Anwedg’, disclosed in U.S. Plant Pat. No. 19,696. In side-by-side comparisons conducted in Dresden, Germany, plants of the new *Angelonia* and ‘Anwedg’ differed in the following characteristics:

1. Plants of the new *Angelonia* were shorter than plants of ‘Anwedg’.
2. Plants of the new *Angelonia* had smaller leaves than plants of ‘Anwedg’.
3. Plants of the new *Angelonia* flowered earlier than plants of ‘Anwedg’.
4. Plants of the new *Angelonia* had larger flowers than plants of ‘Anwedg’.
5. Plants of the new *Angelonia* and ‘Anwedg’ differed in flower color.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying colored photographs illustrate the overall appearance of the new *Angelonia* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Angelonia* plant.

The photograph on the first sheet comprises a side perspective view of a typical flowering plant of ‘Lowhiblu’ grown in a container.

The photograph on the second sheet is a close-up view of a typical flowering plant of ‘Lowhiblu’.

**DETAILED BOTANICAL DESCRIPTION**

Plants used for the aforementioned photographs and the following observations, measurements and values were

grown during the summer and autumn in 13-cm containers in a glass-covered greenhouse in Dresden, Germany and under cultural practices typical of commercial *Angelonia* production. During the production of the plants, day temperatures were about 20° C., night temperatures were about 16° C. and light levels ranged from 15 kilolux to 100 kilolux. Plants were pinched two times, two and six weeks after planting. Plants were five months old when the description was taken and six months old when the photographs were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Angelonia hybrida* 'Lowhiblu'.

Parentage: Naturally-occurring branch mutation of a proprietary selection of *Angelonia hybrida* identified as code number 7036, not patented.

Propagation:

*Type*.—By cuttings.

*Time to initiate roots*.—About 20 days at temperatures about 20° C.

*Time to produce a rooted young plant*.—About four weeks at temperatures about 20° C.

*Root description*.—Fine, fibrous; white in color.

*Rooting habit*.—Freely branching; moderately dense.

Plant description:

*Plant form and growth habit*.—Herbaceous perennial; compact, upright and outwardly slanting plant habit; broad inverted triangle; freely branching habit; when pinched, about six to eight lateral branches develop per plant; moderately vigorous growth habit.

*Plant height*.—About 35 cm to 40 cm.

*Plant width (spread)*.—About 30 cm.

*Lateral branches*.—Length: About 20 cm to 25 cm. Diameter: About 3 mm to 4 mm. Internode length: About 1 cm to 1.5 cm. Strength: Moderately strong to strong. Texture: Slightly pubescent. Color: Close to 145B and 166A.

Leaf description:

*Arrangement*.—Opposite, decussate; simple; sessile.

*Length*.—About 2 cm to 7 cm.

*Width*.—About 1 cm to 1.5 cm.

*Shape*.—Lanceolate to oblong.

*Apex*.—Acute.

*Base*.—Attenuate.

*Margin*.—Serrate.

*Texture, upper and lower surfaces*.—Smooth, glabrous.

*Venation pattern*.—Pinnate.

*Color*.—Developing and fully expanded leaves, upper surface: Close to 137A; venation, close to 137A. Developing and fully expanded leaves, lower surface: Close to 137B; venation, close to 145C.

Flower description:

*Flower type and flowering habit*.—Single flowers arranged in terminal racemes; flowers face mostly outwardly; freely flowering habit; full dense inflorescences.

*Fragrance*.—None detected.

*Natural flowering season*.—Early flowering habit, plants begin flowering about eight to nine weeks after planting; in the garden, flowering is continuous from mid-May until frost in Central Europe.

*Postproduction longevity*.—Flowers last about 20 days on the plant; flowers not persistent.

*Flower buds*.—Height: About 5 mm. Diameter: About 5 mm. Shape: Globose. Color: Close to 137A.

*Inflorescence height*.—About 20 cm.

*Inflorescence diameter*.—About 8 cm.

*Flower size*.—About 3.3 cm by 3.3 cm.

*Flower depth*.—About 1 cm.

*Petals*.—Quantity per flower: Typically five or six in a single whorl; petals fused at the base into a tubular throat. Length: About 1.2 cm. Width: About 1.6 cm. Shape: Roughly spatulate. Apex: Rounded. Margin: Entire, undulate. Texture, upper and lower surfaces: Smooth, glabrous; velvety. Color: When opening and fully opened, upper surface: Close to 155C; center, close to 94B; center color becoming closer to 94C with development. When opening and fully opened, lower surface: Close to 155C; center, close to 94C. Throat: Close to 144C, 155C and 94B.

*Sepals*.—Quantity per flower: Typically five in a single whorl. Length: About 5 mm. Width: About 3 mm. Shape: Lanceolate. Apex: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 137A.

*Pedicels*.—Length: About 1.5 cm to 2 cm. Diameter: About 1 mm. Angle: Outward to slightly upright, less than 90° from vertical. Strength: Moderately strong; flexible. Texture: Smooth, glabrous. Color: Close to 146A.

*Reproductive organs*.—Stamens: Quantity per flower: Typically four. Filament length: About 3 mm. Filament color: Close to 155C. Anther shape: Elliptic. Anther length: About 2 mm. Anther color: Brown to grey. Pollen amount: Moderate. Pollen color: Close to 155D. Pistils: Quantity per flower: One. Pistil length: About 5 mm. Stigma shape: Tapering. Stigma color: Close to 155D. Style length: About 4 mm. Style color: Close to 155D. Ovary color: Close to 142B.

*Fruits*.—Length: About 5 mm. Diameter: About 5 mm. Color: Close to 177B.

*Seeds*.—Quantity per flower: About 30 to 50. Length: Less than 1 mm. Diameter: Less than 1 mm. Color: Light brown.

Disease & pest resistance: Plants of the new *Angelonia* have not been noted to be resistant to pathogens and pests common to *Angelonia* plants.

Garden performance: Plants of the new *Angelonia* have been observed to have good garden performance and tolerate rain, wind and temperatures ranging from about 1° C. to 30° C.

It is claimed:

1. A new and distinct *Angelonia* plant named 'Lowhiblu' as illustrated and described.

\* \* \* \* \*



