



(12) **United States Plant Patent**
Hanes

(10) **Patent No.:** **US PP26,294 P2**
(45) **Date of Patent:** **Jan. 5, 2016**

(54) **PELARGONIUM PLANT NAMED ‘PEQZ0007’**

(50) Latin Name: *Pelargonium interspecific*
Varietal Denomination: **PEQZ0007**

(71) Applicant: **Syngenta Crop Protection AG**, Basel
(CH)

(72) Inventor: **Mitchell E. Hanes**, Gilroy, CA (US)

(73) Assignee: **Syngenta Crop Protection AG**, Basel
(CH)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 73 days.

(21) Appl. No.: **13/998,128**

(22) Filed: **Oct. 4, 2013**

(51) **Int. Cl.**
A01H 5/02 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./329**

(58) **Field of Classification Search**
USPC Plt./325, 328, 329
See application file for complete search history.

Primary Examiner — Keith Robinson

(74) *Attorney, Agent, or Firm* — Joshua L. Price

(57) **ABSTRACT**

A new *Pelargonium* plant named ‘PEQZ0007’ particularly
distinguished by the large light red-purple flowers with a bold
red-purple large-sized splash pattern, fairly large inflores-
cences, relatively long peduncles, medium green foliage
without zonation, good branching and rounded plant habit.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed:
Pelargonium interspecific.

Varietal denomination: ‘PEQZ0007’.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new *Pelargonium*,
botanically known as *Pelargonium interspecific*, and herein-
after referred to by the variety name ‘PEQZ0007’.

‘PEQZ0007’ is a product of a planned breeding program.
The new cultivar ‘PEQZ0007’ has large light red-purple flow-
ers with a bold red-purple large-sized splash pattern, fairly
large inflorescences, relatively long peduncles, medium
green foliage without zonation, good branching and rounded
plant habit.

‘PEQZ0007’ originates from a hybridization made in Sep-
tember 2007 in a greenhouse in Gilroy, Calif. The female
parent was an unpatented, proprietary plant identified as
‘10536-1’ with white flowers with a slight splash pattern and
the flower size and splash pattern is smaller when compared
to ‘PEQZ0007’.

The male parent of ‘PEQZ0007’ was an unpatented, pro-
prietary plant, identified as ‘10499-1’ with pink flowers with
a splash pattern. ‘PEQZ0007’ is more free flowering and has
a more rounded plant habit when compared to ‘10499-1’. The
resultant seed was sown in August 2008.

‘PEQZ0007’ was selected as one flowering plant within the
progeny of the stated cross in November 2008 in a greenhouse
in Gilroy, Calif.

The first act of asexual reproduction of ‘PEQZ0007’ was
accomplished when vegetative cuttings were propagated
from the initial selection in November 2008 in a greenhouse
in Gilroy, Calif.

BRIEF SUMMARY OF INVENTION

Horticultural examination of plants grown from cuttings of
the plant initiated in November 2008 in Gilroy, Calif., and
continuing thereafter, has demonstrated that the combination

2

of characteristics as herein disclosed for ‘PEQZ0007’ are
firmly fixed and are retained through successive generations
of asexual reproduction.

‘PEQZ0007’ has not been observed under all possible envi-
ronmental conditions. The phenotype may vary significantly
with variations in environment such as temperature, light
intensity, and day length.

Plant Breeder’s Rights for this cultivar were applied for in
Canada on Nov. 9, 2012 (12-7790) and at Community Plant
Variety Office on Jun. 4, 2013 (2013/1513). ‘PEQZ0007’ has
not been made publicly available more than one year prior to
the filing of this application.

The following traits have been repeatedly observed and are
determined to be basic characteristics of the new variety. The
combination of these characteristics distinguishes this *Pelar-*
gonium as a new and distinct variety.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographic drawing shows typical
flower and foliage characteristics of ‘PEQZ0007’ with colors
being as true as possible with an illustration of this type.

The photographic drawing shows in FIG. 1 a flowering
plant of the new variety and in FIG. 2 a close-up of an
inflorescence.

DETAILED BOTANICAL DESCRIPTION

The plant descriptions and measurements were taken in
Gilroy, Calif. in August under natural light. These plants were
approximately 25 weeks old and were grown in 6 inch pots,
first in a greenhouse and later outdoors.

The plants shown in the photographs were about 15-18
weeks old, growing in a 6 inch pot in a greenhouse. The
photographs were taken inside a greenhouse in May in
Enkhuizen, The Netherlands.

Color references are made to The Royal Horticultural Soci-
ety Colour Chart (R.H.S.) 2001.

TABLE 1

DIFFERENCES BETWEEN THE NEW VARIETY 'PEQZ0007' AND A MOST SIMILAR VARIETY		
	'PEQZ0007'	'Auld Pikegs' (U.S. Plant Pat. No. 20,481)
Branching habit:	More branches	Fewer branches
Flowering habit:	Is more free-flowering	Is less free-flowering
Plant habit:	Has tighter plant habit	Has a more open plant habit

Plant:

Form, growth and habit.—Upright, outwardly spreading and rounded growth habit. Vigorous, well branched and grows well in sun or semi-shade.
Plant height.—19-21.0 cm.
Plant height (inflorescence included): 21-24.0 cm.
Plant width.—35-40.0 cm.

Roots:

Number of days to initiate roots.—15-18 days at about 22 degrees C.
Number of days to produce a rooted cutting.—21-24 days at 22 degrees C.
Type.—Fine, fibrous, free branching.
Color.—RHS N155B but whiter.

Foliage:

Immature leaf, color upper surface.—Closest to RHS 137A.
Immature leaf, lower surface.—Closest to RHS 137D.
Mature leaf, color upper surface.—Closest to RHS 137A.
Mature leaf, color lower surface.—Closest to RHS 137D.
Length.—4.4-4.9 cm.
Width.—7.2-8.1 cm.
Shape.—Reniform.
Base shape.—Cordate.
Apex shape.—Rounded.
Margin.—Crenate.
Texture upper side.—Hirsute.
Texture lower side.—Hirsute.
Leaf zonation.—Is not present.
Color of veins, upper surface.—Mostly indistinct from foliage color.
Color of veins, lower surface.—Mostly indistinct from foliage color.
Pattern of veining.—Palmate.
Petiole color.—RHS 138B.
Petiole length.—6.2-7.0 cm.
Diameter of petiole.—0.2 cm.
Strength of petiole.—Moderately strong.
Texture.—Moderately smooth and hirsute.
Arrangement.—Typical.

Stem:

Quantity of branches.—4-5.
Color of stem.—RHS 137B to RHS 137C.
Length of stem.—10-12.0 cm.
Diameter.—0.6 cm.
Length of internodes.—0.5-1.0 cm.
Texture.—Hirsute.

Peduncle:

Color of peduncle.—RHS 137B to RHS 137C.
Length of peduncle.—7-8.0 cm.
Peduncle diameter.—0.25 cm.

Texture.—Hirsute, pilose.
Strength of peduncle.—Moderately strong.

Pedicel:

Color of pedicel.—RHS 137C, with anthocyanins of about RHS 178B to RHS 178C at the pedicel apex under the floret.
Length of pedicel.—2.1-3.0 cm.
Diameter of pedicel.—0.15 cm.
Texture.—Hirsute, pilose.
Strength of pedicel.—Moderately strong.

Bud (just before opening):

Color.—RHS N57A but a little darker and more purple.
Length.—1.5-1.7 cm.
Width.—0.5-0.7 cm.
Shape.—Elliptical.
Texture.—Smooth.

Inflorescence:

Type.—Umbel; semi-spherical or nearly semi-spherical.
Duration of flowering.—Continuous upon commencement.
Lastingness of individual flowers.—7-9 days at 18° C. temperature.
Number of inflorescences per plant.—10-15, with 4 immature umbels in various sizes.
Fragrance.—None.
Umbel diameter.—9-10.0 cm.
Umbel depth.—5.5-6.0 cm.

Corolla:

Form.—Single.
Number of petals.—5.
Diameter of flower.—4.5-4.9 cm.
Depth of flower.—1.5-2.0 cm.
Color upper petals, upper surface.—RHS N66C overlaid with a bold blotch of closest to RHS N66A, and a base of RHS N155B and veins of between RHS 60B and RHS 61B.
Color upper petals, lower surface.—RHS 67C with mid-veins of RHS 59B.
Length of upper petals.—2.7-2.9 cm.
Width of upper petals.—1.5-1.8 cm.
Color lower petals, upper surface.—RHS N66C overlaid with a bold blotch of closest to RHS N66A, and a base of between RHS N66B and RHS 66C and veins of between RHS 60B and RHS 61B.
Color lower petals, lower surface.—RHS 67C with mid-veins of RHS 59B.
Length of lower petals.—2.7-2.8 cm.
Width of lower petals.—2.1-2.2 cm.
Petal shape.—Spathulate.
Apex shape.—Rounded.
Margin.—Entire.
Base.—Cuneate.
Petal texture, upper surface.—Papillose.
Petal texture, lower surface.—Papillose.
Number of petaloids.—None present.

Calyx:

Number of sepals.—5.
Color of sepals.—RHS 146B.
Length of sepals.—1-1.1 cm.
Width of sepals.—0.3-0.35 cm.
Sepal shape.—Lanceolate.
Apex shape.—Acute.
Margins.—Mostly fused.
Texture, upper surface.—Glabrous.
Texture, lower surface.—Hirsute, glandular hairs.

Reproductive organs:

Gynoecium.—Pistil: 1. Length: 0.7-0.8 cm. Style color: RHS 155C. Style length: 0.5-0.7 cm. Stigma color: RHS 61B. Sigma type: 5-parted. Ovary color: Closest to RHS 149D, though some are lighter. Ovary length: 0.4 cm. Ovary diameter: 0.15 cm.

Androecium.—Number of stamens: About 8-10; though many are not fully formed or developed. Color of filaments: RHS 155C with between RHS 61B and

RHS 61C at the apex. Length filaments: 0.3-0.6 cm. Anther color: Closest to RHS 48A. Length of anthers: 0.2 cm. Color of pollen: RHS N25A. Pollen amount: Medium. Fertility/Seed set: Moderate seed set. Disease/Pest resistance: Has not been determined to date.

What is claimed is:

1. A new and distinct variety of *Pelargonium* plant named ‘PEQZ0007’ substantially as illustrated and described herein.

* * * * *




FIGURE 1




FIGURE 2