



US00PP26507P2

(12) **United States Plant Patent**
Misato(10) **Patent No.:** US PP26,507 P2
(45) **Date of Patent:** Mar. 15, 2016

- (54) **MANDEVILLA PLANT NAMED
'SUNPARAPREHO'**
- (50) Latin Name: *Mandevilla hybrida*
Varietal Denomination: Sunparapreho
- (71) Applicant: **Tomoya Misato**, Shiga (JP)
- (72) Inventor: **Tomoya Misato**, Shiga (JP)
- (73) Assignee: **Suntory Flowers Limited**, Tokyo (JP)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 177 days.
- (21) Appl. No.: **13/999,243**
- (22) Filed: **Jan. 31, 2014**

- (51) **Int. Cl.**
A01H 5/02 (2006.01)
- (52) **U.S. Cl.**
USPC **Plt./232**
- (58) **Field of Classification Search**
USPC Plt./232
See application file for complete search history.

Primary Examiner — Anne Grunberg
(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Mandevilla* plant named 'Sunparapreho', characterized by its vining plant habit; moderately vigorous growth habit; freely branching habit; dark green-colored leaves; freely flowering habit; long flowering period; and medium-sized white-colored flowers.

1 Drawing Sheet

1

Botanical designation: *Mandevilla hybrida*.
Cultivar denomination: 'SUNPARAPREHO'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Mandevilla* plant, botanically known as *Mandevilla hybrida* and hereinafter referred to by the name 'Sunparapreho'.

The new *Mandevilla* plant is a product of a planned breeding program conducted by the Inventor in Higashiom, Shiga, Japan. The objective of the breeding program is to create new freely-branching and vining *Mandevilla* plants with numerous medium-sized attractive flowers.

The new *Mandevilla* plant originated from an open-pollination in Higashiom, Shiga, Japan in October, 2009 of a proprietary selection of *Mandevilla hybrida* identified as code number 06M-28-09, not patented, as the female, or seed parent with an unknown proprietary selection of *Mandevilla hybrida*, as the male, or pollen, parent. The new *Mandevilla* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated open-pollination in a controlled greenhouse environment in Higashiom, Shiga, Japan in November, 2010.

Asexual reproduction of the new *Mandevilla* plant by cuttings in Higashiom, Shiga, Japan since November, 2010 has shown that the unique features of this new *Mandevilla* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Mandevilla* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Sunpar-

2

preho'. These characteristics in combination distinguish 'Sunparapreho' as a new and distinct *Mandevilla* plant:

1. Vining plant habit.
2. Moderately vigorous growth habit.
3. Freely branching habit.
4. Dark green-colored leaves.
5. Freely flowering habit.
6. Long flowering period.
7. Medium-sized white-colored flowers.

Plants of the new *Mandevilla* can be compared to plants of the female parent selection. Plants of the new *Mandevilla* differ primarily from plants of the female parent selection in the following characteristics:

1. Plants of the new *Mandevilla* have smaller flowers than plants of the female parent selection.
2. Plants of the new *Mandevilla* and the female parent selection differ in flower color as plants of the female parent selection have red-colored flowers.

Plants of the new *Mandevilla* can be compared to plants of the *Mandevillaxamabilis* 'Sunparamiho', disclosed in U.S. Plant Pat. No. 22,868. In side-by-side comparisons conducted in Higashiom, Shiga, Japan, plants of the new *Mandevilla* differed from plants of 'Sunparamiho' in the following characteristics:

1. Plants of the new *Mandevilla* had shorter lateral branches with shorter internodes than plants of 'Sunparamiho'.
2. Plants of the new *Mandevilla* were not as vigorous as plants of 'Sunparamiho'.
3. Plants of the new *Mandevilla* had smaller leaves than plants of 'Sunparamiho'.
4. Plants of the new *Mandevilla* flowered earlier than plants of 'Sunparamiho'.
5. Plants of the new *Mandevilla* had shorter peduncles than plants of 'Sunparamiho'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Mandevilla* plant showing the col-

ors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the actual colors of the new *Mandevilla* plant.

The photograph at the top of the sheet comprises a side perspective view of a typical flowering plant of 'Sunparapreho' grown in a container.

The photograph at the bottom of the sheet is a close-up view of a typical flower of 'Sunparapreho'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown during the late summer in 18-cm containers in an outdoor nursery in Higashiomii, Shiga, Japan and under cultural practices typical of commercial *Mandevilla* production. During the production of the plants, day temperatures averaged 25° C. and night temperatures averaged 15° C. Plants were six months old when the photographs and detailed description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Mandevilla hybrida* 'Sunparapreho'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Mandevilla hybrida* identified as code number 06M-28-09, not patented.

Male, or pollen, parent.—Unknown proprietary selection of *Mandevilla hybrida*, not patented.

Propagation:

Type.—By vegetative cuttings.

Time to initiate roots.—About two weeks at temperatures about 23° C. to 25° C.

Time to produce a rooted young plant.—About five to six weeks at temperatures about 23° C. to 25° C.

Root description.—Fibrous; light brown in color.

Rooting habit.—Freely branching; medium density.

Plant description:

Plant and growth habit.—Vining plant habit; moderately vigorous growth habit; freely branching habit.

Lateral branch description.—Length: About 115 cm. Diameter: About 2.5 mm. Internode length: About 3.2 cm. Strength: Strong, flexible. Texture: Smooth, glabrous; with development, woody. Color, developing: Close to 143A tinted with close to 165A. Color, woody: Close to N199A to N199B.

Leaf description:

Arrangement.—Opposite, simple.

Length.—About 5.6 cm.

Width.—About 3.1 cm.

Shape.—Elliptical.

Apex.—Cuspidate.

Base.—Obtuse.

Margin.—Entire.

Texture, upper and lower surfaces.—Smooth, glabrous.

Venation pattern.—Pinnate, reticulate.

Color.—Developing leaves, upper surface: Close to 144A. Developing leaves, lower surface: Close to 144B. Fully expanded leaves, upper surface: Close to

N137A; venation, close to 144C. Fully expanded leaves, lower surface: Close to 137C; venation, close to 144D.

Petiole length.—About 1.5 cm.

Petiole diameter.—About 1.4 mm.

Petiole texture, upper and lower surfaces.—Smooth, glabrous.

Petiole color, upper and lower surfaces.—Close to 144A.

Flower description:

Flower type and habit.—Single salverform flowers arranged in axillary racemes; flowers face upright to outwardly; freely flowering habit with about five flowers per inflorescence and about 25 inflorescences developing per plant during the flowering season.

Natural flowering season.—Plants begin flowering about six weeks after planting; long flowering period, plants flower continuously from early summer to late autumn in Japan.

Flower longevity on the plant.—About seven to ten days; flowers not persistent.

Fragrance.—None detected.

Inflorescence height.—About 12 cm.

Inflorescence diameter.—About 10 cm.

Flowers.—Appearance: Salverform; flared trumpet, corolla fused and five-parted; flowers roughly star-shaped. Diameter: About 6.6 cm. Depth (length): About 5.1 cm. Throat diameter: About 1.5 cm. Tube length: About 4.2 cm. Tube diameter, mid-section: About 9.8 mm. Tube diameter, base: About 3.6 mm.

Flower buds.—Height: About 5.7 cm. Diameter: About 1.2 cm. Shape: Lenticular. Color: Towards the apex, close to 69A; mid-section, close to 145D; towards the base, close to 145B.

Corolla.—Quantity and arrangement: Five petals arranged in a single whorl and fused towards the base into an elongated tube. Petal lobe length: About 2.6 cm. Petal lobe width: About 2.5 cm. Petal lobe shape: Obovate. Petal lobe apex: Rounded. Petal lobe margin: Entire; slightly undulate. Petal lobe texture, upper and lower surfaces: Smooth, glabrous; satiny. Throat texture: Smooth, glabrous; satiny. Tube texture: Smooth, glabrous; satiny. Color: Petal lobe, when opening, upper surface: Close to 69A to 69C. Petal lobe, when opening, lower surface: Close to 69C to 69D. Petal lobe, fully opened, upper surface: Close to NN155C. Petal lobe, fully opened, lower surface: Close to NN155D. Throat: Close to 17A; towards the petal lobes, close to 12B. Tube: Mid-section and apex, close to N155A to N155B; towards the base, close to 145C tinted with close to 180C.

Calyx.—Quantity and arrangement: Five sepals arranged in a single whorl, fused at the base; calyx, star-shaped. Sepal length: About 6.5 mm. Sepal width: About 3.2 mm. Sepal shape: Lanceolate. Sepal apex: Acute. Sepal margin: Entire. Sepal texture, upper and lower surfaces: Smooth, glabrous. Sepal color: When developing, upper surface: Close to 144C tinted with close to 180C. When developing, lower surface: Close to 144C tinted with close to 180D. Fully opened, upper surface: Close to 144C tinted with close to 180A. Fully opened, lower surface: Close to 144C tinted with close to 180B.

Peduncles.—Length: About 27 cm. Diameter: About 2.1 mm. Texture: Smooth, glabrous. Aspect: Upright to outwardly. Color: Close to 137B.

Pedicels.—Length: About 1.5 cm. Diameter: About 1.9 mm. Texture: Smooth, glabrous. Aspect: Obliquely upright. Color: Close to 146A.

Reproductive organs.—Stamens: Quantity and arrangement: Typically five; filaments fused to corolla; anthers, connivent. Anther shape: Ellipsoidal. Anther size: About 1.8 mm by 9.2 mm. Anther color: Close to 4C. Pollen amount: Moderate. Pollen color: Close to 4D. Pistils: Quantity: Typically one. Pistil length: About 2.4 cm. Style color: Close to 145C. Stigma shape: Conical. Stigma color: Close to 145B. Ovary color: Close to 144B.

Seeds and fruits.—Seed and fruit production have not been observed on plants of the new *Mandevilla*.

Disease & pest resistance: Plants of the new *Mandevilla* have not been noted to be resistant to pathogens and pests common to *Mandevilla* plants.

Garden performance: Plants of the new *Mandevilla* have been observed to tolerate wind, rain and temperatures ranging from about 4° C. to about 30° C.

It is claimed:

1. A new and distinct *Mandevilla* plant named ‘Sunparapreho’ as illustrated and described.

* * * * *

