



US00PP26551P2

(12) **United States Plant Patent**  
**Shadow**(10) **Patent No.:** US PP26,551 P2  
(45) **Date of Patent:** Mar. 29, 2016(54) **ALTHEA PLANT NAMED 'DS04PS'**(50) Latin Name: ***Hibiscus syriacus***  
Varietal Denomination: **DS04PS**(71) Applicant: **Don Odom Shadow**, Winchester, TN  
(US)(72) Inventor: **Don Odom Shadow**, Winchester, TN  
(US)(73) Assignee: **Greenleaf Nursery Company**, Park  
Hill, OK (US)(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 88 days.(21) Appl. No.: **13/986,867**(22) Filed: **Jun. 13, 2013**(51) **Int. Cl.**  
**A01H 5/02** (2006.01)(52) **U.S. Cl.**  
USPC ..... **Plt./257**(58) **Field of Classification Search**USPC ..... Plt./257, 226  
CPC ..... A01H 5/02; A01H 5/00  
See application file for complete search history.(56) **References Cited**

## U.S. PATENT DOCUMENTS

PP24,336 P2 \* 3/2014 Woods ..... Plt./257

\* cited by examiner

Primary Examiner — June Hwu

(74) Attorney, Agent, or Firm — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of Althea plant named 'DS04PS', characterized by its upright, somewhat outwardly spreading and uniformly mounded plant habit; freely branching habit, dense and bushy plant form; dark green-colored leaves; early, uniform and freely flowering habit; long flowering period; large double-type ruffled pink-colored flowers with red purple-colored stripes and bases; and good container and garden performance.

## 2 Drawing Sheets

## 1

Botanical designation: *Hibiscus syriacus*.  
Cultivar denomination: 'DS04PS'.

## BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct Althea plant, botanically known as *Hibiscus syriacus*, also known as Rose of Sharon, and hereinafter referred to by the name 'DS04PS'. 5

The new Althea plant originated from an open-pollination in Winchester, Tenn. of an unnamed proprietary selection of *Hibiscus syriacus*, not patented, as the female, or seed, parent with an unknown selection of *Hibiscus syriacus* as the male, or pollen, parent. The new Althea plant was discovered and selected by the Inventor as a single flowering plant within the progeny of the stated open-pollination in a controlled nursery environment in Winchester, Tenn. in July, 2007. 10

Asexual reproduction of the new Althea plant by softwood cuttings in a controlled environment in Winchester, Tenn. since June, 2008 has shown that the unique features of this new Althea plant are stable and reproduced true to type in successive generations. 15

## SUMMARY OF THE INVENTION

Plants of the new Althea have not been observed under all possible environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype. 20

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'DS04PS'. These characteristics in combination distinguish 'DS04PS' as a new and distinct Althea plant: 25

1. Upright, somewhat outwardly spreading and uniformly mounded plant habit.
2. Freely branching habit, dense and bushy plant form.
3. Dark green-colored leaves.
4. Early, uniform and freely flowering habit.
5. Long flowering period.
6. Large double-type ruffled pink-colored flowers with red purple-colored stripes and bases.
7. Good container and garden performance.

Plants of the new Althea can be compared to plants of the parent selection. Plants of the new Althea differ from plants of the female parent selection in the following characteristics:

1. Plants of the new Althea are more uniform, freely branching and denser than plants of the female parent selection.
2. Plants of the new Althea and the female parent selection differ in flower color as plants of the female parent selection have solid red purple-colored flowers.
3. Plants of the new Althea have not been observed to produce fruits and seeds whereas plants of the female parent selection produce fruits and seeds.

Plants of the new Althea can be compared to plants of the *Hibiscus syriacus* 'Notwood3', disclosed in U.S. Plant Pat. No. 20,574. In side-by-side comparisons conducted in Winchester, Tenn., plants of the new Althea differed from plants of 'Notwood3' in the following characteristics:

1. Plants of the new Althea were smaller and more compact than plants of 'Notwood3'.
2. Plants of the new Althea and 'Notwood3' differed in flower color as plants of 'Notwood3' had blue-colored flowers.

## BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new Althea plant showing the colors as

true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Althea plant.

The photograph on the first sheet is a side perspective view of a typical flowering plant of 'DS04PS' grown in a container.

The photograph on the second sheet is a close-up view of a typical open flower of 'DS04PS'.  
10

#### DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown during the summer in 15-cm containers in an outdoor nursery in Park Hill, Okla. and Fort Worth, Tex. under cultural practices typical of commercial Althea production. During the production of the plants, day temperatures ranged from 20° C. to 38° C., night temperatures ranged from 15° C. to 24° C. and light levels ranged from 3,000 to 4,000 foot-candles. Plants were pinched one time and were four years old when the photographs and the description were taken. In the description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.  
15  
20  
25

Botanical classification: *Hibiscus syriacus* 'DS04PS'.

##### Parentage:

*Female, or seed, parent.*—Unnamed proprietary selection of *Hibiscus syriacus*, not patented.

*Male or pollen parent.*—Unknown selection of *Hibiscus syriacus*, not patented.

##### Propagation:

*Type.*—By softwood cuttings in the summer and semi-hardwood cuttings in the winter; plants are typically propagated during the summer.  
35

*Time to initiate roots, summer.*—About 14 to 18 days at soil temperatures about 24° C. to 27° C.

*Time to initiate roots, winter.*—About 20 to 25 days at soil temperatures about 24° C. to 27° C.  
40

*Time to produce a rooted young plant, summer.*—About two to three months at soil temperatures about 24° C. to 27° C.

*Time to produce a rooted young plant, winter.*—About three to six months at soil temperatures about 16° C. to 21° C.  
45

*Root description.*—Medium in thickness, fibrous; white in color.

*Rooting habit.*—Freely branching; dense.

##### Plant description:

*Plant and growth habit.*—Perennial shrub; upright to outwardly spreading and uniformly mounded plant habit; moderately vigorous to vigorous growth habit.

*Branching habit.*—Freely branching habit with lateral branches potentially develop at every node; pinching enhances lateral branch development; dense and bushy plant form.  
55

*Plant height.*—About 57 cm.

*Plant diameter (area of spread).*—About 65 cm.

##### Lateral branch description:

*Length.*—About 34 cm.  
60

*Diameter.*—About 5 mm.

*Internode length.*—About 4.5 cm.

*Aspect.*—Upright to outwardly spreading.

*Texture, immature.*—Smooth, glabrous.

*Texture, mature.*—Woody and somewhat rough.  
65

*Luster, immature and mature.*—Matte, dull.

*Color, immature.*—Close to 194A.

*Color, mature.*—Close to 194A becoming closer to 199B with development.

##### 5 Leaf description:

*Arrangement.*—Alternate, single.

*Length.*—About 7 cm.

*Width.*—About 5.2 cm.

*Shape.*—Overall outline, ovate; palmately lobed.

*Apex.*—Rounded.

*Base.*—Cuneate.

*Margin.*—Crenate; palmately lobed to varying depths, sinuses divergent.

*Texture, upper and lower surfaces.*—Smooth, glabrous.

*Luster, upper and lower surfaces.*—Matte, dull.

*Venation pattern.*—Pinnate; arcuate.

*Color.*—Developing leaves, upper surface: Close to N137A. Developing leaves, lower surface: Close to 146A. Fully expanded leaves, upper surface: Close to N137A to N137B; venation, close to 144A. Fully expanded leaves, lower surface: More green than 146A; mid-vein, close to 145C and lateral veins, close to 144A.  
25

*Petioles.*—Length: About 1.4 cm. Diameter: About 2 mm. Strength: Strong, flexible. Texture, upper and lower surfaces: Pubescence, minute. Luster, upper and lower surfaces: Matte. Color, upper and lower surfaces: More green than 144A.  
30

##### Flower description:

*Flower arrangement and flowering habit.*—Ruffled double-type flowers arranged at terminal leaf axils; uniform and freely flowering habit with about three flowers per terminal; flowers face mostly outwardly.

*Natural flowering season.*—Early flowering habit, plants of the new Althea begin flowering about seven weeks after pinching; long flowering period with plants flowering continuously from spring until the autumn in Oklahoma and Texas.

*Flower longevity.*—Flowers last about two to three days on the plant depending on temperature; flowers not persistent.

*Flower diameter.*—About 9.75 cm.

*Flower length (height).*—About 4.3 cm.

*Flower buds.*—Length: About 1.5 cm. Diameter: About 9 mm. Shape: Ovoid. Texture: Rough, short pubescence. Luster: Matte, dull. Color: Darker green than 144A.

*Petals/petaloids.*—Arrangement: Corolla consists of multiple petals/petaloids in numerous whorls that are fused at base; petals/petaloids imbricate. Length (largest, outer): About 4.75 cm. Width (largest outer): About 3.2 cm. Shape: Roughly spatulate. Apex: Rounded. Base: Cuneate. Margin: Crenate; undulate, ruffled appearance. Texture, upper and lower surfaces: Smooth, glabrous; velvety; somewhat waxy on the lower surface. Luster, upper and lower surface: Matte. Color: When opening and fully opened, upper surface: Close to 69A to 69C; stripes and towards the base, close to 59A; color does not change with development. When opening and fully opened, lower surface: Close to 69B to 69C; towards the base, close to 59B; color does not change with development.

*Sepals.*—Appearance: Five sepals fused into a star-shaped calyx. Length: About 2.5 cm. Width: About 1 cm. Shape: Broadly lanceolate. Apex: Cuspidate.

US PP26,551 P2

5

Base: Fused. Margin: Entire. Texture, upper surface: Smooth, glabrous. Texture, lower surface: Rough, pubescence, minute. Luster, upper surface: Somewhat glossy. Luster, lower surface: Matte, dull. Color, upper surface: Close to 144A to 144B. Color, lower surface: Darker green than 144A.

*Bracts*.—Appearance: About eight fused at base. Length: About 1.4 cm. Width: About 1.5 mm. Shape: Acicular. Apex: Acute. Base: Fused. Margin: Entire. Texture, upper and lower surfaces: Rough, pubescence, minute. Luster, upper and lower surfaces: Matte, dull. Color, upper and lower surfaces: Close to 144A.

*Peduncles*.—Length: About 7.5 mm. Diameter: About 4 mm. Aspect: Upright to slightly outward. Strength: Strong, flexible. Texture: Rough, short pubescence. Color: Close to 144A.

6

*Reproductive organs*.—None observed, reproductive structures transformed into petaloids.

*Seeds and fruits*.—Seed and fruit production has not been observed on plants of the new Althea as flowers are sterile.

Garden performance: Plants of the new Althea have been observed to have good garden performance and to tolerate wind, rain and temperatures ranging from about -29° C. to about 43° C.

10 Pathogen & pest resistance: Plants of the new Althea have not been observed to be resistant to pathogens and pests common to Althea plants.

It is claimed:

15 1. A new and distinct Althea plant named 'DS04PS' as illustrated and described.

\* \* \* \* \*



