

US00PP26923P2

(12) United States Plant Patent

Luo

(10) Patent No.:

US PP26,923 P2

(45) **Date of Patent:**

Jul. 12, 2016

MANDEVILLA PLANT NAMED 'DREAMNUFLOR3512'

Latin Name: *Mandevilla hybrida* Varietal Denomination: **Dreamnuflor3512**

Applicant: Shuming Luo, Dulwich Hill (AU)

Shuming Luo, Dulwich Hill (AU) Inventor:

Assignee: Nuflora International Pty. Ltd., (73)

Macquarie Fields, NSW (AU)

Subject to any disclaimer, the term of this Notice:

patent is extended or adjusted under 35

U.S.C. 154(b) by 35 days.

Appl. No.: 14/121,805

Oct. 20, 2014 (22)Filed:

Int. Cl. A01H 5/02 (2006.01)

U.S. Cl. (52)

(58)

Field of Classification Search See application file for complete search history.

Primary Examiner — Anne Grunberg

(74) Attorney, Agent, or Firm — C. A. Whealy

ABSTRACT (57)

A new and distinct cultivar of *Mandevilla* plant named 'Dreamnuflor3512', characterized by its upright and somewhat vining to climbing plant habit; vigorous growth habit; early and freely flowering habit; long flowering period; dark pink-colored flowers; and good garden performance.

1 Drawing Sheet

Botanical designation: Mandevilla hybrida. Cultivar denomination: 'DREAMNUFLOR3512'.

CROSS REFERENCED TO CLOSELY-RELATED APPLICATIONS

Title: *Mandevilla* Plant Named 'Dreamnuflor3513'

Applicant: Shuming Luo

Filed: Concurrently with this application having applica-

tion Ser. No. 14/121,806

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of Mandevilla plant, botanically known as Mandevilla hybrida and hereinafter referred to by the name 'Dreamnuflor3512'.

The new *Mandevilla* plant is a product of a planned breeding program conducted by the Inventor in Cobbitty, New South Wales, Australia. The objective of the breeding program is to create new *Mandevilla* plants that flower for a long period of time and have large attractive flowers.

The new Mandevilla plant originated from a cross-pollination made by the Inventor in Cobbitty, New South Wales, Australia in February, 2009 of a proprietary selection of Mandevilla hybrida identified as code number DIP 799, not patented, as the female, or seed parent with a proprietary selection of Mandevilla hybrida identified as code number 25 DIP 7100, not patented, as the male, or pollen, parent. The new *Mandevilla* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Cobbitty, New South Wales, Australia in May, 30 2010.

Asexual reproduction of the new *Mandevilla* plant by cuttings in Macquarie Fields, New South Wales, Australia since May, 2010 has shown that the unique features of this new Mandevilla plant are stable and reproduced true to type in 35 successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Mandevilla* have not been observed under all possible combinations of environmental conditions and ⁴ cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in geno-

type.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Dreamnuflor3512'. These characteristics in combination distinguish 'Dreamnuflor3512' as a new and distinct Mandevilla plant:

1. Upright and somewhat vining to climbing plant habit.

2. Vigorous growth habit.

3. Early and freely flowering habit.

4. Long flowering period.

5. Dark pink-colored flowers.

6. Good garden performance.

Plants of the new *Mandevilla* can be compared to plants of the female parent selection. Plants of the new *Mandevilla* differ primarily from plants of the female parent selection in 20 the following characteristics:

- 1. Plants of the new *Mandevilla* have smaller leaves than plants of the female parent selection.
- 2. Leaves of plants of the new *Mandevilla* are smooth and glabrous whereas leaves of plants of the female parent selection are pubescent.
- 3. Plants of the new *Mandevilla* and the female parent selection differ in flower color as plants of the female parent selection have pale pink-colored flowers.

Plants of the new *Mandevilla* can be compared to plants of the male parent selection. Plants of the new Mandevilla differ primarily from plants of the male parent selection in the following characteristics:

- 1. Plants of the new *Mandevilla* have larger leaves than plants of the male parent selection.
- 2. Plants of the new *Mandevilla* and the male parent selection differ in flower color as plants of the male parent selection have red-colored flowers.

Plants of the new *Mandevilla* can be compared to plants of 'Dreamnuflor3513', disclosed in U.S. Plant patent application Ser. No. 14/121,806, filed concurrently. Plants of the new

Mandevilla and 'Dreamnuflor3513' differ primarily in flower color as plants of 'Dreamnuflor3513' have bright red-colored flowers.

Plants of the new *Mandevilla* can be compared to plants of *Mandevilla hybrida* 'White Fantasy', not patented. In side-by-side comparisons conducted in Cobbitty, New South Wales, Australia, plants of the new *Mandevilla* differed primarily from plants of 'White Fantasy' in the following characteristics:

- 1. Plants of the new *Mandevilla* had smaller leaves than plants of 'White Fantasy'.
- 2. Leaves of plants of the new *Mandevilla* were smooth and glabrous whereas leaves of plants of 'White Fantasy' were pubescent.
- 3. Plants of the new *Mandevilla* and 'White Fantasy' differed in flower color as plants of 'White Fantasy' had white-colored flowers.

Plants of the new *Mandevilla* can also be compared to plants of *Mandevilla hybrida* 'Crimson Fantasy', not patented. In side-by-side comparisons conducted in Cobbitty, New South Wales, Australia, plants of the new *Mandevilla* differed primarily from plants of 'Crimson Fantasy' in the following characteristics:

- 1. Plants of the new *Mandevilla* had smaller leaves than plants of 'Crimson Fantasy'.
- 2. Leaves of plants of the new *Mandevilla* were smooth and glabrous whereas leaves of plants of 'Crimson Fantasy' were pubescent.
- 3. Plants of the new *Mandevilla* and 'Crimson Fantasy' differed in flower color as plants of 'Crimson Fantasy' had crimson-colored flowers.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Mandevilla* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the following detailed botanical description which accurately describe the actual colors of the new *Mandevilla* plant.

The photograph at the bottom of the sheet comprises a side perspective view of a typical flowering plant of 'Dreamnuflor3512' grown in a container.

The photograph at the top of the sheet is a close-up views of a typical flowering plant of 'Dreamnuflor3512'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown in one-gallon containers during the early summer in a polyethylene-covered greenhouse in Vista, Calif. and under cultural practices typical of commercial *Mandevilla* production. During the production of the plants, day temperatures averaged 27° C. and night temperatures averaged 21° C. Plants were pinched two times and were 19 weeks old when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general 65 terms of ordinary dictionary significance are used.

Botanical classification: *Mandevilla hybrida* 'Dreamnuflor3512'.

Parentage:

Female, or seed, parent.—Proprietary selection of Mandevilla hybrida identified as code number DIP 799, not patented.

Male, or pollen, parent.—Proprietary selection of Mandevilla hybrida identified as code number DIP 7100, not patented.

Propagation:

Type.—By cuttings.

Time to initiate roots, summer.—About ten days at temperatures about 25° C. to 30° C.

Time to initiate roots, winter.—About 16 days at temperatures about 15° C. to 20° C.

Time to produce a rooted young plant, summer.—About four weeks at temperatures about 25° C. to 30° C.

Time to produce a rooted young plant, winter.—About six weeks at temperatures about 15° C. to 20° C.

Root description.—Medium in thickness, fibrous; white in color becoming creamy white to golden yellow with development.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Upright and somewhat vining to climbing plant habit; vigorous growth habit.

Plant height.—About 28 cm.

Plant diameter.—About 27 cm by 30 cm.

Lateral branch description.—Branching habit: Moderate branching habit; pinching enhances lateral branch development; about five primary lateral branches each with numerous secondary branches. Length: About 27 cm. Diameter: About 4 mm. Internode length: About 1.8 cm. Texture: Smooth, glabrous; becoming woody with development. Color: Close to 145A; color becoming closer to 199B with development.

40 Leaf description:

35

50

Arrangement.—Opposite, simple.

Length.—About 6.6 cm.

Width.—About 4.8 cm.

Shape.—Elliptic to oval.

Apex.—Acuminate.

Base.—Equilateral to slightly cordate.

Margin.—Entire.

Texture, upper and lower surfaces.—Smooth, glabrous; leathery.

Venation pattern.—Pinnate, arcuate.

Color.—Developing leaves, upper surface: Close to 146A. Developing leaves, lower surface: Close to 146A to 146B. Fully expanded leaves, upper surface: Close to 137B; venation, close to 146C. Fully expanded leaves, lower surface: Close to 146B; venation, close to 145D.

Petioles.—Length: About 1.6 cm. Diameter: About 2 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper surface: Close to 146C. Color, lower surface: Close to 146D.

Flower description:

Flower type and habit.—Salverform flowers arranged alternately in axillary clusters similar to loose racemes; flowers face horizontally to slightly drooping; freely flowering habit, about six to seven flowers per cluster and about 84 flowers developing per plant.

Natural flowering season.—Early flowering habit, plants begin flowering about four to six weeks after planting; long flowering period, plants flower continuously from late winter into the late autumn in Southern California.

5

Flower longevity on the plant.—About four to five days; flowers not persistent.

Fragrance.—None detected.

Flower buds.—Height: About 9.5 cm. Diameter: About 2 cm. Shape: Elongated oblong to pointed. Color: 10 Close to 73C to 73D.

Cluster height.—About 10.4 cm.

close to 18B.

Cluster diameter.—About 6.5 cm by 10.5 cm.

Flower diameter.—About 6.5 cm to 9.5 cm.

Flower depth (length).—About 7.6 cm.

Flower throat diameter.—About 2 cm.

Flower tube length.—About 5.3 cm. Flower tube diameter, base.—About 5 mm.

Corolla.—Arrangement and appearance: Single whorl of five petals, petals slightly imbricate; lobes asym- ²⁰ metrical giving a pinwheel appearance to the flower. Petal lobe length: About 3.5 cm. Petal lobe width: About 3.5 cm. Petal lobe shape: Asymmetrical; recurved. Petal apex: Acuminate. Petal margin: Entire. Petal texture, upper and lower surfaces: ²⁵ Smooth, glabrous; velvety. Throat texture: Smooth, glabrous. Tube texture: Smooth, glabrous. Color: Petal lobe, when opening, upper surface: Close to 73B to 73C. Petal lobe, when opening, lower surface: Close to 73C. Petal lobe, fully opened, upper surface: 30 Striated, close to 73A, 73B and 73C; venation, close to 73B; color becoming closer to 73B with development. Petal lobe, fully opened, lower surface: Close to 73D; venation, close to 73C to 73D. Throat: Close to 21A; at throat apex, central ring, close to 54A, 54B ³⁵ and 54C; venation, close to 21A. Tube: Proximally, close to 1C; distally, close to 18C to 18D; venation,

Sepals.—Arrangement and appearance: Five per flower fused in a single whorl; calyx short funnelform and five-pointed. Length: About 5 mm. Width: About 3 mm. Shape: Lanceolate. Apex: Acute to slightly acuminate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 145C.

Peduncles.—Length: About 2.7 cm. Diameter: About 2 mm. Texture: Smooth, glabrous. Strength: Strong, flexible. Aspect: About 35° to 45° from lateral branch axis. Color: Close to 145A.

Pedicels.—Length: About 2 cm to 2.5 cm. Diameter: About 2.5 mm. Texture: Smooth, glabrous. Strength: Strong, flexible. Aspect: About 25° to 30° from peduncle axis. Color: Close to 145A.

Reproductive organs.—Stamens: Quantity per flower: Typically five. Filament length: About 7 mm. Filament color: Close to 1D. Anther length: About 8 mm. Anther shape: Lanceolate. Anther color: Close to 161B. Pollen amount: None observed. Pistils: Quantity per flower: Typically one. Pistil length: About 3.2 cm. Style length: About 2.7 cm. Style color: Close to 145D. Stigma shape: Rounded. Stigma color: Close to 145B. Ovary color: Close to 144A.

Seeds and fruits.—Seed and fruit production has not been observed on plants of the new Mandevilla.


Disease & pest tolerance: Plants of the new *Mandevilla* have been observed to be somewhat tolerant to mites. Plants of the new *Mandevilla* have not been noted to be resistant to pathogens and other pests common to *Mandevilla* plants.

Garden performance: Plants of the new *Mandevilla* have been observed to have good garden performance and to tolerate rain, wind, full sunlight and temperatures from about 5° C. to about 45° C.

It is claimed:

1. A new and distinct *Mandevilla* plant named 'Dreamnuflor3512' as illustrated and described.

* * * *



.....

