



(12) **United States Plant Patent**
van den Hoogen

(10) **Patent No.:** **US PP27,122 P2**
(45) **Date of Patent:** **Aug. 30, 2016**

(54) **HYPERICUM PLANT NAMED**
‘ALLLANGANO’

(50) Latin Name: *Hypericum hybrida*
Varietal Denomination: **Alllangano**

(71) Applicant: **Wilhelmus T. J. van den Hoogen**, Cuijk
(NL)

(72) Inventor: **Wilhelmus T. J. van den Hoogen**, Cuijk
(NL)

(73) Assignee: **Allplants Holding B. V.**, Cuijk (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 10 days.

(21) Appl. No.: **14/544,618**

(22) Filed: **Jan. 28, 2015**

(51) **Int. Cl.**
A01H 5/02 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./442**

(58) **Field of Classification Search**
USPC Plt./442, 226
CPC A01H 5/02; A01H 5/00
See application file for complete search history.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Hypericum* plant named ‘All-
langano’, characterized by its upright and outwardly spread-
ing plant habit; moderately vigorous growth habit; dark
green-colored leaves; freely flowering habit and high fruit
density; and glossy red purple-colored fruits.

1 Drawing Sheet

1

Botanical designation: *Hypericum hybrida*.
Cultivar denomination: ‘ALLLANGANO’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Hypericum* plant, botanically known as *Hypericum*
hybrida, typically grown as a cut-flower and hereinafter
referred to by the name ‘Alllangano’.

The new *Hypericum* plant is a product of a planned breed-
ing program conducted by the Inventor in Cuijk, The Neth-
erlands. The objective of the breeding program is to create
new *Hypericum* plants with strong stems and numerous
attractive fruits.

The new *Hypericum* plant originated from an open-poll-
ination in September, 2009 in Cuijk, The Netherlands of a
proprietary selection of *Hypericum hybrida* identified as code
number KZ05-54-3, not patented, as the female, or seed par-
ent with an unknown selection of *Hypericum hybrida* as the
male, or pollen, parent. The new *Hypericum* plant was dis-
covered and selected by the Inventor as a single flowering
plant from within the progeny of the stated open-pollination
in a controlled greenhouse environment in Cuijk, The Neth-
erlands in September, 2010.

Asexual reproduction of the new *Hypericum* plant by veg-
etative cuttings in a controlled greenhouse environment in
Cuijk, The Netherlands since April, 2012 has shown that the
unique features of this new *Hypericum* plant are stable and
reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Hypericum* have not been observed under
all possible combinations of environmental conditions and
cultural practices. The phenotype may vary somewhat with
variations in environmental conditions such as temperature
and light intensity without, however, any variance in geno-
type.

2

The following traits have been repeatedly observed and are
determined to be the unique characteristics of ‘Alllangano’.
These characteristics in combination distinguish ‘Alllan-
gano’ as a new and distinct *Hypericum* plant:

1. Upright and outwardly spreading plant habit.
2. Moderately vigorous growth habit.
3. Dark green-colored leaves.
4. Freely flowering habit and high fruit density.
5. Glossy red purple-colored fruits.

Plants of the new *Hypericum* differ primarily from plants
of the female parent selection in the following characteristics:

1. Leaves of plants of the new *Hypericum* curl upwards
whereas leaves of plants of the female parent selection
curled strongly downwards.
2. Plants of the new *Hypericum* and the female parent
selection differ in fruit shape as plants of the female
parent selection have oval-shaped fruits.

Plants of the new *Hypericum* can be compared to plants of
the *Hypericum hybrida* ‘Allrio’, not patented. In side-by-side
comparisons, conducted in Cuijk, The Netherlands, plants of
the new *Hypericum* differed primarily from plants of ‘Allrio’
in the following characteristics:

1. Plants of the new *Hypericum* had larger fruits than plants
of ‘Allrio’.
2. Plants of the new *Hypericum* and ‘Allrio’ differed in fruit
color as plants of ‘Allrio’ had red-colored fruits.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the over-
all appearance of the new *Hypericum* plant showing the col-
ors as true as it is reasonably possible to obtain in colored
reproductions of this type. Colors in the photograph may
differ slightly from the color values cited in the detailed
botanical description which accurately describe the actual
colors of the new *Hypericum* plant.

The photograph comprises a side perspective view of a
typical fruiting stem of ‘Alllangano’.

DETAILED BOTANICAL DESCRIPTION

Plants used for the aforementioned photographs and fol-
lowing description were grown in September in an outdoor

nursery in Timau, Kenya and under cultural practices typical of commercial cut-flower *Hypericum* production. During the production of the plants, day temperatures ranged from 15° C. to 25° C. and night temperatures ranged from 5° C. to 15° C. Plants were one year old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Hypericum hybrida* 'Alllangano'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Hypericum hybrida* identified as code number KZ05-54-3, not patented.

Male, or pollen, parent.—Unknown selection of *Hypericum hybrida*, not patented.

Propagation:

Type cutting.—Vegetative cuttings.

Time to initiate roots, summer.—About 10 to 14 days at temperatures about 12° C. to 30° C.

Time to initiate roots, winter.—About 14 to 20 days at temperatures about 12° C. to 22° C.

Time to produce a rooted young plant, summer.—About 24 to 32 days at temperatures ranging from 12° C. to 30° C.

Time to produce a rooted young plant, winter.—About 28 to 36 days at temperatures ranging from 12° C. to 22° C.

Root description.—Fine, fleshy; white in color.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Perennial shrub; upright and outwardly spreading plant habit, narrow inverted triangle; moderately vigorous growth habit.

Branching habit.—Moderately freely basal branching habit; pinching enhances lateral branch development.

Plant height.—About 63.3 cm.

Plant width (spread).—About 40.5 cm.

Lateral branch description.—Length: About 64.5 cm. Diameter: About 4 mm. Internode length: About 5.7 cm. Strength: Moderately strong to strong. Texture: Smooth, glabrous. Luster: Moderately glossy. Color, developing: Close to 145A moderately tinged with close to 182C to 182D. Color, fully developed: Close to N199C to N199D.

Leaf description:

Arrangement.—Opposite, simple; sessile.

Length.—About 7 cm.

Width.—About 4.7 cm.

Shape.—Ovate.

Apex.—Retuse.

Base.—Truncate to shallowly cordate.

Margin.—Entire.

Texture, upper and lower surfaces.—Smooth to slightly rugose; glabrous.

Luster, upper and lower surfaces.—Slightly glossy.

Venation pattern.—Pinnate.

Color.—Developing leaves, upper surface: Close 137A. Developing leaves, lower surface: Close to 138B. Fully expanded leaves, upper surface: Darker than between 147A and N189A; venation, close to 144B. Fully expanded leaves, lower surface: Close to between 138B and 147B; venation, close to 144B.

Flower description:

Flower type, arrangement and flowering habit.—Single rotate flowers arranged in terminal and axillary compound cymes; freely flowering habit with about 16 flowers per cyme; flowers face mostly upright.

Fragrance.—None detected.

Natural flowering season.—Plants flower continuously from mid-June to mid-August in Kenya; plants begin flowering about ten months after planting.

Flower longevity.—Flowers last about two or three days on the plant; flowers not persistent.

Fruit longevity.—About one month on the plant.

Flower buds.—Length: About 8 mm. Diameter: About 7 mm. Shape: Globular. Color: Close to 12B to 12C; towards the apex, close to 30D.

Inflorescence height.—About 8.4 cm.

Inflorescence diameter.—About 8.2 cm.

Flower diameter.—About 2.5 cm.

Flower depth (height).—About 1.8 cm.

Petals.—Quantity and arrangement: Five in a single whorl. Length: About 1.2 cm. Width: About 8 mm. Shape: Ovate; concave; very slightly reflexed. Apex: Bluntly acute. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Luster, upper surface: Matte. Luster, lower surface: Slightly glossy. Color: When opening and fully opened, upper surface: Close to 12A; color becoming closer to 14B with development. When opening and fully opened, lower surface: Close to 13B.

Sepals.—Quantity and arrangement: Five in a single whorl. Length: About 1 cm. Width: About 8 mm. Shape: Broadly ovate. Apex: Broadly acute to obtuse. Base: Broadly cuneate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Luster, upper and lower surfaces: Dull, matte. Color: When opening, upper surface: Close to 143A. When opening, lower surface: Close to 143B tinged with close to 180C. Fully opened, upper surface: Close to between 141A and 143A. Fully opened, lower surface: Close to 138B.

Peduncles.—Length: About 3.4 cm. Diameter: About 2 mm. Texture: Smooth, glabrous. Strength: Moderately strong. Aspect: Erect to about 40° from vertical. Color: Close to 144B to 144C.

Pedicels.—Length: About 1.3 cm. Diameter: About 1.5 mm. Texture: Smooth, glabrous. Strength: Moderately strong. Aspect: Erect to about 45° from vertical. Color: Close to 144B.

Reproductive organs.—Stamens: Quantity per flower: About 80. Filament length: About 1.3 mm. Filament color: Close to 12A. Anther length: About 0.5 mm. Anther shape: Broadly oval. Anther color: Close to 22A. Pollen amount: Scarce. Pollen color: Close to 11A. Pistils: Quantity per flower: Single pistil with three stigmas. Pistil length: About 4 mm. Stigma shape: Club-shaped. Stigma color: Close to 35A. Style length: About 3.5 mm. Style color: Close to 150C. Ovary color: Close to 150D.

Fruits.—Length: About 1.6 cm. Diameter: About 1.6 cm. Shape: Spherical. Texture: Smooth, glabrous. Luster: Glossy. Color: Slightly darker than between 60A and 185A; towards the base, close to 145D.

Seeds.—Length: About 0.9 mm. Diameter: About 0.5 mm. Color: Close to 200A.

Disease & pest resistance: Plants of the new *Hypericum* have not been observed to be resistant to pathogens and pests common to *Hypericum* plants.
Temperature tolerance: Plants of the new *Hypericum* have been observed to tolerate high temperatures about 30° C. 5
and to be hardy to USDA Hardiness Zone 7.

It is claimed:
1. A new and distinct *Hypericum* plant named ‘Alllangano’ as illustrated and described.
* * * * *

