



(12) **United States Plant Patent**  
**Zaiger et al.**

(10) **Patent No.:** **US PP27,188 P2**  
(45) **Date of Patent:** **Sep. 27, 2016**

(54) **INTERSPECIFIC TREE NAMED ‘REDONDO’**

(50) Latin Name: **Interspecific *Prunus* species**  
Varietal Denomination: **Redondo**

(71) Applicants: **Gary Neil Zaiger**, Modesto, CA (US);  
**Leith Marie Gardner**, Modesto, CA  
(US); **Grant Gene Zaiger**, Modesto,  
CA (US)

(72) Inventors: **Gary Neil Zaiger**, Modesto, CA (US);  
**Leith Marie Gardner**, Modesto, CA  
(US); **Grant Gene Zaiger**, Modesto,  
CA (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 37 days.

(21) Appl. No.: **14/545,068**

(22) Filed: **Mar. 23, 2015**

(51) **Int. Cl.**  
**A01H 5/08** (2006.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./180**

(58) **Field of Classification Search**  
USPC ..... **Plt./180**  
See application file for complete search history.

*Primary Examiner* — Anne Grunberg

(57) **ABSTRACT**

A new and distinct variety of interspecific tree. The follow-  
ing features of the tree and its fruit are characterized with the  
tree budded on ‘Nemaguard’ Rootstock (non-patented),  
grown on Handford sandy loam soil with Storie Index rating  
95, in USDA Hardiness Zone 9, near Modesto, Calif., with  
standard commercial fruit growing practices, such as prun-  
ing, thinning, spraying, irrigation and fertilization. Its nov-  
elty consist of the following combination of desirable fea-  
tures:

1. Tree having a vigorous, semi-spreading growth habit.
2. Tree being a regular and productive bearer of medium  
to large size fruit.
3. Fruit with an attractive orange skin color.
4. Fruit with very good flavor and eating quality.
5. Fruit with an average Brix of 20.1°.

**1 Drawing Sheet**

**1**

Botanical designation: Interspecific *Prunus* species.  
Variety denomination: ‘REDONDO’.

**BACKGROUND OF THE VARIETY**

**Field of the Invention**

In the field of plant genetics, we conduct an extensive and  
continuing plant-breeding program including the organiza-  
tion and asexual reproduction of orchard trees, and of which  
plums, peaches, nectarines, apricots, cherries, almonds and  
interspecifics are exemplary. It was against this background  
of our activities that the present variety of interspecific tree  
was originated and asexually reproduced by us in our  
experimental orchard located near Modesto, Stanislaus  
County, Calif.

**PRIOR VARIETIES**

Among the existing varieties of interspecific trees, which  
are known to us, and mentioned herein, ‘Late Brittney’  
Interspecific (U.S. Plant Pat. No. 18,921), ‘Summer Cot’  
Interspecific (U.S. Plant Pat. No. 23,864) and our propri-  
etary non-patented interspecific selections ‘15M39’,  
‘39ZK329’ and ‘13MA480’.

**STATEMENT REGARDING FEDERALLY  
SPONSORED RESEARCH AND  
DEVELOPMENT**

Not applicable.

**ORIGIN OF THE VARIETY**

The new variety of interspecific tree, a combination of  
crosses between *Prunus armeniaca*, *Prunus salicina* and

**2**

*Prunus persica* was originated by us in our experimental  
orchard located near Modesto, Calif. from crosses between  
the proprietary non-patented interspecific varieties with the  
field identification numbers ‘15M39’ and ‘39ZK329’. The  
seed parent (15M39) originated as an open pollinated seed-  
ling selection of ‘Late Brittney’ Interspecific (U.S. Plant Pat.  
No. 18,921). The pollen parent (39ZK329) originated as an  
open pollinated seedling selection of the proprietary non-  
patented interspecific seedling ‘13MA480’. A large number  
of these first generation seedlings were budded onto older  
established trees of ‘Nemaguard’ Rootstock (non-patented)  
to enhance earlier fruit production. Under close and careful  
observation the present seedling exhibited desirable fruit and  
tree characteristics and was selected in 2010 for additional  
asexual propagation and commercialization.

**ASEXUAL REPRODUCTION OF THE VARIETY**

In 2010 asexual reproduction of the new and distinct  
variety of interspecific tree was by budding to ‘Nemaguard’  
Rootstock (non-patented), as performed by us in our experi-  
mental orchard located near Modesto, Calif., and shows that  
reproductions run true to the original tree and all character-  
istics of the tree and its fruit are established and transmitted  
through succeeding asexual propagations.

**SUMMARY OF THE NEW VARIETY**

The new and distinct variety of interspecific tree Apricot×  
Plum×Peach is of large size, vigorous, semi-spreading  
growth and a regular and productive bearer of medium to  
large size, orange flesh, freestone fruit. The fruit is further  
characterized by having an attractive orange skin color, very  
good flavor and eating quality and having good storage and



shipping quality. In comparison to its non-patented interspecific seed parent '15M39' the fruit of the new variety is approximately 8 days later in maturity. In comparison to its non-patented interspecific pollen parent '39ZK329' the fruit of the new variety is approximately 24 days later in maturity. In comparison to the commercial variety 'Summer Cot' Interspecific (U.S. Plant Pat. No. 23,864) the fruit of the new variety is approximately 28 days earlier in maturity.

#### DESCRIPTION OF THE PHOTOGRAPH

The accompanying color photographic illustration shows typical specimens of the foliage and fruit of the present new interspecific variety.

The illustration shows the upper and lower surface of the leaves, an exterior and sectional view of a single fruit divided in its suture plane to show flesh color, pit cavity and the stone remaining in place:

The photographic illustration was taken shortly after being picked (shipping ripe) from a 5 year old tree and the colors are as nearly true as is reasonably possible in a color representation of this type.

#### DESCRIPTION OF THE VARIETY

The following is a detailed botanical description of the new variety of interspecific tree, its flowers, foliage and fruit, as based on observations of 5 year old specimens grown near Modesto, Calif., with color in accordance with Munsell Book of Color published in 1958.

##### Tree:

*Size*.—Large, usually pruned to 3 to 3.5 meters in height and width for economical harvesting of fruit. Varies with different cultural practices.

*Vigor*.—Vigorous, tree growth of approximately 1.5 to 2 meters the first growing season. Varies with cultural practice, soil type, fertility of soil and climatic conditions.

*Form*.—Semi-spreading, usually pruned to vase shape.

*Branching habit*.—Semi-spreading, crotch angle approximately 35°, increases with heavy crop load.

*Productivity*.—Productive, thinning and spacing of fruit necessary for desired market size. Fruit set varies with climatic conditions during bloom time.

*Bearer*.—Regular, adequate fruit set 3 consecutive years. No alternate bearing observed.

*Fertility*.—Self fertile, sets fruit under bag.

*Density*.—Medium dense, usually pruned to vase shape to increase air movement and sunlight to center of tree to enhance fruit color and health of fruit spurs.

*Hardiness*.—Hardy in all stone fruit growing areas of California. Tree grown in USDA Hardiness Zone 9. Winter chilling requirement approximately 900 hours at or below 45° F.

##### Trunk:

*Size*.—Medium to large. Average circumference 59.7 cm at 25.4 cm above ground on a 5 year old tree.

*Stocky*.—Medium stocky.

*Texture*.—Medium shaggy, roughness increases with age.

*Color*.—Varies from 2.5Y 5/2 to 2.5Y 4/2.

##### Branches:

*Size*.—Medium. Average circumference 13.7 cm at 1.2 meters above ground on a 5 year old tree. Crotch angle approximately 35°, increases with heavy crop load.

*Surface texture*.—New growth relatively smooth. Mature growth medium rough, roughness increases with age.

*Lenticels*.—Average number 41 in a 25.8 square cm area. Average length 3.1 mm. Average width 1.5 mm. Color varies from 2.5Y 7/8 to 2.5Y 6/8.

*Color*.—New growth varies from 5GY 5/8 to 7.5R 2/8. Mature growth varies from 7.5YR 2/4 to 10YR 3/4, varies with age of growth.

##### Leaves:

*Size*.—Small to medium. Average length 72.3 mm. Average width 57.5 mm.

*Form*.—Ovate.

*Apex*.—Acuminate.

*Base*.—Cuneate.

*Margin*.—Doubly serrate.

*Thickness*.—Medium.

*Surface texture*.—Upper surface relatively smooth, slight indentations over midrib and leaf veins. Lower surface relatively smooth, except for small ridges created by midrib and pinnate venation. Both upper and lower surfaces glabrous.

*Petiole*.—Medium. Average length 24.5 mm. Average width 1.3 mm. Longitudinally grooved. Surface — glabrous. Color varies from 2.5GY 6/6 to 7.5R 3/4.

*Glands*.—Type — reniform. Size — medium. Average length 1.1 mm. Average diameter 0.9 mm. Average number 3, varies from 2 to 4. Located primarily on the upper portion of the petiole and base of leaf blade. Color varies from 7.5R 3/4 to 7.5R 2/4.

*Stipules*.—Average length 7.7 mm. Average number 2. Edges — pectinate. Color varies from 2.5GY 6/6 to 7.5R 3/4.

*Color*.—Upper surface varies from 5GY 4/6 to 5GY 3/6. Lower surface varies from 2.5GY 4/4 to 5GY 4/4. Midvein color varies from 2.5GY 6/6 to 5GY 6/6.

##### Flower buds:

*Size*.—Large. Average length 14.3 mm. Average diameter 8.4 mm.

*Hardiness*.—Hardy with respect to California winters.

*Form*.—Conical, becoming elongated just before opening.

*Pedicel*.—Average length 1.8 mm. Average width 1.2 mm. Color varies from 2.5GY 6/6 to 5GY 7/6. Surface — glabrous.

*Color*.—Varies from 7.5RP 9/2 to 7.5RP 7/8.

*Number of buds per spur*.—Varies from 7 to 13, average 9. Varies with age of spur.

##### Flowers:

*Blooming period*.—Date of First Bloom Feb. 24, 2014. Date of Petal Fall Mar. 5, 2014, varies slightly with climatic conditions.

*Size*.—Medium to large. Average height 15.8 mm. Average diameter 22.3 mm.

*Petals*.—Normally 5, alternately arranged to sepals. Size — large. Average length 12.0 mm. Average width 12.2 mm. Shape — orbicular. Arrangement — overlapping. Petal apex — rounded. Petal base — rounded to somewhat truncated. Margin — sinuate. Color varies from 5RP 9/2 to 7.5RP 9/2, fades with age of flower. Both upper and lower surfaces glabrous.

*Sepals*.—Normally 5, alternately arranged to petals. Size — large. Average length 5.4 mm. Average width



5.0 mm. Shape — ovate. Margin — entire. Apex — rounded to triangular. Both upper and lower surfaces glabrous. Color — upper surface varies from 2.5R 2/6 to 5R 2/8. Lower surface varies from 5R 2/4 to 5R 2/6.

*Stamens*.—Average number per flower 33. Average filament length 10.5 mm. On average, the stamens are even with the height of the petals. Filament color N 9.5/ (white). Anther color varies from 5Y 8/8 to 5Y 8/10.

*Pollen*.—Self fertile, sets fruit under bag. Color varies from 2.5Y 7/10 to 5Y 7/8.

*Pistil*.—Number — normally one. Surface — pubescent. Average length 14.0 mm. Position of stigma even with anthers. Color varies from 7.5Y 8/6 to 10Y 8/6.

*Fragrance*.—Heavy.

*Color*.—Varies from 7.5RP 9/2 to 10RP 9/2, depending on age of flower.

*Pedicel*.—Average length 2.5 mm. Average width 1.4 mm. Color varies from 10Y 7/6 to 2.5GY 7/6. Surface — glabrous.

*Number flowers per flower bud*.—Average 2, varies from 1 to 3.

#### Fruit:

*Maturity when described*.—Firm ripe and ready for consumption.

*Date of first picking*.—Jul. 17, 2014.

*Date of last picking*.—Jul. 27, 2014, varies slightly with climatic conditions.

*Size*.—Medium to large. Average diameter axially 56.1 mm. Average 63.7 mm transversely in suture plane. Average across suture plane 57.3 mm. Average weight 119.3 grams, varies slightly with fertility of the soil, amount of thinning and climatic conditions.

*Form*.—Globose to slightly elongated.

*Suture*.—Slightly lipped, extends from base toward apex.

*Ventral surface*.—Slightly lipped.

*Apex*.—Retuse.

*Base*.—Retuse.

*Stem cavity*.—Rounded to slightly elongated in suture plane. Average depth 8.8 mm. Average diameter 4.5 mm.

#### Stem:

*Size*.—Small. Average length 4.9 mm. Average diameter 3.0 mm.

*Color*.—Varies from 2.5GY 6/6 to 2.5GY 6/8.

#### Flesh:

*Ripens*.—Evenly.

*Texture*.—Firm, meaty.

*Fibers*.—Few, small, tender.

*Firmness*.—Firm, comparable to other commercial apricot varieties.

*Aroma*.—Moderate.

*Amydgalin*.—Undetected.

*Eating quality*.—Very good.

*Flavor*.—Very good, with a good balance between acid and sugar.

*Juice*.—Moderate amount, enhances flavor.

*Acidity*.—Not available.

*Brix*.—Average Brix 20.1°, varies slightly with amount of fruit per tree and climatic conditions.

*Color*.—Varies from 7.5YR 7/12 to 10YR 8/10.

*Pit cavity*.—Average length 32.0 mm. Average width 26.9 mm. Average depth 9.3 mm. Color varies from 7.5YR 7/12 to 7.5YR 7/14.

#### Skin:

*Thickness*.—Medium.

*Surface*.—Smooth to very slightly waffled.

*Pubescence*.—Light, short in length.

*Tendency to crack*.—None.

*Color*.—Color varies from 7.5YR 7/10 to 10YR 7/10.

*Tenacity*.—Tenacious to flesh.

*Astringency*.—Undetected.

#### Stone:

*Type*.—Freestone, weak adherence.

*Size*.—Large. Average length 29.2 mm. Average width 24.9 mm. Average thickness 14.5 mm.

*Form*.—Ovoid.

*Base*.—Flat.

*Apex*.—Rounded.

*Surface*.—Lightly pitted throughout. One shallow groove on each side of suture extending from base toward apex.

*Sides*.—Unequal, one side extending further outward from suture plane.

*Ridges*.—Very small and short, extending from base towards apex.

*Tendency to split*.—None.

*Color*.—Varies from 10YR 4/2 to 10YR 3/2 when dry.

#### Kernel:

*Size*.—Large. Average length 18.7 mm. Average width 15.1 mm. Average depth 7.6 mm.

*Form*.—Ovoid.

*Viability*.—Viable, complete embryo development.

*Skin color*.—Varies from 5Y 9/2 to 10YR 6/4.

#### Use:

*Dessert*.—Market — local and long distance.

Keeping quality: Good, held firm in cold storage for 3 weeks at 38° to 42° F. without shriveling, internal breakdown of flesh or appreciable loss of eating quality.

Shipping quality: Good, showed minimal skin scarring or flesh bruising during picking, packing and shipping trials.

Plant/fruit disease resistance/susceptibility: No specific testing for relative plant/fruit disease resistance/susceptibility has been designed. Under close observation during planting, growing and harvesting of fruit, under normal cultural and growing conditions near Modesto, Calif., no particular plant/fruit disease resistance or susceptibility has been observed. Any variety observed during indexing of plant characteristics with abnormal fungus, bacterial, virus or insect susceptibility is destroyed and eliminated from our breeding program. No atypical resistances/susceptibilities have been noted under normal cultural practices. The present new variety of interspecific tree, its flowers foliage and fruit herein described may vary in slight detail due to climate, soil conditions and cultural practices under which the variety may be grown. The present description is that of the variety grown under the ecological conditions prevailing near Modesto, Calif.

The invention claimed is:

1. A new and distinct variety of interspecific tree, substantially as illustrated and described.

\* \* \* \* \*



