



(12) **United States Plant Patent**
Uchneat

(10) **Patent No.:** **US PP27,474 P2**
(45) **Date of Patent:** **Dec. 20, 2016**

(54) **CRAPEMYRTLE PLANT NAMED**
‘G2X133271’

(50) Latin Name: *Lagerstroemia indica*
Varietal Denomination: **G2X133271**

(71) Applicant: **Michael S. Uchneat**, Bellefonte, PA
(US)

(72) Inventor: **Michael S. Uchneat**, Bellefonte, PA
(US)

(73) Assignee: **GardenGenetics LLC**, Bellefonte, PA
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/545,625**

(22) Filed: **May 29, 2015**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./252**

(58) **Field of Classification Search**
USPC **Plt./252**
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

PLUTO Plant Variety Database Jul. 19, 2016. p. 1.*

* cited by examiner

Primary Examiner — Annette Para

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of Crapemyrtle plant named
‘G2X133271’, characterized by its relatively compact,
upright to outwardly spreading plant habit; freely branching
habit; dark green-colored leaves; numerous inflorescences
with rose magenta-colored flowers; and good garden per-
formance.

1 Drawing Sheet

1

Botanical designation: *Lagerstroemia indica*.
Cultivar denomination: ‘G2X133271’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of Crapemyrtle plant, botanically known as *Lagerstroemia*
indica and hereinafter referred to by the name
‘G2X133271’.

The new Crapemyrtle plant is a product of a planned
breeding program conducted by the Inventor in Bellefonte,
Pa. The objective of the breeding program was to develop
new compact, mounding and freely-branching Crapemyrtle
plants with large inflorescences, high temperature tolerance,
winter hardiness and resistance to Powdery Mildew.

The new Crapemyrtle plant originated from an open-
pollination conducted in August, 2010 of *Lagerstroemia*
indica ‘Red Filli’, disclosed in U.S. Plant Pat. No. 14,353,
as the female, or seed, parent with an unknown selection of
Lagerstroemia indica as the male, or pollen, parent. The new
Crapemyrtle plant was discovered and selected by the Inven-
tor on Jan. 9, 2011 as a single flowering plant from within
the progeny of the stated open-pollination in a controlled
nursery environment in Bellefonte, Pa.

Asexual reproduction of the new Crapemyrtle plant by
vegetative tip cuttings in a controlled greenhouse environ-
ment in Bellefonte, Pa. since Feb. 2, 2011 has shown that the
unique features of the new Crapemyrtle plant are stable and
reproduced true to type in successive generations of asexual
reproduction.

SUMMARY OF THE INVENTION

Plants of the new Crapemyrtle have not been observed
under all possible combinations of environmental conditions

2

and cultural practices. The phenotype may vary somewhat
with variations in environmental conditions such as tem-
perature and light intensity without, however, any variance
in genotype.

5 The following traits have been repeatedly observed and
are determined to be the unique characteristics of
‘G2X133271’. These characteristics in combination distin-
guish ‘G2X133271’ as a new and distinct Crapemyrtle plant:

- 10 1. Relatively compact, upright to outwardly spreading and
arching plant habit.
2. Freely branching habit.
3. Dark green-colored leaves.
4. Numerous inflorescences with rose magenta-colored
flowers.
- 15 5. Good garden performance.

Plants of the new Crapemyrtle can be compared to plants
of the female parent, ‘Red Filli’. Plants of the new Crape-
myrtle differ primarily from plants of ‘Red Filli’ in the
following characteristics:

- 20 1. Plants of the new Crapemyrtle are more freely branch-
ing than plants of ‘Red Filli’.
2. Plants of the new Crapemyrtle have rose magenta-
colored flowers whereas plants of ‘Red Filli’ have
bright red-colored flowers.
- 25

Plants of the new Crapemyrtle can be compared to plants
of the *Lagerstroemia indica* ‘Tonto’, not patented. In side-
by-side comparisons conducted in Bellefonte, Pa., plants of
the new Crapemyrtle differed primarily from plants of
‘Tonto’ in the following characteristics:

- 30 1. Plants of the new Crapemyrtle were more compact than
plants of ‘Tonto’.
2. Plants of the new Crapemyrtle were more freely
branching than plants of ‘Tonto’.

3. Flowers of plants of the new Crapemyrtle were rose magenta in color whereas flowers of plants of 'Tonto' were fuchsia in color.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new Crapemyrtle plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Crapemyrtle plant.

The photograph is a side perspective view of a typical plant of 'G2X133271' grown in a ground bed in an outdoor nursery.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown in Grand Haven, Mich. during the summer in a polypropylene-covered shadehouse and under cultural conditions which closely approximate commercial Crapemyrtle production. During the production of the plants, day temperatures ranged from 18° C. to 27° C. and night temperatures ranged from 5° C. to 10° C. Plants were two years when the photographs and the description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. Botanical classification: *Lagerstroemia indica* 'G2X133271'.

Parentage:

Female, or seed, parent.—*Lagerstroemia indica* 'Red Filli', disclosed in U.S. Plant Pat. No. 14,353.

Male, or pollen, parent.—Unknown selection of *Lagerstroemia indica*, not patented.

Propagation:

Type.—By vegetative tip cuttings.

Time to initiate roots, summer and winter.—About two to three weeks at temperatures about 26° C.

Time to produce a rooted young plant, summer.—About four to five weeks at temperatures about 23° C.

Time to produce a rooted young plant, winter.—About four to five weeks at temperatures about 19° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching; medium density.

Plant description:

Plant form and growth habit.—Perennial shrub; relatively compact, upright to outwardly spreading and arching plant habit; vigorous growth habit.

Branching habit.—Freely branching habit with about 40 lateral branches developing per plant.

Plant height.—About 52.5 cm.

Plant diameter (area of spread).—About 111 cm.

Lateral branch description:

Length.—About 70 cm.

Diameter.—About 1 cm.

Internode length.—About 1 cm.

Strength.—Strong.

Aspect.—About 15° to 40° from vertical.

Texture.—Smooth, glabrous; woody with age.

Color, immature.—Close to 183B.

Color, mature.—Close to 183B and 177A.

Leaf description:

Arrangement.—Alternate to sub-opposite; simple.

Length.—About 5 cm.

Width.—About 3 cm.

Shape.—Ovate.

Apex.—Acute.

Base.—Cordate.

Margin.—Entire.

Texture, upper surface.—Smooth to slightly ribbed, glabrous.

Texture, lower surface.—Venation is prominent, ribbed.

Venation pattern.—Pinnate.

Color.—Developing leaves, upper and lower surfaces: Close to 144B and 185A. Fully expanded leaves, upper surface: Close to 137A; venation, close to 138B. Fully expanded leaves, lower surface: Close to 138B; venation, close to 138B.

Petioles.—Length: About 1 mm. Diameter: About 1 mm. Texture, upper surface: Slightly pubescent; coriaceous. Texture, lower surface: Smooth, glabrous; coriaceous. Color, upper and lower surfaces: Close to 143A.

Flower description:

Flower type, arrangement and habit.—Showy single ruffled flowers arranged in terminal panicles; freely flowering habit with usually about twelve flowers per inflorescence and numerous inflorescences developing during the flowering season; flowers face upright and outwardly; flowers not persistent.

Natural flowering season.—Plants of the new Crapemyrtle flower from June through October in Grand Haven, Mich.

Fragrance.—None detected.

Inflorescence height.—About 12.8 cm.

Inflorescence diameter.—About 9.4 cm.

Flower length.—About 4.6 cm.

Flower diameter.—About 4.5 cm.

Flower depth.—About 1.3 cm.

Flower buds.—Length: About 8 mm. Diameter: About 7.2 mm. Shape: Obovate. Texture: Smooth, glabrous. Color: Close to 145B overlain with close to 184A.

Petals.—Quantity per flower and arrangement: Six arranged in a single whorl. Length: About 2 cm. Width: About 1.4 cm. Shape: Roughly orbicular. Apex: Rounded, undulate. Base: Rounded. Margin: Entire, undulate. Texture, upper and lower surfaces: Smooth, glabrous; soft; delicate. Color: When opening, upper and lower surfaces: Close to 63A. Fully opened, upper and lower surfaces: Close to 58A; color becoming closer to 63A slightly shaded with close to 58D with development.

Sepals.—Quantity per flower and arrangement: Six arranged in a single whorl. Length: About 9 mm. Width: About 3.2 mm. Shape: Narrowly deltoid. Apex: Acute. Base: Obtuse. Margin: Entire. Texture, upper surface: Rugose, glabrous; coriaceous. Texture, lower surface: Smooth, glabrous. Color: When opening and fully opened, upper surface: Close to 144B overlain with close to 176B. When opening and fully opened, lower surface: Close to 145C and 155C.

Pedicels.—Length: About 1.3 cm. Diameter: About 8 mm. Strength: Strong, flexible. Aspect: About 5° to

35° from stem axis. Texture: Slightly pubescent. Color: Close to 145B; striations, tinged with close to 176B.

Reproductive organs.—Androecium: Quantity per flower: About 29. Filament length: About 1 cm. 5
Filament color: Close to 70D; towards the base, blended with close to 155B. Anther length: About 1 mm. Anther shape: Lanceolate. Anther color: Close to 163A; edges, close to 165A. Amount of pollen: Scarce. Pollen color: Close to 163A. Gynoecium: 10
Quantity per flower: One. Pistil length: About 2 cm. Style length: About 1.5 cm. Style color: Close to 184D. Stigma appearance: Spherical. Stigma color: Close to 189A. Ovary: Close to 11B.

Fruits and seeds.—Fruit and seed development has not been observed on plants of the new Crapemyrtle.

Garden performance: Plants of the new Crapemyrtle have been observed to have good garden performance and to tolerate rain, wind and temperatures ranging from about -28° C. to about 35° C.

Pathogen & pest resistance: Plants of the new Crapemyrtle have not been observed to be resistant to pathogens and pests common to Crapemyrtle plants.

It is claimed:

1. A new and distinct Crapemyrtle plant named 'G2X133271' as illustrated and described.

* * * * *

