



US00PP27586P3

(12) **United States Plant Patent**
Adams

(10) **Patent No.:** **US PP27,586 P3**
(45) **Date of Patent:** **Jan. 24, 2017**

- (54) **RASPBERRY PLANT NAMED ‘GLEAM’**
- (50) Latin Name: *Rubus idaeus*
Varietal Denomination: **Gleam**
- (71) Applicant: **PLANT SCIENCES, INC.,**
Watsonville, CA (US)
- (72) Inventor: **Scott W. Adams**, Watsonville, CA (US)
- (73) Assignee: **Plant Sciences, Inc.**, Watsonville, CA (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 175 days.
- (21) Appl. No.: **14/121,145**
- (22) Filed: **Aug. 6, 2014**
- (65) **Prior Publication Data**
US 2015/0223380 P1 Aug. 6, 2015
- Related U.S. Application Data**
- (60) Provisional application No. 61/934,890, filed on Feb. 3, 2014.

- (51) **Int. Cl.**
A01H 5/08 (2006.01)
- (52) **U.S. Cl.**
USPC **Plt./204**
- (58) **Field of Classification Search**
USPC Plt./204
See application file for complete search history.

Primary Examiner — Keith Robinson
(74) *Attorney, Agent, or Firm* — Foley & Lardner LLP

(57) **ABSTRACT**
This invention relates to a new and distinct everbearing variety of raspberry plant named ‘GLEAM’. The new variety is primarily adapted to the growing conditions of the central coast of California and is characterized by the following: early primocane production with medium to large sized fruit of light-red coloration. Fruit is of consistent conic shape, releases easily from receptacle and is of very good flavor. Foliage is strongly concave, medium green; possessing weak rugosity and mostly 3 foliate. Primocanes have an absent or very weak waxy coat, medium thorn density and have very weak anthocyanin coloration.

4 Drawing Sheets

1

Latin name of the genus and species of the plant claimed: *Rubus idaeus*.
Variety denomination: ‘GLEAM’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct fall bearing raspberry variety designated as ‘GLEAM’. This new variety is a result of a controlled cross made in 2005 in Watsonville, Calif. between raspberry variety ‘PS-1852’ (patented, U.S. Plant Pat. No. 18,806) as the female parent and raspberry variety ‘RADIANCE’ (patented, U.S. Plant Pat. No. 20,342) as the male parent in an ongoing breeding program. The variety is botanically known as *Rubus idaeus*.

The seedling resulting from the aforementioned cross was asexually propagated by dormant canes in Santa Cruz County, Calif. and was subsequently selected by the inventor from a controlled breeding plot in Watsonville, Calif. in 2007. After its selection, the new variety was further asexually propagated by dormant canes, roots and non-dormant root shoot cuttings in Santa Cruz County, Calif., San Joaquin County, Calif. and Siskiyou County, Calif. The new variety was then extensively tested over the next several years in fruiting fields in Santa Cruz County, Calif. This propagation has demonstrated that the combination of traits disclosed herein as characterizing the new variety are fixed and remain true to type through successive generations of asexual reproduction.

BRIEF SUMMARY OF THE INVENTION

‘GLEAM’ is primarily adapted to the climate and growing conditions of the central coast of California. This region

2

provides the necessary year-round temperatures required for it to produce and maintain a strong vigorous plant and to remain in fruit production from July through December on primocanes and in the ensuing year from May through July on the floricanes. The following traits have been repeatedly observed and are determined to be unique characteristics of ‘GLEAM’, which in combination distinguish this raspberry plant as a new and distinct variety:

1. Medium fruit size
2. High fruit yield
3. Light to medium red fruit color
4. Absent to very weak primocane glaucosity
5. Very good flavor

The raspberry variety that is believed to be most closely related to the new raspberry variety ‘GLEAM’ is the raspberry variety ‘GRANDEUR’ (patented, U.S. Plant Pat. No. 20,459). In comparison to the similar raspberry variety ‘GRANDEUR’, ‘GLEAM’ differs by the following combination of characteristics described in Table 1:

TABLE 1

Characteristic	‘GLEAM’	‘GRANDEUR’ (U.S. Plant Pat. No. 20,459)
Primocane waxy coat (glaucosity)	Absent to very weak	Absent to very weak
Predominate number of leaflets	Mostly 3	Always 3
Leaf shape (cross section)	Strongly concave	Flat to slightly convex
Rugosity	Weak	Strong
Color mature fruit	RHS 43A Light to medium red	RHS 42A Medium red

TABLE 1-continued

Characteristic	'GLEAM'	'GRANDEUR' (U.S. Plant Pat. No. 20,459)
Primocane fruit weight (g)	3.8	3.9
Florican color	RHS 165B Greyed-orange group	RHS 164A Greyed-orange group

'GLEAM' differs from its parents, 'PS-1852' and 'RADIANCE' by the following combination of characteristics described in Table 2:

TABLE 2

Characteristic	'GLEAM'	'PS-1852' (U.S. Plant Pat. No. 18,806)	'RADIANCE' (U.S. Plant Pat. No. 20,342)
Productivity	High	Medium to low	Medium
Glossiness (fruit)	Medium	Medium	High
Adherence of receptacle	Weak	Medium	Weak
Primocane time of fruiting	Early	Late	Medium
Fruit size	Medium	Medium	Small
Flavor	Very good	Good	Very good

For identification, a series of molecular markers have been determined for this new variety.

BRIEF DESCRIPTIONS OF THE DRAWINGS

The accompanying color photographs illustrate the overall appearance of typical specimens of the new raspberry variety, 'GLEAM' at various stages of development as true as reasonably possible with color reproductions of this type. Color in the photographs may differ slightly from the color value cited in the detailed botanical description which accurately describes the color of 'GLEAM'. The depicted plant and plant parts of the new raspberry variety 'GLEAM' were taken in Watsonville, Calif. and are approximately 2 to 16 months old:

FIG. 1 shows typical primocane foliage and fruit color; foliate and rugosity characteristics of 'GLEAM' taken in the month of July 2014;

FIG. 2 shows typical coloration of apical growing tip during early primocane rapid growth of 'GLEAM' taken in the month of May 2013;

FIG. 3 shows typical harvested fruit of 'GLEAM' taken in the month of August 2013;

FIG. 4 shows typical dormant cane color characteristics of 'GLEAM' taken in the month of January 2013.

DETAILED BOTANICAL DESCRIPTION

'GLEAM' has not been observed under all possible environmental conditions. The characteristics of the new variety may vary in detail, depending upon variations in environmental factors, including weather (temperature, humidity and light intensity), day length, soil type, and location.

The aforementioned photographs, together with the following description of the new raspberry variety 'GLEAM', unless otherwise noted, are based upon observations taken during the 2013-2014 growing season in Watsonville, Calif. Primocane measurements and ratings were taken from plants of 'GLEAM' dug from a nursery located in Siskiyou County, Calif. during the middle of October 2012 and planted approximately 3 to 4 weeks later in Watsonville, Calif. The approximate age of the observed primocane plants is 7 to 8 months. Florican measurements and ratings were taken from the same planting of 'GLEAM' at an approximate age

of 16 to 18 months. Yield observations and fruit quality characteristics are averaged from five years of data collected from the 2009 through 2013 production seasons. Flower measurements and characteristics are from secondary flowers unless otherwise noted. Fruit characteristics and measurements are from secondary fruit unless otherwise noted. Foliage characteristics and measurements are from 3-foliolate foliage unless otherwise noted.

Color terminology where noted follows The R.H.S. Colour Chart Fifth Edition, Royal Horticultural Society, London, United Kingdom (1966).

The following tables 3-7 describe fruit, plant, foliage, flower and pest/disease characteristics of the new raspberry 'GLEAM' in comparison to the similar raspberry varieties 'GRANDEUR' (patented, U.S. Plant Pat. No. 20,459).

TABLE 3

FRUIT CHARACTERISTICS		
Characteristic	'GLEAM'	'GRANDEUR' (U.S. Plant Pat. No. 20,459)
Color mature fruit	RHS 43A Light to medium red	RHS 42A Medium red
Color achenes	RHS 159A Orange-white group	RHS 159A Orange-white group
Fruit length (mm)	21.74	22.91
Fruit width (mm)	20.65	20.12
Length/Width ratio	1.05	1.14
	Longer than broad	Longer than broad
Seed weight (mg)	1.73	1.75
Drupelets per berry	86	93
Weight of single drupe (g/drupe)	0.047	0.043
Relative size of drupes	Medium	Medium
Fruit Size	Medium	Medium
Predominant shape	Conical	Conical
Evenness of color	Even	Even
Glossiness	Medium	Medium
Adherence of receptacle	Weak	Weak
Firmness of flesh	Firm	Very firm
Firmness of skin	Medium	Very firm
Soluble Solids (% brix)	10.6	10.5
Flavor	Very Good	Good

TABLE 4

PLANT CHARACTERISTICS		
Characteristic	'GLEAM'	'GRANDEUR' (U.S. Plant Pat. No. 20,459)
General:		
Habit	Upright	Upright
Size	Medium	Medium
Productivity	High	High
Self-fruitfulness	Yes	Yes
Type of bearing	Everbearing	Everbearing
Primocane:		
Color (true)	RHS 145B Yellow-green group	RHS 145B Yellow-green group
Length (cm)	147.3	143.6
Basal diameter (mm)	16.53	21.80
Diameter central 1/3 (mm)	9.04	12.55
Lateral length at central 1/3 (cm)	48.0	43.6
No. fruiting laterals per cane	10.8	17.4
Internode length at central 1/3 (mm)	71.35	48.84

TABLE 4-continued

PLANT CHARACTERISTICS		
Characteristic	'GLEAM'	'GRANDEUR' (U.S. Plant Pat. No. 20,459)
Anthocyanin coloration	RHS 59B Red-purple group	RHS 59B Red-purple group
Anthocyanin intensity	Absent to very weak	Weak
Pubescence	Absent	Absent
Length of vegetative bud (mm)	6.82	11.01
Strength of waxy coat (glaucosity)	Absent to very weak	Absent to very weak
Time of flowering	Early	Medium
Time of fruiting	Early	Medium
Length of fruiting season	Long	Long
% of total yield	52%	50%
Flowering period	Late June to Late November	Late June to Late November
Harvest period	Late July to Late December	Late July to Late December
Primocane fruit weight (g)	3.8	3.9
Primocane yield (g/plant)	2,429	2,662
Young Shoots:		
Number	Medium	Medium
Anthocyanin presence	Present	Present
Anthocyanin coloration	RHS 59B Red-purple group	RHS 179A Greyed-red group
Anthocyanin intensity	Medium	Medium
Thorns:		
Thorn coloration	RHS 182A Greyed-red group	RHS 180C Greyed-red group
Thorn length at central 1/3 (mm)	0.55	0.76
Thorn base at central 1/3 (mm)	0.94	1.53
Thorn presence	Present	Present
Thorn density per cm at central 13	3.94	3.38
Thorn texture	Medium	Medium
Thorn attitude of the tip	Rigid	Rigid
Floricane:	Horizontal	Horizontal
Color (true)	RHS 165B Greyed-orange group	RHS 164A Greyed-orange group
Length (cm)	104.1	119.8
Fruiting lateral attitude	Erect	Erect
Time bud burst	Medium	Medium
Time of flowering	Medium	Medium
Time of fruiting	Medium	Medium
Length of fruiting season	Medium to long	Medium to long
% of total yield	48%	50%
Flowering period	Late April to Late June	Late April to Late June
Harvest period	Late May to Late July	Late May to Late July
Floricane fruit weight (g)	3.6	3.7
Floricane yield (g/plant)	2,234	2,680

TABLE 5

FOLIAGE CHARACTERISTICS		
Characteristic	'GLEAM' (3 Foliate)	'GRANDEUR' (U.S. Plant Pat. No. 20,459) (3 Foliate)
General:		
Color of upper surface	RHS 137B Green group	RHS N137A Green group
Color of lower surface	RHS 190B Greyed-green group	RHS 190B Greyed-green group

TABLE 5-continued

FOLIAGE CHARACTERISTICS		
Characteristic	'GLEAM' (3 Foliate)	'GRANDEUR' (U.S. Plant Pat. No. 20,459) (3 Foliate)
Shape in cross section	Strongly concave	Flat to slightly convex
Arrangement	Compound	Compound
Relief between veins (rugosity)	Weak	Strong
Glossiness	Weak	Weak
Number of leaflets/leaf	Mostly 3	Always 3
Terminal Leaflet:		
Length (mm)	126.3	127.5
Width (mm)	76.7	91.6
Length/Width Ratio	1.6	1.4
Size	Longer than broad	Longer than broad
Shape	Medium	Medium to large
Shape of base	Ovate	Cordate
Shape of tip	Rounded	Cordate
Margins	Acuminate	Acuminate
Lateral Leaflet:		
Length (mm)	Biserrate	Biserrate
Width (mm)	103.5	105.5
Length/Width Ratio	59.4	65.1
Length/Width Ratio	1.7	1.6
Rachis length (mm)	Longer than broad	Longer than broad
Orientation	33.4	36.2
Arrangement	Opposite	Opposite
Shape	Compound	Compound
Overlapping	Ovate	Ovate
Shape of the base	Free	Touching
Shape of the tip	Rounded	Oblique rounded
Margins	Acuminate	Acuminate
Petiole:	Biserrate	Biserrate
Length (mm)	62.98	59.00
Width (mm)	2.82	3.99
Thorn presence	Yes	Yes
Thorn orientation	Erect	Erect
Anthocyanin coloration of upper surface	RHS 186C	RHS 184C
Anthocyanin intensity of upper surface	Greyed-purple group	Greyed-purple group
Stipule length (mm)	Weak	Weak
Stipule orientation	9.72	10.23
	Erect	Erect to horizontal

TABLE 6

FLOWER CHARACTERISTICS		
Characteristic	'GLEAM'	'GRANDEUR' (U.S. Plant Pat. No. 20,459)
Petal color	155C White group	155C White group
Flower diameter (mm)	18.14	22.55
Petal Length (mm)	6.79	6.42
Petal width (mm)	3.27	3.11
Petal length/width ratio	2.08	2.06
No. petals/flower	Much longer than broad	Much longer than broad
No. sepals/flower	5.4	5.2
Relative number of pedicel thorns	5.4	5.2
Peduncle anthocyanin presence	Medium	Medium
Peduncle anthocyanin coloration	Present	Present
Peduncle anthocyanin intensity	RHS 183B Greyed-purple group	RHS 184A Greyed-purple group
	Strong	Medium

TABLE 7

PEST AND DISEASE REACTIONS		
Characteristic	'GLEAM'	'GRANDEUR' (U.S. Plant Pat. No. 20,459)
Spotted wing drosophila (<i>Drosophila suzukii</i>)	Susceptible	Susceptible
Two spotted spider mite (<i>Tetranychus urticae</i>)	Susceptible	Susceptible
Grey fruit mold (<i>Botrytis cinerae</i>)	Susceptible	Susceptible
Powdery mildew (<i>Podosphaera aphanis</i> var. <i>aphanis</i>)	Moderately susceptible	Moderately susceptible

TABLE 7-continued

PEST AND DISEASE REACTIONS		
Characteristic	'GLEAM'	'GRANDEUR' (U.S. Plant Pat. No. 20,459)
Yellow rust (<i>Phragmidium rubi-idaei</i>)	Moderately resistant	Moderately susceptible

5

10

15

I claim:

1. A new and distinct variety of raspberry plant named 'GLEAM', as herein described and illustrated by the characteristics set forth above.

* * * * *

FIG. 1



FIG. 2




FIG. 3



FIG. 4

