

US00PP27744P3

(12) **United States Plant Patent**  
**Grosser**(10) **Patent No.:** US PP27,744 P3  
(45) **Date of Patent:** Mar. 7, 2017

- (54) **CITRUS ROOTSTOCK NAMED 'UFR-3'**
- (50) Latin Name: [(*Citrus reticulata*×*Citrus paradisi*)+*Citrus grandis*]×[*Citrus reticulata*+*Poncirus trifoliata* (trifoliate orange)]  
Varietal Denomination: UFR-3
- (71) Applicant: **Florida Foundation Seed Producers, Inc.**, Marianna, FL (US)
- (72) Inventor: **Jude W. Grosser**, Winter Haven, FL (US)
- (73) Assignee: **Florida Foundation Seed Producers, Inc.**, Marianna, FL (US)
- (\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/544,427**(22) Filed: **Jan. 5, 2015**(65) **Prior Publication Data**

US 2015/0195976 P1 Jul. 9, 2015

**Related U.S. Application Data**

- (60) Provisional application No. 61/923,586, filed on Jan. 3, 2014.
- (51) **Int. Cl.**  
**A01H 5/08** (2006.01)
- (52) **U.S. Cl.**  
USPC ..... **Plt./201**
- (58) **Field of Classification Search**  
USPC ..... Plt./201  
CPC ..... A01H 5/0806

See application file for complete search history.

- (56) **References Cited**
- U.S. PATENT DOCUMENTS
- PP21,535 P2 11/2010 Grosser
- OTHER PUBLICATIONS
- Florida Foundation Seed Producers, Inc.; 9 New Citrus Rootstocks—Descriptions and Data; May 2014, 14 pages.\*  
U.S. Appl. No. 14/544,424, filed Jan. 5, 2015, Grosser.  
U.S. Appl. No. 14/544,425, filed Jan. 5, 2015, Grosser.  
U.S. Appl. No. 14/544,423, filed Jan. 5, 2015, Grosser.  
U.S. Appl. No. 14/544,572, filed Jan. 22, 2015, Grosser.  
U.S. Appl. No. 14/544,571, filed Jan. 22, 2015, Grosser.  
U.S. Appl. No. 14/544,573, filed Jan. 22, 2015, Grosser.  
U.S. Appl. No. 14/545,063, filed Mar. 20, 2015, Grosser.  
U.S. Appl. No. 14/544,570, filed Jan. 22, 2015, Grosser.  
Grosser et al., "Protoplast fusion and citrus improvement," *Plant Breeding Reviews* 8:339-374, 1990.  
Grosser et al., "Development of "tetrazyg" rootstocks tolerant of the diaprepes/phytophthora complex under greenhouse conditions," *Proc. Fla. State Hort. Soc.* 116:263-267, 2003.  
Grosser et al., "Continued Development of Rootstocks Tolerant of the Phytophthora-Diaprepes Complex via Greenhouse Screening," *Proc. Fla. State Hort. Soc.* 120:103-109, 2007.  
Grosser et al., "Protoplast fusion for production of tetraploids and triploids: Applications for scion and rootstock breeding in citrus," *Plant Cell Tissue Organ Culture* 104:343-357, 2011.  
Letter and corresponding informational document to Florida citrus nurseries for the purpose of a possible license agreement, dated Jan. 9, 2014.

\* cited by examiner

**Primary Examiner** — Kent L Bell(74) **Attorney, Agent, or Firm** — Dentons US LLP(57) **ABSTRACT**

'UFR-3' is a new and distinct allotetraploid *citrus* rootstock for tree size control and improved disease resistance. 'UFR-3' has shown a positive reaction to the Huanglongbing disease (HLB, or *citrus* greening disease) in multiple experimental field trials. Scion trees grafted on this rootstock show a reduced frequency of infection and reduced disease symptoms once infected as compared to commercial diploid rootstocks.

**6 Drawing Sheets****1****BACKGROUND OF THE INVENTION**

Latin name of the genus and species of the plant claimed: allotetraploid hybrid of [(*Citrus reticulata*×*Citrus paradisi*)+*Citrus grandis*]×[*Citrus reticulata*+*Poncirus trifoliata* (trifoliate orange)].

Variety denomination: 'UFR-3'.

The present invention relates to a new and distinct variety of *citrus* rootstock named 'UFR-3'. The Plant Improvement Team in Lake Alfred, Fla. has pioneered the development and testing of allotetraploid *citrus* rootstocks. 'UFR-3' (identified as Orange #15 in field trials) is an allotetraploid zygotic hybrid derived from a conventional cross of two somatic hybrids previously produced by protoplast fusion. The somatic hybrid seed parent is 'Nova' mandarin hybrid+

**2**

'Hirado Buntan' pummelo (zygotic seedling) and the somatic hybrid pollen parent is 'Cleopatra' mandarin+'Argentine' trifoliate orange.

**5 BRIEF SUMMARY OF THE INVENTION**

'UFR-3' was selected on the basis of its positive reaction to Huanglongbing disease (HLB, or *citrus* greening disease) in multiple experimental field trials. The claimed plant was first asexually reproduced by grafting to 'Swingle' citrumelo and 'Volkameriana' rootstocks and planted in Lake Alfred, Fla. The resulting trees were true to type. Scion trees grafted onto this rootstock show a reduced frequency of infection and reduced disease symptoms once infected when compared to commercial diploid rootstocks. Although trees grafted to 'UFR-3' grow off quickly, trees remain small in

size, suitable for use in Advanced *Citrus* Production Systems that feature high density plantings. 'UFR-3' produces adequate nucellar seed, making it amenable to standard nursery propagation practices for uniform liner production. This selection exhibited a lower incidence of HLB infection than commercial rootstocks after 5 years in an exposed field trial. Long-term performance of trees on this rootstock selection is unknown. Yield and fruit quality data is limited, but 3 years of data from young trees (the St. Helena project, see Table 1) indicates good productivity and fruit quality. Tolerance to *citrus* blight is also unknown. Fruit quality for 'Valquarius'® 'SF14W-62' (U.S. Plant Pat. No. 21,535) scion was significantly less than that of 'Vernia'.

TABLE 1

		Rootstock Data from 5-year old trees in the St. Helena trial - Dundee, FL				
Scion	Rootstock	Lbs Solids/Box		Yield Boxes/Tree 2011		
		2012	2013	(35 mo.)		
'VAL-QUARIUS'	Som.	'UFR-6'	5.65	5.43	0.5	
'VERNIA'	Hyb.					
'VERNIA'	Som.	'UFR-6'	5.67	6.01	0.4	
'VAL-QUARIUS'	Hyb.					
'VAL-QUARIUS'	Tetrazyg	'UFR-1'	5.5	4.87	NS	
'VERNIA'	Tetrazyg	'UFR-1'	5.61	6.28	0.31	
'VERNIA'	Tetrazyg	'UFR-2'	5.47	5.93	0.35	
'VAL-QUARIUS'	Tetrazyg	'UFR-2'	4.57	5.37	NS	
'VAL-QUARIUS'	Tetrazyg	'UFR-3'	4.84	5.05	NS	
'VERNIA'	Tetrazyg	'UFR-3'	5.46	5.82	0.37	
'VERNIA'	Tetrazyg	'UFR-4'	5.79	6.07	0.54	
'VAL-QUARIUS'	Tetrazyg	'UFR-4'	4.65	5.07	NS	
'VAL-QUARIUS'	Tetrazyg	'UFR-5'	5.76	5.72	0.33	
'VERNIA'	Tetrazyg	'UFR-5'	5.89	5.34	0.42	
'VAL-QUARIUS'	Diploid	'FG 1731'	5.83	6.81	NS	
'VAL-QUARIUS'	Diploid	'FG 1731'	5.12	5.63	NS	
'VERNIA'	Diploid	'SWINGLE'*	5.11	5.79	0.33	
'VAL-QUARIUS'	Diploid	'SWINGLE'*	NS	5.61	NS	
'VERNIA'	Diploid	'CLEO'*	4.79	5.51	NS	
'VAL-QUARIUS'	Diploid	'CLEO'*	NS	5.21	NS	
'VERNIA'	Diploid	'R. LEMON'*	3.67	na	NS	
'VAL-QUARIUS'	Diploid	'VOLK'*	NS	4.12	NS	
'VERNIA'	Diploid	'VOLK'*	3.6	4.73	0.4	
'VAL-QUARIUS'	Diploid	'KU-HARSKE'*	NS	5.75	NS	
'VERNIA'	Diploid	'KU-HARSKE'*	4.34	5.83	0.15	

Scion	Rootstock	Yield Boxes/Tree		
		2012 (47 mo.)	2013 (59 mo.)	Cumulative Yield (Boxes)
'VAL-QUARIUS'	Som.	'UFR-6'	0.78	1.94
'VERNIA'	Hyb.			3.22
'VERNIA'	Som.	'UFR-6'	0.63	1.41
'VAL-QUARIUS'	Hyb.			2.44
'VAL-QUARIUS'	Tetrazyg	'UFR-1'	0.72	2.23
				2.95

TABLE 1-continued

Rootstock Data from 5-year old trees in the St. Helena trial - Dundee, FL						
5	'VERNIA'	Tetrazyg	'UFR-1'	0.67	1.33	2.31
	'VERNIA'	Tetrazyg	'UFR-2'	0.25	1.38	1.98
	'VAL-QUARIUS'	Tetrazyg	'UFR-2'	0.75	1.73	2.48
10	'VAL-QUARIUS'	Tetrazyg	'UFR-3'	0.81	1.97	2.78
	'VERNIA'	Tetrazyg	'UFR-3'	0.38	1.82	2.57
	'VERNIA'	Tetrazyg	'UFR-4'	0.71	1.73	2.98
	'VAL-QUARIUS'	Tetrazyg	'UFR-4'	0.65	1.59	2.64
15	'VAL-QUARIUS'	Tetrazyg	'UFR-5'	0.56	1.80	2.69
	'VERNIA'	Tetrazyg	'UFR-5'	0.25	1.93	2.60
	'VAL-QUARIUS'	Diploid	'FG 1731'	0.68	2.20	2.88
20	'VAL-QUARIUS'	Diploid	'FG 1731'	0.67	2.77	3.44
	'VERNIA'	Diploid	'SWINGLE'*	0.85	1.08	2.26
	'VAL-QUARIUS'	Diploid	'SWINGLE'*	NS	1.50	1.50
25	'VAL-QUARIUS'	Diploid	'VOLK'*	NS	2.58	2.58
	'VERNIA'	Diploid	'VOLK'*	1.13	0.83	2.36
	'VAL-QUARIUS'	Diploid	'KU-HARSKE'*	NS	2.2	2.2
30	'VERNIA'	Diploid	'KU-HARSKE'*	0.75	1.08	1.98

NS—not significant fruit; na—data not available;

\*control commercial rootstock

One box contains approximately 90 lbs. fruit.

## BRIEF DESCRIPTION OF THE DRAWINGS

'UFR-3' is illustrated by the accompanying photographs, which show the tree's form, foliage, and fruit. The colors shown are as true as can be reasonably obtained by conventional photographic procedures. The photographs are of a tree approximately 6 years old. All figures were taken of the same tree during the fall.

FIG. 1.—Shows a close-up of the nearly mature fruits with the rind and cross-sectional view of the fruit when cut in the center.

FIG. 2.—Shows the overall mature plant growth habit in the fall of 2013.

FIG. 3.—Shows nearly mature fruits hanging on the tree.

FIG. 4.—Shows a close-up of leaves and nearly mature fruits.

FIG. 5.—Shows a close-up of nearly mature fruits.

FIG. 6.—Shows a close-up of seeds from nearly mature fruit.

## DETAILED BOTANICAL DESCRIPTION

The following detailed description sets forth the distinctive characteristics of 'UFR-3'. The colors (except those in common terms) are described from R.H.S. Colour Chart published by The Royal Horticultural Society in London (second edition), in association with the Flower Council of Holland.

Phenotypic Description of *Citrus reticulata*  
‘UFR-3’

## Classification:

*Botanical*.—/[*Citrus reticulata* hybrid ('Clementine' mandarin×'Orlando' tangelo) *Citrus reticulata*×*Citrus paradisi*]+*Citrus grandis*]×[*Citrus reticulata*+*Poncirus trifoliata* (trifoliate orange)].

*Common name*.—Complex allotetraploid 'tetrazyg' rootstock hybrid.

## Parentage:

*Female parent*.—'Nova' mandarin+‘Hirado Buntan’ pummelo (zygotic seedling) somatic hybrid (unpatented).

*Male parent*.—‘Cleopatra’ mandarin+‘Argentine’ trifoliate orange; somatic hybrid (unpatented).

## Tree:

*Ploidy*.—Tetraploid.

*Size*.—Medium.

*Height*.—3.70 meters.

*Tree spread*.—3.8 to 4.3 meters.

*Vigor*.—Vigorous.

*Density*.—Canopies are quite dense.

*Form*.—The tree is obloid-shaped with lateral and upright branches growing toward low to medium angles. Branches with fruit exhibit drooping.

*Growth habit*.—Both upright and lateral growth with low medium angle.

## Trunk:

*Trunk diameter*.—20 cm in diameter at 30 cm above the ground on a 6-year-old tree.

*Trunk texture*.—Smooth.

*Trunk bark color*.—RHS 195A (greyed-green); irregularly striated with RHS N189A (greyed-green).

## Branches:

*Crotch angle*.—First crotch forms a 55- to 60-degree angle, middle crotch forms a 50-degree angle.

*Branch length*.—Branch reaches 4.2 meters from the first crotch to the tip of the branch.

*Branch texture*.—Relatively rough with small thorns or spines.

*Branch color (shoots from previous flush, hardened and 4 to 5 mm in diameter)*.—RHS 137A (green).

## Leaves:

*Size (lamina average)*.—Length: 91.5 mm. Width: 62.5 mm. L/W ratio: 1.46.

*Thickness*.—Thicker than regular and average for diploid commercial rootstock hybrids.

*Type*.—Simple leaves with a small amount of trifoliate and biofoliate leaves.

*Shape*.—Elliptical.

*Apex*.—Retuse.

*Base*.—Acute to sub-obtuse.

*Margin*.—Entire and slightly undulate.

*Surface*.—Upper surface: Glabrous. Lower surface: Medium veins that are pinnately netted.

*Color*.—Upper surface (adaxial): RHS 137A (green). Lower surface (abaxial): RHS 137C (green).

*Petiole*.—Shape: Brevipetiolate (shorter than leaf lamina); junction between petiole and lamina is articulate. Width (petiole wing): Narrow. Shape (petiole wing): Obovate. Length: 17 to 19.5 mm. Width: 5.1 to 6.5 mm. Color: RHS 137A (green).

## Flowers and flower buds:

*Type*.—Hermaphrodite.

*Bearing*.—Flower grown from leaf axillaries and leaf terminals singly and in small clusters; most single flowers grow from leaf axillaries.

*Flower bud size*.—Shape: Initial visible flower bud has a round ball shape; mature flower bud has an elongated olive shape.

*Flower petals*.—Shape: Flat, spatula-shaped. Apex shape: Smooth, acute-shaped. Base shape: Even obtuse. Margin: Smooth.

*Flower sepal*.—Shape: Delta-shaped with an acute angle at the apex. Apex shape: Triangle-shaped. Margin: Smooth.

*Fragrance*.—Fragrant/Moderately fragrant.

*Reproductive organs*.—Fertility: Appears self-fertile. Pollen amount: Abundant/Moderately amount. Pollen color (general): Bright-yellow. Ovary shape: Oval-shaped.

## Fruit:

*Size*.—Uniform.

*Height*.—70.5 to 72.8 mm on average.

*Width*.—86 to 94 mm on average.

*Average weight (per individual fruit)*.—302.1 grams.

*Shape*.—Round.

*Shape (cross-section)*.—Round.

*Apex*.—Truncated with slight dent.

*Apex cavity diameter*.—N/A.

*Base cavity diameter*.—6.9 to 8.2 mm.

*Base*.—No neck.

*Harvesting*.—Fruit can be harvested from October through December in Florida.

*Fruit stem (short stem connecting the fruit)*.—Length: 8.0 mm. Diameter: 5.8 mm. Color: RHS N137A (green) with RHS 195B (greyed-green) strip.

## Rind:

*Adherence*.—Adherence between albedo (mesocarp) and flesh (endocarp) is medium. The adherence is evenly distributed from based to apex.

*Thickness*.—6.0 to 7.8 mm on average.

*Texture*.—Smooth.

*Color*.—Flavedo (epicarp): Ranges between RHS 150C (yellow-green) to RHS 151D (yellow-green). Albedo (mesocarp): RHS 150C (white).

*Stylar end*.—Closed.

*Rind oil cell density*.—145 oil cells/square cm.

## Flesh:

*Number of segments*.—Between 9 and 10 segments per fruit on average.

*Segment walls*.—Medium-firm with sufficient strength to maintain integrity as separated.

*Juice*.—Abundant.

*Color*.—Uniformly RHS 14D (yellow-orange).

*Texture*.—Medium soft.

*Vesicles*.—Length: Arranged from 13.5 to 21.5 mm on average. Diameter (thickness): 3.2 to 3.6 mm on average.

*Eating quality*.—N/A.

*Juice index*.—Soluble solids (average): 8.5 Brix.

## Seeds:

*Type*.—Polyembryonic and monoembryonic.

*Number*.—Ranges from 5 to 23. Most fruit has less than 10 seeds.

*Shape.*—Seed shapes are not uniform. Normal seeds are mostly ventricose/swollen-shaped and clavate club shaped. Many seeds appear to be wrinkled and flatter than normal.

*Size.*—Length: 13.6 to 18.6 mm. Width: 5.5 to 7.2 mm.

*Seed coat color.*—Outer Surface: RHS 155C (white) and wrinkled. Inner surface: RHS 164B (greyed-orange). Cotyledon color: RHS 144D (green).

Resistance to disease: ‘UFR-3’ rootstock was selected on the basis of its positive reaction to HLB disease (huanglong-


5

bing or *citrus* greening disease) in multiple field trials. Trees on this rootstock show a reduced frequency of infection, and reduced disease symptoms once infected as compared to commercial diploid rootstocks.

What is claimed is:

1. A new and distinct *citrus* rootstock plant as illustrated and described herein.

\* \* \* \* \*



**FIG. 1**




**FIG. 2**



**FIG. 3**



**FIG. 4**



**FIG. 5**



**FIG. 6**