



US00PP27928P2

(12) **United States Plant Patent**
Zaiger et al.

(10) **Patent No.:** **US PP27,928 P2**
(45) **Date of Patent:** **Apr. 25, 2017**

(54) **PEACH TREE NAMED ‘ZEE RICH’**

(50) Latin Name: *Prunus persica*
Varietal Denomination: **Zee Rich**

(71) Applicants: **Gary Neil Zaiger**, Modesto, CA (US);
Leith Marie Gardner, Modesto, CA
(US); **Grant Gene Zaiger**, Modesto,
CA (US)

(72) Inventors: **Gary Neil Zaiger**, Modesto, CA (US);
Leith Marie Gardner, Modesto, CA
(US); **Grant Gene Zaiger**, Modesto,
CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/998,650**

(22) Filed: **Jan. 29, 2016**

(51) **Int. Cl.**
A01H 5/08 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./197**

(58) **Field of Classification Search**
USPC **Plt./197**
See application file for complete search history.

Primary Examiner — Annette Para

(57) **ABSTRACT**

A new and distinct variety of peach tree (*Prunus persica*).
The following features of the tree and its fruit are charac-
terized with the tree budded on ‘Nemaguard’ Rootstock
(non-patented), grown on Handford sandy loam soil with
Storie Index rating 95, in USDA Hardiness Zone 9, near
Modesto, Calif., with standard commercial fruit growing
practices, such as pruning, thinning, spraying, irrigation and
fertilization. Its novelty consist of the following combina-
tion of desirable features:

1. Tree having a vigorous, upright growth habit.
2. Tree being a regular and productive bearer of large size
fruit.
3. Fruit with an attractive, red skin color.
4. Fruit with very good flavor and eating quality.
5. Clingstone fruit with firm, yellow flesh.

1 Drawing Sheet

1

Botanical designation: *Prunus persica*.
Variety denomination: ‘Zee Rich’.

BACKGROUND OF THE VARIETY

Field of the Invention

In the field of plant genetics, we conduct an extensive and
continuing plant-breeding program including the organiza-
tion and asexual reproduction of orchard trees, and of which
plums, peaches, nectarines, apricots, cherries, almonds and
interspecifics are exemplary. It was against this background
of our activities that the present variety of peach tree was
originated and asexually reproduced by us in our experi-
mental orchard located near Modesto, Stanislaus County,
Calif.

PRIOR VARIETIES

Among the existing varieties of peach trees, which are
known to us, and mentioned herein, ‘Vista’ Peach (U.S.
Plant Pat. No. 9,549), ‘Sweet Henry’ Peach (U.S. Plant Pat.
No. 16,068), ‘Sierra Rich’ Peach (U.S. Plant Pat. No.
12,391) and our proprietary non-patented peach seedling
selection ‘11ZL1010’.

**STATEMENT REGARDING FEDERALLY
SPONSORED RESEARCH AND
DEVELOPMENT**

Not applicable.

ORIGIN OF THE VARIETY

The new and distinct variety of peach tree (*Prunus
persica*) was developed by us in our experimental orchard

2

located near Modesto, Calif. from open pollinated seed
collected from our proprietary non-patented peach seedling
selection with the field identification number ‘11ZL1010’.
The seed parent (11ZL1010) originated as a first generation
cross between ‘Vista’ Peach (U.S. Plant Pat. No. 9,549) and
‘Sweet Henry’ Peach (U.S. Plant Pat. No. 16,068). A large
number of these open pollinated seedlings were planted and
grown on their own root system, during which time we
recognized the desirable fruit and tree characteristics of the
present seedling and selected it in 2011 for additional
asexual propagation and commercialization.

ASEXUAL REPRODUCTION OF THE VARIETY

In 2011 asexual reproduction of the new and distinct
variety of peach tree was by budding to ‘Nemaguard’
Rootstock (non-patented), as performed by us in our experi-
mental orchard located near Modesto, Calif., and shows that
reproductions run true to the original tree and all character-
istics of the tree and its fruit are established and transmitted
through succeeding asexual propagations.

SUMMARY OF THE NEW VARIETY

The present new variety of peach tree (*Prunus persica*) is
of large size, vigorous, upright growth and a regular and
productive bearer of large size, yellow flesh, clingstone fruit
with very good flavor and eating quality. The fruit is further
characterized by having firm flesh and an attractive red skin
color. In comparison to its proprietary non-patented peach
seed parent ‘11ZL1010’ the fruit of the new variety is three
days later in maturity. In comparison to the commercial
variety ‘Sierra Rich’ Peach (U.S. Plant Pat. No. 12,391) the

fruit of the new variety is 10 days later in maturity and has a more attractive red skin color.

DESCRIPTION OF THE PHOTOGRAPH

The accompanying color photographic illustration shows typical specimens of the foliage and fruit of the present new peach variety.

The illustration shows the upper and lower surface of the leaves, an exterior and sectional view of a single fruit divided in its suture plane to show flesh color, pit cavity and the stone remaining in place.

The photographic illustration was taken shortly after being picked (shipping ripe) from a 4 year old tree and the colors are as nearly true as is reasonably possible in a color representation of this type.

DESCRIPTION OF THE VARIETY

The following is a detailed botanical description of the new variety of peach tree, its flowers, foliage and fruit, as based on observations of 4 year old specimens grown near Modesto, Calif., with color in accordance with Munsell Book of Color published in 1958.

Tree:

Size.—Large, usually pruned to 3 to 3.5 meters in height and width for economical harvesting of fruit. Size varies with different cultural practices.

Vigor.—Vigorous, growth of 1.5 to 2 meters in height the first growing season. Varies slightly with type and fertility of soil, climatic conditions and cultural practices.

Form.—Upright, usually pruned to vase shape.

Branching habit.—Upright, crotch angle approximately 35°, increases with heavy crop load.

Productivity.—Productive, thinning and spacing of fruit necessary for desired market size fruit. Number of fruit set varies with climatic conditions during blooming period.

Bearer.—Regular, has had adequate fruit set 3 consecutive years. No alternate bearing observed.

Fertility.—Self fertile.

Density.—Medium dense, usually pruned to vase shape to increase air movement and sunlight to enhance fruit color and health of fruit wood.

Hardiness.—Hardy in all stone fruit growing areas of California. Tree grown in USDA Hardiness Zone 9. Winter chilling requirement approximately 700 hours at or below 45° F.

Trunk:

Size.—Large, average circumference 53.4 cm at 25.4 cm above ground on a 4 year old tree.

Stocky.—Medium stocky.

Texture.—Medium shaggy, roughness increases with age.

Color.—Varies from 2.5GY 4/2 to 7.5YR 3/2.

Branches:

Size.—Medium. Average circumference 9.7 cm at 1.2 meters above ground. Crotch angle approximately 35°, increases with heavy crop load.

Surface texture.—New growth relatively smooth. Mature growth medium rough, roughness increases with age.

Lenticels.—Average number 36 in a 25.8 square cm area. Average length 4.0 mm. Average width 2.0 mm. Color varies from 7.5YR 6/12 to 10YR 6/12.

Color.—New growth varies from 5GY 6/6 to 2.5GY 5/8. Mature growth varies from 5YR 3/4 to 2.5YR 2/4, varies with age of growth.

Leaves:

Size.—Medium to large. Average length 110.1 mm. Average width 32.6 mm.

Form.—Lanceolate.

Apex.—Acuminate.

Base.—Cuneate.

Margin.—Crenate.

Thickness.—Medium.

Surface texture.—Upper surface relatively smooth, slight indentations over midrib and leaf veins. Lower surface relatively smooth, except for small ridges created by midrib and pinnate venation. Both upper and lower surfaces glabrous.

Petiole.—Average length 8.6 mm. Average width 1.5 mm. Longitudinally grooved. Surface — glabrous. Color varies from 2.5GY 6/6 to 10Y 6/6.

Glands.—Type — reniform. Size — medium. Average length 1.0 mm. Average diameter 0.5 mm. Number varies from 2 to 3, average number 2. Located primarily on the base of the leaf blade and upper portion of the petiole. Color varies from 10R 2/6 to 10Y 5/6.

Stipules.—Average number 2. Average length 6.3 mm. Edges — pectinate. Color varies from 2.5GY 6/8 to 5GY 6/8.

Color.—Upper surface varies from 2.5GY 3/4 to 10Y 3/4. Lower surface varies from 2.5GY 5/4 to 5GY 4/4. Midvein color varies from 7.5Y 8/6 to 10Y 7/8.

Flower buds:

Size.—Large. Average length 18.5 mm. Average diameter 10.0 mm.

Hardiness.—Hardy with respect to California winters.

Density.—Medium dense.

Form.—Conical, becoming elongated just before opening.

Pedicel.—Average length 4.3 mm. Average width 1.4 mm. Surface — glabrous. Color varies from 7.5GY 8/6 to 7.5GY 7/8.

Color.—Varies from 7.5RP 9/2 to 10RP 8/6, color fades with age.

Flowers:

Blooming period.—Date of First Bloom Feb. 11, 2015. Date of Petal Fall Feb. 21, 2015, varies slightly with climatic conditions.

Size.—Large, showy. Average height 25.6 mm. Average diameter 45.9 mm.

Petals.—Normally 5, alternately arranged to sepals. Size — large. Average length 23.2 mm. Average width 21.9 mm. Form — globose to obovate. Petal apex — rounded. Petal base — truncate. Margin — sinuate. Arrangement — slightly overlapping to free. Both upper and lower surfaces glabrous. Color varies from 5RP 9/2 to 5RP 5/12.

Sepals.—Normally 5, alternately arranged to petals. Size — large. Average length 6.5 mm. Average width 6.0 mm. Shape — ovate, apex rounded to triangular. Margin — entire. Color — upper surface varies from 5GY 5/8 to 5GY 4/8. Lower surface varies from 5RP 3/10 to 5RP 2/8. Surface — upper surface glabrous, lower surface pubescent.

Stamens.—Average number per flower 52. On average, the stamens are below the height of the petals.

Filament color N 9.5/(white) to 2.5RP 7/8, depending on age of flower. Anther color varies from 7.5R 4/2 to 5Y 8.5/8.

Pollen.—Self fertile. Color varies from 2.5Y 7/12 to 5Y 7/12.

Pistil.—Normally one. Average length 18.9 mm. Position of stigma an average of 1.4 mm below anthers. Surface — pubescent. Color varies from 10Y 8/6 to 2.5GY 7/6.

Fragrance.—Slight.

Color.—Varies from 5RP 9/2 to 5RP 5/12.

Pedice.—Average length 6.4 mm. Average width 1.4 mm. Surface — glabrous. Color varies from 5GY 6/8 to 5GY 5/10.

Number flowers per flower bud.—Normally one.

Fruit:

Maturity when described.—Firm ripe and ready for consumption.

Date of first picking.—Jul. 4, 2015.

Date of last picking.—Jul. 14, 2015, varies slightly with climatic conditions.

Size.—Large. Average diameter axially 69.1 mm. Average transversely in suture plane 81.8 mm. Average weight 261.3 grams, varies slightly with fertility of the soil, amount of thinning and climatic conditions.

Form.—Globose.

Suture.—Slightly lipped, extends from base to apex.

Ventral surface.—Slightly lipped.

Apex.—Rounded to slight tip.

Base.—Retuse.

Stem cavity.—Rounded to slightly elongated in suture plane. Average depth 11.1 mm. Average diameter 10.4 mm.

Stem:

Size.—Small to medium. Average length 8.4 mm. Average diameter 3.1 mm.

Color.—Varies from 10Y 6/8 to 2.5GY 5/6.

Flesh:

Ripens.—Evenly.

Texture.—Firm, meaty.

Fibers.—Few, small, tender.

Firmness.—Firm, comparable to other commercial peach varieties.

Aroma.—Moderate.

Amygdalin.—Undetected.

Eating quality.—Very good, good balance between acid and sugar.

Flavor.—Very good, slightly tart.

Juice.—Moderate amount, enhances flavor.

Acidity.—Not available.

Brix.—Average Brix 10.0°, varies slightly with amount of fruit per tree and climatic conditions.

Color.—Varies from 2.5Y 8.5/10 to 2.5Y 8/10.

Pit cavity.—Average length 36.3 mm. Average width 28.8 mm. Average depth 11.0 mm. Color varies from 2.5Y 8/8 to 7.5R 3/12.

Skin:

Thickness.—Medium.

Surface.—Smooth.

Pubescence.—Moderate amount, short in length.

Tendency to crack.—None.

Color.—Ground color varies from 5Y 8/8 to 5Y 7/8. Overspread with 7.5R 4/10 to 7.5R 2/4.

Tenacity.—Tenacious to the flesh.

Astringency.—Slight to none.

Stone:

Type.—Clingstone, strong adherence to flesh.

Size.—Large. Average length 35.3 mm. Average width 27.8 mm. Average thickness 20.0 mm.

Form.—Obovoid.

Base.—Flat.

Apex.—Pointed. Average length 1.7 mm.

Surface.—Pitted throughout, pits vary from rounded to elongated.

Sides.—Unequal, one side extending further from suture plane.

Ridges.—Relatively smooth, extending from base to apex.

Tendency to split.—None.

Color.—Varies from 7.5YR 4/8 to 7.5R 3/10 when dry.

Kernel:

Size.—Medium. Average length 18.5 mm. Average width 11.5 mm. Average depth 5.7 mm.

Form.—Ovoid.

Viability.—Viable, complete embryo development.

Skin color.—Varies from 5Y 9/6 to 5Y 8.5/6.

Use: Dessert.

Market.—Local and long distance.

Keeping quality: Good, held firm in cold storage 3 weeks at 38° to 42° F. without shriveling, internal breakdown of flesh or appreciable loss of flavor.

Shipping quality: Good, showed minimal skin scarring or flesh bruising during picking, packing and shipping trials.

Plant/fruit disease resistance/susceptibility: No specific testing for relative plant/fruit disease resistance/susceptibility has been designed. Under close observation during planting, growing, and harvesting of fruit, under normal cultural and growing conditions near Modesto, Calif., no particular plant/fruit disease resistance or susceptibility has been observed. Any variety or selection observed during indexing of plant characteristics with abnormal fungus, bacterial, virus or insect susceptibility is destroyed and eliminated from our breeding program. No atypical resistances/susceptibilities have been noted under normal cultural practices.

The present new variety of peach tree, its flowers, foliage and fruit herein described may vary in slight detail due to climate, soil conditions and cultural practices under which the variety may be grown. The present description is that of the variety grown under the ecological conditions prevailing near Modesto, Calif.

The invention claimed is:

1. A new and distinct variety of peach tree, substantially as illustrated and described.

* * * * *

