

US00PP28258P3

(12) United States Plant Patent Hanes

(10) Patent No.: US PP28,258 P3

(45) **Date of Patent:** Aug. 8, 2017

(54) PELARGONIUM PLANT NAMED 'PEQZ0010'

(50) Latin Name: *Pelargonium interspecific* Varietal Denomination: **PEQZ0010**

(71) Applicant: SYNGENTA PARTICIPATIONS AG,

Basel (CH)

(72) Inventor: Mitchell E. Hanes, Gilroy, CA (US)

(73) Assignee: Syngenta Participations AG, Basel

(CH)

(*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/999,661**

(22) Filed: Jun. 9, 2016

(65) Prior Publication Data

US 2016/0366800 P1 Dec. 15, 2016

(51) Int. Cl. A01H 5/02

(2006.01)

(56) References Cited

PUBLICATIONS

PLUTO Plant Variety Database Apr. 26, 2017.*

* cited by examiner

Primary Examiner — Annette Para

(74) Attorney, Agent, or Firm — Dale Skalla

(57) ABSTRACT

A new *Pelargonium* plant named 'PEQZ0010' particularly distinguished dark lavender single petal inflorescences held above the medium-green foliage, very heat and drought tolerant with continuous color through the most extreme summer heat, and exceptional edema tolerance on a well-branched plant habit.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed: *Pelargonium interspecific*.

Varietal denomination: 'PEQZ0010'.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new *Pelargonium*, botanically known as *Pelargonium interspecific*, and hereinafter referred to by the variety name 'PEQZ0010'.

'PEQZ0010' is a product of a planned breeding program. The new cultivar 'PEQZ0010' has intense dark lavender colored inflorescences held above the green foliage, very heat and drought tolerant with continuous color through the medium extreme summer heat, and exceptional edema tolerance on a vigorous plant.

'PEQZ0010' originates from a hybridization in a controlled breeding program made in February 2007, in a greenhouse in Gilroy, Calif. The female parent was an unpatented, proprietary plant of *P. interspecific* parentage, identified as '10518-4' a hot pink single flowered variety with more open plant habit when compared to 'PEQZ0010'.

The male parent of 'PEQZ0010' was an unpatented, proprietary plant of *P. interspecific* parentage, identified as '10622-1' with more blue-rose colored florets and a more restricted plant habit when compared to 'PEQZ0010'. The resultant seed was sown in August 2007.

'PEQZ0010' was selected as one flowering plant within the progeny of the stated cross in December 2007 in a greenhouse in Gilroy, Calif.

The first act of asexual reproduction of 'PEQZ0010' was accomplished when vegetative cuttings were propagated from the initial selection in the December 2007 in a greenhouse in Gilroy, Calif.

2

BRIEF SUMMARY OF INVENTION

Horticultural examination of plants grown from cuttings of the plant initiated in Jun. 2, 2016 in Gilroy, Calif., and continuing thereafter, has demonstrated that the combination of characteristics as herein disclosed for 'PEQZ0010' are firmly fixed and are retained through successive generations of asexual reproduction.

'PEQZ0010' has not been observed under all possible environmental conditions. The phenotype may vary significantly with variations in environment such as temperature, light intensity, and day length.

A Plant Breeder's Right for this cultivar was applied for in the European Union on Jun. 10, 2015, No. 2015/1216. 'PEQZ0010' has not been made publicly available more than one year prior to the filing of this application.

The following traits have been repeatedly observed and are determined to be the basic characteristics of the new variety. The combination of these characteristics distinguishes this *Pelargonium* as a new and distinct variety.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographic drawing shows typical flower and foliage characteristics of 'PEQZ0010' with colors being as true as possible with an illustration of this type. The photographic drawing shows in FIG. 1, 3 flowering plants of the new variety and in FIG. 2, a close-up of an inflorescence.

DETAILED BOTANICAL DESCRIPTION

The plant descriptions and measurements were taken in Gilroy, Calif. in Jun. 7, 2016 under natural light. These

3

plants were approximately 7 weeks old and were grown in Inflorescence: *Type.*—Umbel; semi-spherical or nearly semi-spheria 4 cell packs in a greenhouse trial. cal. Color references are made to The Royal Horticultural *Lastingness of individual flowers.*—7-9 days at 18° C. Society Colour Chart (R.H.S.) 2001. temperature. Plant: Number of inflorescences per plant.—12, with 15 Form, growth and habit.—Upright, outwardly spreadimmature umbels in various stages. ing and rounded growth habit, heat and drought Fragrance.—None. tolerant with continuous color through the most Umbel diameter.—11-12 cm. extreme summer heat, edema tolerance, well-*Umbel depth.*—5.5-6 cm. branched plant habit. Corolla: Plant height.—9-10.5 cm. Form.—Single. Plant height (inflorescence included).—12-13 cm. *Number of petals.*—4-5. Plant width.—16-17 cm. Diameter of flower.—4.5-4.6 cm. Roots: Depth of flower.—2-2.2 cm. *Number of days to initiate roots.*—15-18 days at about 15 Color upper petals, upper surface.—RHS N74A with 22 degrees C. RHS 75B veining. Number of days to produce a rooted cutting.—21-23 Color upper petals, lower surface.—RHS N74B. days at 22 degrees C. Length of upper petals.—2.2-2.5 cm. *Type.*—Fine, fibrous, free branching. Width of upper petals.—1.7-1.8 cm. Color.—RHS 156B. Color lower petals, upper surface.—RHS 71D. Foliage: Color lower petals, lower surface.—RHS 72D. Immature leaf, color upper surface.—RHS 137C. Length of lower petals.—2.6 cm. Immature leaf, lower surface.—RHS 138B. Width of lower petals.—1.4 cm. Mature leaf, color upper surface.—RHS 137A. *Petal shape.*—Obovate to spathulate. Mature leaf, color lower surface.—RHS 137C. *Apex shape.*—Rounded. *Length.*—4.0-4.3 cm. *Margin*.—Entire. *Width.*—7.2-7.5 cm. Base.—Attenuate. Shape.—Cordate. Petal texture.—Papillose on both surfaces. Base shape.—Cordate. 30 Calyx: Apex shape.—Reniform. *Number of sepals.*—4-5. *Margin.*—Slightly dentate. Color of sepals.—RHS N144C. Texture upper side.—Hirsute. Length of sepals.—1.1 cm. Texture lower side.—Hirsute. Width of sepals.—0.4-0.5 cm. Leaf zonation color.—None. Sepal shape.—Lanceolate to linear. Color of veins, upper surface.—RHS 137B becoming 35 *Apex shape.*—Acute. indistinct. Margins.—Mostly fused. Color of veins, lower surface.—RHS 137C becoming Texture, upper surface.—Glabrous. indistinct. Lower surface.—Glandular hairs, hirsute. Pattern of veining.—Palmate. 40 Reproductive organs: Petiole color.—RHS 137C. Gynoecium: Petiole length.—6.8-8.5 cm. Pistil.—1. Diameter of petiole.—0.2 cm. Length.—0.9 cm. *Texture*.—Pilose, hirsute, glandular hairs. Style color.—RHS 50A. Stem: Style length.—0.4-0.5 cm. Quantity of branches.—4-5. Stigma color.—RHS 53D. Color of stem.—RHS 137C. Ovary color.—RHS 149A. Length of stem.—6.5-7.0 cm. Ovary length.—0.5-0.6 cm. Diameter.—0.6 cm. Ovary diameter.—0.2-0.3 cm. Length of internodes.—1.2 cm. 50 Androecium: *Texture.*—Sparsely hirsute, pilose, glandular hairs. *Number of stamens.*—7. Peduncle: Color of filaments.—RHS N57C. Color of peduncle.—RHS 137A. Length filaments.—0.6 cm. Length of peduncle.—10.8-11 cm. Anther color.—RHS 71D. Peduncle diameter.—0.3 cm. Length of anthers.—0.3 cm. *Texture*.—Hirsute, glandular hairs. Color of pollen.—RHS 25A. Pedicel: Pollen amount.—Normal. Color of pedicel.—RHS 187A. Fertility/seed set.—Has not been determined to date. Length of pedicel.—3.5-3.9 cm. Disease/pest resistance.—Has not been determined to Diameter of pedicel.—0.2 cm. date. 60 Texture.—Sparsely pilose, glandular hiars. What is claimed is: Bud (just before opening): 1. A new and distinct variety of *Pelargonium* plant named Color.—RHS N74A with more blue.

Length.—1.5 cm.

Width.-0.5-0.6 cm.

Shape.—Elliptical.

65 * * * *

herein.

'PEQZ0010' substantially as illustrated and described



FIGURE 1



FIGURE 2