

US00PP28369P3

(12) **United States Plant Patent**
Finn

(10) **Patent No.:** **US PP28,369 P3**
(45) **Date of Patent:** **Sep. 12, 2017**

(54) **BLACKBERRY PLANT NAMED ‘COLUMBIA GIANT’**

(50) Latin Name: *Rubus* subg. *Rubus* Watson
Varietal Denomination: **Columbia Giant**

(71) Applicant: **The United States of America, as represented by the Secretary of Agriculture, Washington, DC (US)**

(72) Inventor: **Chad E. Finn, Corvallis, OR (US)**

(73) Assignee: **THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY OF AGRICULTURE, Washington, DC (US)**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 24 days.

(21) Appl. No.: **14/756,637**

(22) Filed: **Sep. 28, 2015**

(65) **Prior Publication Data**
US 2017/0094856 P1 Mar. 30, 2017

(51) **Int. Cl.**
A01H 5/08 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./203**

(58) **Field of Classification Search**
USPC Plt./230, 203
See application file for complete search history.

Primary Examiner — June Hwu

(74) *Attorney, Agent, or Firm* — Gail Poulos; John D. Fado

(57) **ABSTRACT**

A new and distinct blackberry cultivar that originated from seed produced from a cross between the female blackberry plant ‘NZ 9629-1’ (unpatented) and the male parent blackberry plant ‘ORUS 1350-2’ (unpatented). This new blackberry cultivar can be distinguished by its high yields of extremely large and very uniformly shaped berries with very good firmness, color, and flavor and that are borne on vigorous, completely thornless trailing plants.

4 Drawing Sheets

1

Latin name of the genus and species of the plant claimed: ‘COLUMBIA GIANT’ is a blackberry plant that is *Rubus* subg. *Rubus* Watson.

Variety denomination: The new blackberry plant claimed is of the variety denominated ‘Columbia Giant’ *Rubus* subg. *Rubus* Watson.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct blackberry cultivar designated ‘Columbia Giant’ and botanically known as *Rubus* subg. *Rubus* Watson. This new blackberry cultivar was discovered in Corvallis, Oreg. in July 2008 and originated from a cross between the female blackberry plant ‘NZ 9629-1’ (unpatented) and the male parent blackberry plant ‘ORUS 1350-2’ (unpatented). ‘Columbia Giant’'s spinelessness is derived from ‘Lincoln Logan’ (unpatented) that can be found as a parent four and five generations back in ‘Columbia Giant’'s pedigree. ‘Columbia Giant’ was selected from the same population as ‘Columbia Star’ (U.S. Plant patent application Ser. No. 13/815,074, now U.S. Plant Pat. No. 25,532). The original seedling of the new cultivar was asexually propagated at a nursery in Benton County, Oreg. The new cultivar was established in vitro from a cane cutting and microcuttings have been taken and rooted from this sort of culture. The present invention has been found to be stable and to reproduce true to type through successive asexual propagations.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

This new blackberry plant is illustrated by the accompanying photographs that show the fruit of the plant and

2

machine harvested fruit, as well as canes and entire plants; the colors shown are as true as can be reasonably obtained by conventional photographic procedures.

FIG. 1. shows a section of a spineless primocane and a leaf.

FIG. 2. shows flowers on the plant. As is typical for commercial production, trailing primocanes are lifted from the ground in late summer and tied to a two wire trellis with the lower wire approximately 1.0 m above the ground and the upper wire approximately 1.5 m above the ground.

FIG. 3. shows typical fruit in a fruit cluster in the field.

FIG. 4. shows an entire 4-year old plant. As is typical for commercial production, trailing primocanes are lifted from the ground in late summer and tied to a two wire trellis with the lower wire approximately 1.0 m above the ground and the upper wire approximately 1.5 m above the ground.

DETAILED DESCRIPTION OF THE NEW VARIETY

The following description of ‘Columbia Giant’ is based on observations taken from 2011 to 2014 growing seasons in trials in Corvallis and Aurora, Oreg. This description is in accordance with UPOV terminology. Color designations, color descriptions and other phenotypical descriptions may deviate from the stated values and descriptions depending upon variation in environmental, seasonal, climatic and cultural conditions. ‘Columbia Giant’ has not been observed under all possible environmental conditions. Color terminology follows The Royal Horticultural Society Colour Chart. London (R.H.S.) (5th edition, 2007).

Table 1 shows plant characteristics of the new cultivar. Characteristics include plant vigor growth habit, date of

budbreak, timing of primocane emergence, winter tolerance in Aurora, Oreg. (45° 16' 49" N/122° 44' 50" W), and winter tolerance in Lynden, Wash. (48° 56' 48" N/122° 27' 2" W).

TABLE 1

Plant Characteristics of 'Columbia Giant'.	
Characteristic	Columbia Giant
Plant vigor	High compared to Black Diamond
Growth habit	Trailing
Date of budbreak	27 March
Timing of primocane emergence	Early to medium compared to Black Diamond
Winter tolerance in Aurora, Oregon (45° 16' 49" N/122° 44' 50" W)	Very good-excellent (better than Marion)
Winter tolerance in Lynden, Washington (48° 56' 48" N/122° 27' 2" W)	Good-very good (better than Marion)

Table 2 shows floricanes and mature primocane characteristics of the new cultivar. Characteristics include diameter at base, diameter at midpoint, diameter at terminus, internode length at base, internode length at midpoint, internode length at terminus, presence of spines further than 0.6 m from the soil surface, presence of spines less than 0.6 m from the soil surface, floricanes color at base, floricanes color at midpoint, floricanes color at terminus, floricanes lateral length, floricanes lateral strength, primocane color at base, primocane color at midpoint, primocane color at terminus, and floricanes length (range).

TABLE 2

Floricanes and Mature Primocane Characteristics of 'Columbia Giant'.	
Characteristic	Columbia Giant
Diameter at base	1.17 cm
Diameter at midpoint	0.84 cm
Diameter at terminus	0.19 cm
Internode length at base	3.35 cm
Internode length at midpoint	6.32 cm
Internode length at terminus	3.95 cm
Presence of spines further than 0.6 m from the soil surface	Absent
Presence of spines less than 0.6 m from the soil surface	Absent
Floricanes color at base	144A
Floricanes color at midpoint	144A
Floricanes color at terminus	144B
Floricanes lateral length	Medium-long
Floricanes lateral strength	Medium
Primocane color at base	144A with overtones of 185B
Primocane color at midpoint	144A
Primocane color at terminus	146C
Floricanes length (range)	2.12-6.22 m

Table 3 shows primocane foliage characteristics of the new cultivar. Primocane characteristics include mature compound leaf width, mature compound leaf length, number of leaflets per primocane compound leaf, mature leaflet shape, mature leaflet apex, mature leaflet base, mature terminal leaflet width, mature terminal leaflet length, mature first lateral leaflet width, mature first lateral leaflet length, leaflet margin, leaflet serration teeth length, leaflet serration teeth width at base, spine presence on leaves, pubescence on primocane leaflet upper surface, pubescence on primocane leaflet undersurface, primocane leaf color abaxial, primocane leaf color adaxial, petiole length, petiole color: upper surface, petiole color: undersurface, petiolule length termi-

nal leaflet, petiolule length first distal leaflet, petiolule color abaxial, petiolule color adaxial, stipule length, stipule width, and stipule attitude. Fruit is borne only on floricanes.

TABLE 3

Primocane Foliage Characteristics of 'Columbia Giant'.	
Characteristic	Columbia Giant
10 Mature compound leaf width	21.70 cm
Mature compound leaf length	18.50 cm
Number of leaflets per primocane compound leaf	Usually 5
Mature leaflet shape	Ovate
Mature leaflet apex	Broadly acuminate
15 Mature leaflet base	Cordate
Mature terminal leaflet width	8.47 cm
Mature terminal leaflet length	11.88 cm
Mature first lateral leaflet width	6.35 cm
Mature first lateral leaflet length	10.53 cm
Leaflet margin	Double serrate
20 Leaflet serration teeth length	0.31 cm
Leaflet serration teeth width at base	0.28 cm
Spine presence on leaves	Absent
Pubescence on primocane leaflet: upper surface	Light
Pubescence on primocane leaflet: undersurface	Light
25 Primocane leaf color abaxial	137C
Primocane leaf color adaxial	148B
Petiole length	8.68 cm
Petiole color: upper surface	1440 with gradation to 177B
Petiole color: undersurface	145A
Petiolule length: terminal leaflet	2.61 cm
30 Petiolule length: first distal leaflet	0.80 cm
Petiolule color: abaxial	146B
Petiolule color: adaxial	145A
Stipule length	2.11 cm
Stipule width	0.17 cm
Stipule attitude	Crossed near base and curled

Table 4 shows floricanes foliage characteristics of the new cultivar. Floricanes characteristics include mature compound leaf width, mature compound leaf length, number of leaflets per floricanes compound leaf, mature leaflet shape, mature leaflet apex, mature leaflet base, mature terminal leaflet width, mature terminal leaflet length, mature first lateral leaflet width, mature first lateral leaflet length, leaflet margin, leaflet serration teeth length, leaflet serration teeth width at base, pubescence on floricanes leaflet: upper surface, pubescence on floricanes leaflet undersurface, floricanes leaf color abaxial, floricanes leaf color adaxial, petiole length, petiole color adaxial, petiole color abaxial, petiolule length terminal leaflet, petiolule length first distal leaflet, petiolule color abaxial, petiolule color adaxial, stipule length, and stipule width.

TABLE 4

Floricanes Foliage Characteristics of 'Columbia Giant'.	
Characteristic	Columbia Giant
55 Mature compound leaf width	16.67 cm
Mature compound leaf length	9.81 cm
Number of leaflets per floricanes compound leaf	Usually 3
60 Mature leaflet shape	Ovate
Mature leaflet apex	Broadly acuminate
Mature leaflet base	Ovate
Mature terminal leaflet width	5.30 cm
Mature terminal leaflet length	7.85 cm
Mature first lateral leaflet width	4.0 cm
65 Mature first lateral leaflet length	6.55 cm

TABLE 4-continued

Florican Foliage Characteristics of 'Columbia Giant'.	
Characteristic	Columbia Giant
Leaflet margin	Double serrate
Leaflet serration teeth length	0.28 cm
Leaflet serration teeth width at base	0.39 cm
Pubescence on florican leaflet: upper surface	Light
Pubescence on florican leaflet: undersurface	Light
Florican leaf color abaxial	146B
Florican leaf color adaxial	147C
Petiole length	7.22 cm
Petiole color adaxial	144C
Petiole color abaxial	N144D
Petiolule length: terminal leaflet	1.96 cm
Petiolule length: first distal leaflet	0.26 cm
Petiolule color: abaxial	144C
Petiolule color: adaxial	N144D
Stipule length	1.04 cm
Stipule width	0.09 cm

Table 5 shows flower and flowering characteristics of the new cultivar. Flower and flowering characteristics include date 1st bloom, date full bloom, date last bloom, petal color, number flowers per cluster, number of petals per flower, flower diameter, petal length, petal width, number of sepals per flower, peduncle length, rachis length, peduncle color, and cyme type.

TABLE 5

Flower and Flowering Characteristics of 'Columbia Giant'.	
Characteristic	Columbia Giant
Date 1 st bloom	4 May
Date full bloom	11 May
Date last bloom	42145
Petal color	157C
Number flowers per cluster	7.33
Number of petals per flower	5.83
Flower diameter	4.23 cm
Petal length	1.43 cm
Petal width	0.91 cm
Number of sepals per flower	6.17
Peduncle length (cm)	25.69 cm
Rachis length (cm)	9.08 cm
Peduncle color	146B
Cyme type	Simple

Table 6 shows fruit and fruiting characteristics of the new cultivar. Fruit and fruiting characteristics include date 5% of fruit were ripe, date 50% of fruit were ripe, date 95% of fruit were ripe, weight of primary fruit, weight of secondary fruit, weight of tertiary fruit, diameter of primary fruit at equator, diameter of 2° fruit at equator, diameter of 3° fruit at equator, diameter of 1° fruit at poles: tip, diameter of 1° fruit at poles: base, diameter of 2° fruit at poles: tip, diameter of 2° fruit at poles: base, diameter of 3° fruit at poles: tip, diameter of 3° fruit at poles: base, berry length primary fruit, berry length 2° fruit, berry length 3° fruit, ratio of primary fruit length to width, shape description, uniformity of berry shape, color when full ripe, number of drupelets per fruit, drupelet weight, individual seed weight, glossiness, firmness, flavor, texture of fruit when chewed, drupelet skin resistance to abrasion, ease of separation of fruit from pedicel, machine harvestability, resistance to heat damage of fruit, berries per inflorescence—mean, berries per inflorescence range, soluble solids (%; in brix), pH, titratable acidity (% as citric acid), and yield (actual kg·plt⁻¹).

TABLE 6

Fruit and Fruiting Characteristics of 'Columbia Giant'.	
Characteristic	Columbia Giant
Date 5% of fruit were ripe	25 June
Date 50% of fruit were ripe	5 July
Date 95% of fruit were ripe	19 July
Weight of primary fruit	14.07 g
Weight of secondary fruit	13.27 g
Weight of tertiary fruit	14.47 g
Diameter of primary fruit at equator	2.14 cm
Diameter of 2° fruit at equator	2.00 cm
Diameter of 3° fruit at equator	1.97 cm
Diameter of 1° fruit at poles: tip	1.17 cm
Diameter of 1° fruit at poles: base	1.90 cm
Diameter of 2° fruit at poles: tip	1.31 cm
Diameter of 2° fruit at poles: base	2.08 cm
Diameter of 3° fruit at poles: tip	1.32 cm
Diameter of 3° fruit at poles: base	1.91 cm
Berry length primary fruit	5.02 cm
Berry length 2° fruit	4.67 cm
Berry length 3° fruit	4.71 cm
Ratio of primary fruit length to width	2.36
Shape description	Long conic
Uniformity of berry shape	Excellent
Color when full ripe	Mostly 203B with N186C on ~5% drupelets
Number of drupelets per fruit	181.70
Drupelet weight	77.0 mg
Individual seed weight	26.9 mg
Glossiness	Medium glossy to dull
Firmness	Very good
Flavor	Very good
Texture of fruit when chewed	Excellent
Drupelet skin resistance to abrasion	Good
Ease of separation of fruit from pedicel	Easy
Machine harvestability	Excellent
Resistance to heat damage of fruit	Medium to good
Berries per inflorescence - mean	7.00
Berries per inflorescence range	5-10
Soluble solids (%; in Brix)	11.25
pH	3.26
Titratable acidity (% as citric acid)	18.5
Yield (actual kg · plt ⁻¹)	6.86
Disease Response	Under a typical, minimal, disease management program does not exhibit any particular disease problems

COMPARISON WITH PARENTAL AND COMMERCIAL VARIETIES

'Columbia Giant' differs from the female parent 'NZ 9629-1' (unpatented) in that 'Columbia Giant' has medium-sized, glossy, very large fruit (12-17 g), while 'NZ 9629-1' has slightly pubescent fruit that are smaller (4.5 g).

'Columbia Giant' differs from the male parent blackberry plant 'ORUS 1350-2' (unpatented) in that it is spineless, has conic berries with excellent flavor, while 'ORUS 1350-1' (unpatented) is spiny and has barrel shaped berries with poor flavor.

'Columbia Giant' differs from the commercial variety 'Marion' (unpatented) in that 'Columbia Giant' is spineless and has large yields of very large, firm, and very uniformly shaped fruit while 'Marion' is spiny and bears medium yields of medium sized, and soft fruit that are unevenly shaped. 'Columbia Giant' differs from the commercial variety 'Black Diamond' (unpatented) in that 'Columbia Giant' carries the 'Lincoln Logan' (unpatented) source of spinelessness and therefore the canes are completely spineless and the plants are vigorous with fruit that are very large and have excellent, aromatic flavor, while 'Black Diamond'

(unpatented) carries the 'Austin Thornless' (unpatented) source of spinelessness and so has spines on the base of the canes and the plants are not vigorous and they produce large fruit with a mild flavor. 'Columbia Giant' differs from the commercial variety 'Columbia Star' (U.S. Plant patent application Ser. No. 13/815,074, now U.S. Plant Pat. No. 25,532) based predominantly on a fruit size. 'Columbia Giant' fruit ranged from 12.2-16.3 g with mean primary fruit size of 14.1 g while 'Columbia Star' fruit ranged from 6.0-10.5 g with mean primary fruit size of 9.7 g.

The market use of Columbia Giant is primarily local fresh such as local farmers markets but it machine harvests well and processes well.

I claim:

1. A new and distinct cultivar of blackberry plant, substantially as illustrated and described, characterized by its high yields of extremely large and very uniformly shaped berries with very good firmness, color, and flavor and that are borne on vigorous, completely thornless trailing plants.

* * * * *



Fig. 1



Fig. 2




Fig. 3



Fig. 4