



(12) **United States Plant Patent**
van den Haak

(10) **Patent No.:** **US PP28,809 P3**
(45) **Date of Patent:** **Dec. 26, 2017**

(54) **PHLOX PLANT NAMED ‘APPOTTPI’**

(50) Latin Name: *Phlox paniculata*
Varietal Denomination: **Appottpi**

(71) Applicant: **Jelle van den Haak**, Amsterdam (NL)

(72) Inventor: **Jelle van den Haak**, Amsterdam (NL)

(73) Assignee: **Gootjes-Allplant B.V.**, Schagen (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 15 days.

(21) Appl. No.: **14/999,228**

(22) Filed: **Apr. 11, 2016**

(65) **Prior Publication Data**

US 2017/0295703 P1 Oct. 12, 2017

(51) **Int. Cl.**
A01H 5/02 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./320**

(58) **Field of Classification Search**
USPC Plt./320
See application file for complete search history.

Primary Examiner — Susan McCormick Ewoldt

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Phlox* plant named ‘Appottpi’, characterized by its broadly upright and relatively compact plant habit; freely flowering habit, numerous small dark pink-colored flowers; long flowering period; and good garden performance.

2 Drawing Sheets

1

Botanical designation: *Phlox paniculata*.
Cultivar denomination: ‘APPOTTPI’.

CROSS-REFERENCED TO CLOSELY-RELATED APPLICATIONS

Title: *Phlox* Plant Named ‘APPOTTWH’
Applicant: Jelle van den Haak
Filed: Concurrently with this application as Ser. No. 14/999,226

Title: *Phlox* Plant Named ‘APPOTWWH’
Applicant: Jelle van den Haak
Filed: Concurrently with this application as Ser. No. 14/999,225

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Phlox* plant, botanically known as *Phlox paniculata* and hereinafter referred to by the name ‘Appottpi’.

The new *Phlox* plant is a product of a planned breeding program conducted by the Inventor in Andijk and Hazerswoude-Dorp, The Netherlands. The objective of the breeding program was to create new compact and vigorous *Phlox* plants with numerous small flowers.

The new *Phlox* plant originated from a cross-pollination in August, 2012 in Andijk, The Netherlands, of a proprietary selection of *Phlox paniculata* identified as code number 047-11-K121-05, not patented, as the female, or seed, parent, with a proprietary selection of *Phlox paniculata* identified as code number 047-08-007, not patented, as the male, or pollen, parent. The new *Phlox* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled environment in Hazerswoude-Dorp, The Netherlands in August, 2013.

Asexual reproduction of the new *Phlox* plant by terminal cuttings in a controlled greenhouse environment in Andijk, The Netherlands since 2014 has shown that the unique

2

features of this new *Phlox* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

5

Plants of the new *Phlox* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

10

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Appottpi’. These characteristics in combination distinguish ‘Appottpi’ as a new and distinct *Phlox* plant:

15

1. Broadly upright and relatively compact plant habit.
2. Freely flowering habit, numerous small dark pink-colored flowers.
3. Long flowering period.
4. Good garden performance.

20

Plants of the new *Phlox* and the female parent selection differ primarily in flower color as flowers of plants of the female parent selection are purple in color.

25

Plants of the new *Phlox* and the male parent selection differ primarily in the following characteristics:

1. Plants of the new *Phlox* have smaller flowers than plants of the male parent selection.
2. Plants of the new *Phlox* and the male parent selection differ in flower color as plants of the male parent selection have white-colored flowers with dark pink-colored centers.

30

Plants of the new *Phlox* can be compared to plants of *Phlox paniculata* ‘Appottwh’, disclosed in a U.S. Plant Patent application filed concurrently and *Phlox paniculata* ‘Appotwwh’, disclosed in a U.S. Plant Patent application filed concurrently. Plants of the new *Phlox* differ primarily from plants of ‘Appottwh’ and ‘Appotwwh’ in flower color as plants of ‘Appottwh’ have purplish white-colored flowers and plants of ‘Appotwwh’ have white-colored flowers.

35

Plants of the new *Phlox* can also be compared to plants of *Phlox paniculata* 'Barseventyone', disclosed in U.S. Plant Pat. No. 23,262. In side-by-side comparisons, plants of the new *Phlox* and 'Barseventyone' differ in the following characteristics:

1. Plants of the new *Phlox* are more vigorous than plants of 'Barseventyone'.
2. Plants of the new *Phlox* have larger flowers than plants of 'Barseventyone'.
3. Plants of the new *Phlox* and 'Barseventyone' differ in flower color as plants of 'Barseventyone' have purple-colored flowers.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Phlox* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Phlox* plant.

The photograph on the first sheet is a side perspective view of a typical flowering plant of 'Appottpi' grown in a container.

The photographs on the second sheet are close-up views of typical inflorescences (upper photograph) and typical leaves (lower photograph) of 'Appottpi'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations and measurements describe plants grown during the summer in 15-cm containers in an outdoor nursery in Mijdrecht, The Netherlands and under cultural practices typically used in commercial *Phlox* production. During the production of the plants, day temperatures ranged from 10° C. to 25° C. and night temperatures ranged from 4° C. to 15° C. Plants were pinched five weeks after planting and were 19 weeks old when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Phlox paniculata* 'Appottpi'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Phlox paniculata* identified as code number 047-11-K121-05, not patented.

Male, or pollen, parent.—Proprietary selection of *Phlox paniculata* identified as code number 047-08-007, not patented.

Propagation:

Type.—By terminal cuttings.

Time to initiate roots, summer.—About 12 days at temperatures about 20° C.

Time to initiate roots, winter.—About 16 days at temperatures about 20° C.

Time to produce a rooted young plant, summer.—About 36 days at temperatures about 18° C.

Time to produce a rooted young plant, winter.—About 42 days at temperatures about 18° C.

Root description.—Medium in thickness, fleshy; typically white in color, actual color of the roots is dependent on substrate composition, water quality,

fertilizer type and formulation, substrate temperature and physiological age of roots.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Herbaceous perennial; broadly upright and relatively compact plant habit; overall shape, broadly obovate; low vigor to moderately vigorous in growth habit.

Plant height.—About 34.8 cm.

Plant width (spread).—About 32.5 cm.

Lateral branches.—Quantity: About seven primary lateral branches per plant. Length: About 15.1 cm. Diameter: About 5.5 mm. Internode length: About 2.1 cm. Strength: Strong. Aspect: Upright to about 45° from vertical. Texture and luster: Smooth, glabrous; slightly glossy. Color: Close to 144A to 144B.

Leaf description:

Arrangement.—Opposite, simple.

Length.—About 10.2 cm.

Width.—About 3.7 cm.

Shape.—Narrowly ovate to elliptic; slightly to moderately carinate.

Apex.—Long apiculate.

Base.—Truncate.

Margin.—Entire; very finely serrate, inconspicuous.

Texture and luster, upper and lower surfaces.—Glabrous; slightly rugose; slightly glossy.

Venation pattern.—Pinnate.

Color.—Developing leaves, upper surface: Close to between 137C and 143A. Developing leaves, lower surface: Close to 138B. Fully expanded leaves, upper surface: Close to N137B; venation, close to 144A. Fully expanded leaves, lower surface: Close to 147B; venation, close to 144B.

Petioles.—Length: About 3 mm. Diameter: About 3 mm by 4 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 144B; towards the margins, moderately tinged with close to 187A.

Flower description:

Flower type and flowering habit.—Single rotate and salverform flowers arranged in compound terminal panicles; flowers face upright to outwardly; panicles roughly hemispherical in shape; freely flowering habit with about 330 flowers developing per inflorescence and about 1,300 flowers developing per plant during the flowering season.

Fragrance.—Moderately fragrant; sweet, pleasant.

Natural flowering season.—Plants begin flowering about nine months after planting; long flowering period, plants flower continuously from July through September in The Netherlands.

Flower longevity.—Flowers last about ten days on the plant; flowers not persistent.

Flower buds.—Height: About 1.5 cm. Diameter: About 4 mm. Shape: Narrowly elliptic. Color: Close to N74C; developing calyx, close to 148D heavily flushed and tinged with close to 200B.

Inflorescence height.—About 15.8 cm.

Inflorescence diameter.—About 16.4 cm.

Flower diameter.—About 2.4 cm.

Flower depth.—About 1.5 cm.

Petals.—Quantity per flower: Typically five in a single whorl; petals fused at the base into a narrow tube; free parts separate to slightly imbricate. Length:

Overall, about 2.1 cm; lower fused portion, about 1 cm. Lobe width: About 1 cm. Shape: Free part, spatulate. Apex: Obtuse. Margin: Entire. Texture and luster, upper surface: Smooth, glabrous; moderately velvety; matte. Texture and luster, lower surface: Smooth, glabrous; slightly glossy. Color: When opening and fully opened, upper surface: Close to N66C; towards the throat, close to N66A; throat, close to N79C; with development, main color becoming closer to N66D and towards the throat, close to N66B. When opening and fully opened, lower surface: Close to 68C to 68D flushed with close to 70B; tube, close to N79C.

Sepals.—Quantity per flower: Typically five in a single whorl, fused towards the base; calyx, campanulate. Length: About 9 mm. Width: About 1 mm. Shape: Lanceolate. Apex: Narrowly apiculate. Margin: Entire. Texture and luster, upper and lower surfaces: Smooth, glabrous; slightly glossy. Color: When opening and fully opened, upper surface: Close to 148D. When opening and fully opened, lower surface: Close to 148D heavily flushed and tinged with close to 200B.

Peduncles.—Length, primary peduncles: About 12.6 cm. Diameter, primary peduncles: About 3 mm. Length, secondary peduncles: About 5.8 cm. Diameter, secondary peduncles: About 1.5 mm. Aspect, primary peduncles: Erect. Aspect, secondary

peduncles: About 40° from vertical. Strength: Strong. Texture: Smooth, glabrous. Color: Close to 144A.

Pedicels.—Length: About 4 mm. Diameter: About 1 mm. Angle: About 40° from the peduncle axis. Strength: Strong. Texture: Smooth, glabrous. Color: Close to 144B.

Reproductive organs.—Stamens: Quantity per flower: Typically five; filaments fused with petals. Filament length: About 0.75 mm. Filament color: Close to NN155C. Anther length: About 1 mm. Anther shape: Oblong; basifixed. Anther color: Close to 164C. Pollen amount: Scarce. Pollen color: Close to 11D. Pistils: Quantity per flower: One. Pistil length: About 1.1 cm. Stigma shape: Cleft, three-parted. Stigma color: Close to 145D. Style length: About 1 cm. Style color: Close to 150D. Ovary color: Close to 143B.

Seeds and fruits.—Seed and fruit development have not been observed on plants of the new *Phlox*.

Garden performance: Plants of the new *Phlox* have been observed to have good garden performance and tolerate rain, wind, high temperatures about 35° C. and to be winter hardy to USDA Hardiness Zone 6.

Pathogen & pest resistance: Plants of the new *Phlox* have been not been observed to be resistant to pathogens and pests common to *Phlox* plants.

It is claimed:

1. A new and distinct *Phlox* plant named 'Appottpi' as illustrated and described.

* * * * *



