

(12) **United States Plant Patent**
Park

(10) **Patent No.:** **US PP29,145 P3**
(45) **Date of Patent:** **Mar. 20, 2018**

(54) **COREOPSIS PLANT NAMED ‘URIBLO2’**

CPC A01H 5/025; A01H 5/02; A01H 5/00
See application file for complete search history.

(50) Latin Name: **Coreopsis**
rosea×**Coreopsis verticillata**
Varietal Denomination: **URIBLO2**

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP15,455 P3 * 12/2004 Faria Plt./417

OTHER PUBLICATIONS

Jala Morphological Change Due to Effects of Acute Gamma Ray on
Wishbone Flower (*Torenia fourmieri*) in Vitro, 2011 International
Transaction Journal of Engineering, Management, & Applied Sci-
ences & Technologies, vol. 2 No. 4, 375-383.*

Park et al. Development of a New *Coreopsis* Variety ‘Uridream
Pink’ by Gamma-ray Irradiation, Kor. J. Hort. Sci. Technol.
32(6):906-911, 2014.*

English translation of Park et al. 2017, 16 pp.*

* cited by examiner

Primary Examiner — June Hwu

(74) Attorney, Agent, or Firm — Penny J. Aguirre

(57) **ABSTRACT**

A new cultivar of hybrid *Coreopsis* plant named
‘URIBLO2’, that is characterized by its dwarf plant habit, by
its large inflorescences, its ray florets ray florets that are light
red-purple in color with a yellow base, and its disk florets
that are orange-red in color.

2 Drawing Sheets

1

Botanical Classification: *Coreopsis rosea* x *Coreopsis*
verticillata.

Variety Denomination: ‘URIBLO2’.

CROSS REFERENCE TO RELATED
APPLICATIONS

This application is co-pending with U.S. Plant Patent
Applications filed for plants derived from the same breeding
programs that is entitled *Coreopsis* Plant Named
‘URIBLO1’ (U.S. Plant patent application Ser. No. 14/999,
120).

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Coreopsis* plant, botanically of hybrid origin and known
as *Coreopsis* ‘URIBLO2’ referred to hereinafter by its
cultivar name, ‘URIBLO2’. The new cultivar of *Coreopsis*
is an herbaceous perennial grown for landscape and con-
tainer use.

The new invention arose from an ongoing controlled
breeding program in Icheon-si, Korea. The objective of the
breeding program is to develop hybrid cultivars of *Coreopsis*
with unique flower colors and dwarf plant habit.

2

The new cultivar arose as a mutation using gamma
radiation of rooted cuttings of unknown, unnamed propri-
etary plants from the inventor’s breeding program in January
of 2010. ‘Limerock Ruby’ (U.S. Plant Patent No. 15,455)
and addition cultivars were initially used in the breeding
program in prior years to obtain initial primary mutations
and subsequent rounds of gamma radiation were performed
on proprietary seedlings until ‘URIBLO2’ was selected in
September 2010 as a single unique plant amongst the
resulting seedlings.

Asexual propagation of the new cultivar was first accom-
plished by stem cuttings under the direction of the inventor
in Icheon-si, Korea in September of 2010. Asexual propa-
gation by stem cuttings has determined that the character-
istics of the new cultivar are stable and are reproduced true
to type in successive generations.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
represent the characteristics of the new cultivar. These
attributes in combination distinguish ‘URIBLO2’ as unique
cultivar of *Coreopsis*.

1. ‘URIBLO2’ exhibits a dwarf plant habit.
2. ‘URIBLO2’ exhibits large inflorescences.

3. 'URIBLO2' exhibits ray florets that are light red-purple in color with a yellow base.

4. 'URIBLO2' exhibits disk florets that are orange-red in color.

'URIBLO2' can be most closely compared to 'URIBLO1', 'Limerock Ruby' and 'Uridream Pink' (not patented). 'URIBLO1' is similar to 'URIBLO2' in having a dwarf plant habit. 'URIBLO1' differs from 'URIBLO2' in having any ray florets that are crimson red in color, in having a slightly smaller plant size, and in having inflorescences that are slightly smaller in size. 'Limerock Ruby', differs from 'URIBLO2' in having a taller plant height and wider plant spread, in having inflorescences that are smaller in size with ray florets that are smaller in size. 'Uridream Pink' differs from 'URIBLO2' in having a taller plant height, longer internode lengths, smaller inflorescences, and ray florets that are much darker pink in color.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying colored photographs illustrate the overall appearance and distinct characteristics of the new *Coreopsis*. The photographs were taken of a 3 month-old plant of 'URIBLO2' as grown outdoors in BosKoop, The Netherlands.

The photograph in FIG. 1 provides a side view of a plant of 'URIBLO2' in bloom.

The photograph in FIG. 2 provides a close-up view of the inflorescences of 'URIBLO2'.

The photograph in FIG. 3 provides a close-up view of the foliage of 'URIBLO2'.

The colors in the photograph are as close as possible with the photographic and printing technology utilized and the color values cited in the Detailed Botanical Description accurately describe the colors of the new *Coreopsis*.

DETAILED BOTANICAL DESCRIPTION

The following is a detailed description of 3 month-old plants of the new cultivar as grown outdoors in 13-cm containers Boskoop, The Netherlands. The phenotype of the new cultivar may vary with variations in environmental, climatic, and cultural conditions, as it has not been tested under all possible environmental conditions. The color determination is in accordance with the 2007 RHS Colour Chart of the Royal Horticultural Society, London, England, except where general color terms of ordinary dictionary significance are used.

General Description:

Blooming period.—Early summer to late summer in The Netherlands.

Plant type.—Herbaceous perennial.

Plant habit.—Dwarf and upright.

Height and spread.—An average of 23.7 cm in height and 26.4 cm in width.

Cold hardiness.—At least to U.S.D.A Zone 4.

Diseases and Pests.—No susceptibility or resistance to diseases or pests has been observed.

Root description.—Fibrous when young, becoming fleshy with age, 162D in color.

Propagation.—Stem cuttings.

Root development.—An average of 10 days for root initiation and about 5 weeks to produce a young rooted plant in a 10-cm container.

Growth rate.—Moderate.

Stem Description:

Shape.—Oval and slightly ridged.

Stem color.—143B.

Stem size.—An average of 7.3 cm in length (excluding inflorescence) and 1.5 mm in diameter.

Stem surface.—Glabrous, glossy, slightly axially ribbed.

Stem aspect.—Upward and outward.

Stem strength.—Strong.

Branching habit.—Well-branched, an average of 38 lateral branches.

Internode length.—An average of 2.4 cm.

Foliage Description:

Leaf type.—Single.

Leaf division.—Simple.

Leaf margins.—Entire to two-lobed.

Leaf size.—Variable; up to 4.3 cm in length and 1.7 cm in width when entire and up to 5.4 cm in length and 4 mm in width when two-lobed.

Leaf shape.—Narrow lanceolate.

Leaf base.—Narrow curieate

Leaf apex.—Acute.

Leaf venation.—Pinnate, color; upper surface 137B and lower surface 144A.

Leaf attachment.—Sessile.

Leaf arrangement.—Opposite.

Leaf surface.—Both surfaces dull and glabrous.

Leaf color.—Young and mature upper surface; 137B and young and mature lower surface; 143A.

Leaf number.—An average of 8 leaves (4 pairs) per stem.

Inflorescence Description:

Inflorescence type.—Terminal capitulate, with a single row of ray florets surrounding disk florets in the center.

Lastingness of inflorescence.—An average of 3 weeks, self-cleaning.

Fragrance.—Faint, of musk and straw.

Quantity of inflorescence size.—An average of 1 per lateral branch.

Inflorescence size.—An average of 4.0 cm in diameter and 1.2 cm in height, disk portion an average of 7 mm in diameter.

Inflorescence buds.—An average of 5 mm in depth and in diameter, globular in shape, color; 147C with apex, 168B to 168C.

Peduncle.—Primary; on average of 7.4 cm in length and 1 mm in width, held straight upright, secondary; an average of 5.2 cm in length and 1 mm in width, held at an average angle of 30° to primary peduncle, slightly glossy and glabrous on all surfaces and 143B in color.

Involucral bracts:

Bract number.—Two rows of 8.

Bract arrangement.—Rotate.

Bract size.—Outer bracts; an average of 3 mm in length and 2 mm in width, inner bracts; an average of 7 mm in length and 4 mm in width.

Bract color.—Inner bracts; 24A with base 146A and outer bracts; 143A.

Bract texture.—Glabrous and slightly glossy on outer and inner surfaces of outer and inner bracts.

Bract apex.—Acute on outer and inner bracts.

Bract base.—Broadly cuneate on outer and inner bracts.

Bract margins.—Entire.

Bract shape.—Outer bracts; narrow ovate, inner bracts; ovate.

Ray Florets (sterile):

Number.—An average of 8 arranged primarily in one row,

Shape.—Obovate.

Size.—An average of 1.8 cm in length and 9 mm in width.

Apex.—Irregularly lobed with an average of 3 acute lobes, center lobe is about 2 mm longer than the lateral lobes.

Base.—Rounded.

Margins.—Entire with apex lobed.

Aspect.—Held horizontal.

Texture.—Both surfaces; glabrous and dull, upper surface very slightly velvety and lower surface moderately carinate.

Color.—Upper surface when opening; 70B to 70C with base 3C to 3D, lower surface when opening; 73D with apex 75A to 75B and base 2D, upper surface when fully open; 75B to 75C marbled 76C to 76D with base 2D, lower surface when fully open; 158D suffused with 75D base 2D, color not fading.

Petaloids.—Ray floret like; average of 4 per inflorescence, an average of 5 mm in length and 2 mm in width, narrow obovate to linear in shape, rounded to acute apex, cuneate base, coloration matches ray floret color.

Disk florets (perfect):

Shape.—Tubular, spirally arranged on disc.

Apex.—Acute.

Base.—Lower 87.5% fused into tube.

Number.—About 80.

Size.—An average of 4.5 mm in length and 1.0 mm in width at the apex and 0.4 mm in width at the base.

5 Color.—En masse; a mix of 15B to 15C and 179A, corolla; base of tube is 15B to 15C in color, flared portion is 179A.

Texture.—Both surfaces; glabrous and dull.

Receptacle.—An average of 1.5 mm in diameter and 1 mm in depth, 143B in color.

10 Reproductive Organs:

Presence.—Disk florets are perfect, ray florets are sterile.

Gynoecium.—Pistil; 1, an average of 3 mm in length, style; an average of 2.5 mm in length and 16A in color, stigma; decurrent and 23A in color, ovary is 151C in color.

15 Androecium.—Stamens; 5, anther; basiFixed and linear in shape, an average of 1.5 mm in length and 203D in color, pollen; moderate in quantity and 17B in color.

Fruit/seed.—No fruit or seed development was observed.

20 It is claimed:

1. A new and distinct cultivar of *Coreopsis* plant named 'URIBLO2' as herein illustrated and described.

* * * * *



FIG. 1



FIG. 2



FIG. 3