



US00PP29787P3

(12) **United States Plant Patent**
Finn

(10) **Patent No.:** **US PP29,787 P3**
(45) **Date of Patent:** **Nov. 6, 2018**

(54) **BLUEBERRY PLANT NAMED ‘ECHO’**

(50) Latin Name: *Vaccinium* spp. section *Cyanococcus*
L.
Varietal Denomination: **Echo**

(71) Applicant: **The United States of America, as
Represented by the Secretary of
Agriculture**, Washington, DC (US)

(72) Inventor: **Chad E. Finn**, Corvallis, OR (US)

(73) Assignee: **The United States of America, as
Represented by the Secretary of
Agriculture**, Washington, DC (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/530,947**

(22) Filed: **Mar. 28, 2017**

(65) **Prior Publication Data**
US 2018/0288913 P1 Oct. 4, 2018

(51) **Int. Cl.**
A01H 5/08 (2018.01)
A01H 6/36 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./157**
CPC *A01H 6/368* (2018.05)

(58) **Field of Classification Search**
USPC Plt./157
See application file for complete search history.

Primary Examiner — Susan McCormick Ewoldt

(74) *Attorney, Agent, or Firm* — John Fado; Ariel
Atkinson

(57) **ABSTRACT**

‘Echo’ is a new and distinct repeat flowering, ornamental
blueberry (*Vaccinium* section *Cyanococcus* L.) cultivar that
originated from seed produced from the cross of the female
blueberry plant ‘Perpetua’ (U.S. Plant Pat. No. 24,209) and
the male blueberry plant ORUS 55-1 (unpatented). ‘Echo’ is
a blueberry clone distinguished by its ornamental charac-
teristics that include a spreading bush habit with glossy
leaves, repeat flowering and fruiting, and small, mild fla-
vored dark blue fruit.

3 Drawing Sheets

1

Latin name of genus and species of the plant claimed:
‘ECHO’ is a new blueberry plant that is a genus *Vaccinium*
spp. section *Cyanococcus* L.

Variety denomination: The new blueberry plant claimed is
of the cultivar denomination ‘Echo’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct blue-
berry cultivar botanically known as *Vaccinium* section *Cya-
nococcus* L., and herein referred to as ‘Echo’, as herein
described and illustrated.

‘Echo’ is a blueberry clone distinguished by its ornamental
characteristics that include a spreading bush with glossy
leaves, repeat flowering and fruiting, and small, mild fla-
vored dark blue fruit. This new blueberry cultivar was
developed in Corvallis, Oreg. in July 2010 and originated
from the cross of the female blueberry plant ‘Perpetua’ (U.S.
Plant Pat. No. 24,209) and the male blueberry plant ORUS
55-1 (unpatented) made in Corvallis, Oreg. in 2006. Both
parents are repeat flowering and their ancestry can be traced
in part to selections CVAC 45 (PI 296412), which was
collected from the wild in 1963 in Monmouth, Me. and
CVAC (296397), which was collected from the wild in 1964
in Farmington, Me., and are listed as *Vaccinium corymbo-
sum* L. (highbush blueberry) by the USDA—ARS, National
Genetic Resources Program, Germplasm Resources Infor-
mation Network—(GRIN) (Online Database). Since the
wild germplasm ancestry originated from where the ranges
of *V. corymbosum* and *V. angustifolium* Aiton (lowbush
blueberry) overlap and since many of the plant’s character-
istics for fruit size, leaf shape, and leaf size are intermediate
between these two species, ‘Echo’ is presumed to contain

2

both species in its pedigree. ‘Echo’s overwintering buds
break in the spring, flower and produce a crop like most
blueberry plants. ‘Echo’, as with its ‘Perpetua’ (U.S. Plant
Pat. No. 24,209) maternal parent, is unique in that the new
growth produces flower buds that proceed to break bud
without winter dormancy and then flower and ripen a
late-summer into fall crop. While it is not uncommon for
blueberry genotypes to produce a few fruit in the fall on 1-2
buds at the tip of the new growth, these repeat flowering
genotypes will flower on up to 12-16 nodes on the new
growth. The original seedling of the new cultivar was
asexually propagated at a nursery in Benton County, Oreg.
The new cultivar was established in vitro from a stem cutting
and microcuttings have been taken and rooted from this sort
of culture. The present invention has been found to be stable
and reproduce true to type through successive asexual
propagations.

SUMMARY OF THE INVENTION

The following are the most outstanding and distinguish-
ing characteristics of this new cultivar when grown under
normal horticultural practices in Oregon. 1. Ability to flower
multiple times in the season on overwintered canes and on
new growth; 2. Attractive combination of flowers and fruit
in late summer and fall; 3. Green, glossy, attractive leaves;
4. Edible dark blue fruit; and 5. Vigorous, spreading plant
habit.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

This new blueberry plant is illustrated by the accompa-
nying color photographs that show typical specimens of the

flower clusters in bloom (FIG. 1), ripe fruit on a four-year-old plant (FIG. 2), and the entire plant with a ripening crop in late August (2nd crop) (FIG. 3); the colors shown are as true as can be reasonably obtained by conventional photographic procedures.

DETAILED DESCRIPTION OF THE NEW CULTIVAR

The following detailed description of 'Echo' is based on observations taken from 2013 to 2016 in trials planted in 2012 in Corvallis and Aurora, Oreg. The measurements/observations for one trait always being made on samples from the same location. This description is in accordance with UPOV terminology. Color designations, color descriptions and other phenotypical descriptions may deviate from the stated values and descriptions depending upon variation in environmental, seasonal, climatic and cultural conditions. 'Echo' has not been observed under all possible environmental conditions. Color terminology follows The Royal Horticultural Society Colour Chart. London (R.H.S.) (5th edition, 2007).

Table 1 shows most the bush characteristics of the new cultivar. Characteristics include plant size, height of plant, canopy diameter, growth habit, plant vigor, twigginess, suckering, tendency towards evergreenness, surface texture 1 year-old-wood, surface texture 3 year-old-wood, color new wood, color rough bark 1 year-old-wood, color rough bark 3 year-old-wood, internode length and winter hardiness in Aurora, Oreg. (45° 16' 49" N/122° 44' 50" W).

TABLE 1

Characteristic	Echo
Plant size	Small; comparable to 'Perpetua'
Height of plant	0.88 m
Canopy diameter	0.90 m
Growth habit	Upright and spreading compared to 'Perpetua'
Plant vigor	Moderate vigor, comparable to 'Perpetua'
Twigginess	Moderate
Suckering	None
Tendency towards evergreenness	Deciduous but holds leaves into winter
Surface texture new wood	Smooth, slightly pubescent
Surface texture 1 year-old-wood	Smooth, slightly pubescent
Surface texture 3 year-old-wood	Rough
Color new wood	Gray-Red group 180A
Color rough bark 1 year-old-wood	Gray-Red group 178B
Color rough bark 3 year-old-wood	Greyed-Orange group 177A
Internode length	1.84 cm
Winter tolerance in Aurora, Oregon (45° 16' 49" N/122° 44' 50" W)	Excellent

Table 2 shows the foliage characteristics of the new cultivar. Characteristics include leaf length, leaf width, leaf shape, leaf margin, leaf color upper leaf surface, leaf color lower leaf surface, pubescence upper leaf surface, pubescence lower leaf surface, pubescence leaf margins, and leaf venation.

TABLE 2

Characteristic	Echo
Leaf length	4.72 cm
Leaf width	1.89 cm
Leaf shape	Elliptic

TABLE 2-continued

Characteristic	Echo
Leaf margin	Entire
Leaf color upper leaf surface	Green group 139A
Leaf color lower leaf surface	Green group 138B
Pubescence upper leaf surface	None visible
Pubescence lower leaf surface	None visible
Pubescence leaf margins	None visible
Leaf venation	Net
Shape of leaf apex	Acuminate
Shape of leaf base	Acute
Petiole length	1.84 mm
Petiole diameter	1.23
Petiole color	Yellow-Green Group (145B)

Table 3 shows the flower characteristics of the new cultivar. Characteristics include fragrance, pedicel length flower—spring, peduncle length flower—spring, flower shape, pollen abundance, pollen color, flower length, corolla length, corolla diameter, style length, calyx diameter, corolla aperture, calyx surface, corolla color, corolla texture, calyx color, pistil color, pedicel color, peduncle color, flowering cluster, spring flowering period (mean date ~5%), spring flowering period (mean date ~50%), summer flowering period (mean date ~5%), number flowers per cluster, Fruit set when pollinated with bulk pollen of other cultivars, Fruit set when pollinated with own pollen (selfed), stigma location (distance from corolla aperture to stigma), pollen abundance, pollen color, calyx aperture, calyx lobes, calyx depth and flower arrangement.

TABLE 3

Characteristic	Echo
Fragrance	None
Pedicel length flower - spring	4.90 mm
Peduncle length flower - spring	7.66 mm
Flower shape	Urceolate
Pollen abundance	High
Pollen color	Yellow group 10D
Flower length	8.8 mm
Corolla length	7.6 mm
Corolla diameter	5.0 mm
Style length	7.1 mm
Calyx diameter	4.5 mm
Corolla aperture	3.4 mm
Calyx surface	Smooth
Corolla color	White group 155A
Corolla texture	Smooth
Calyx color	Yellow-Green group 146B
Pistil color	Yellow-Green group 145A
Pedicel color	Yellow-Green group 146C
Peduncle color	Yellow-Green group 145A
Flowering cluster	Loose
Spring flowering period (mean date ~5%)	Mar. 18
Spring flowering period (mean date ~50%)	Apr. 1
Summer flowering period (mean date ~5%)	Jul. 7
Number flowers per cluster	7.3
Fruit set when pollinated with bulk pollen of other cultivars	95.4%
Fruit set when pollinated with own pollen	55.2%
Stigma location	Below corolla aperture, inside corolla
Pollen abundance	High
Pollen color	Yellow group 10D
Calyx aperture	2.48 mm
Calyx lobes	2.26 mm
Calyx depth	1.75 mm
Flower arrangement	Monopodial branching

Table 4 shows the mature fruit characteristics of the new cultivar. Characteristics include berry cluster, berry

peduncle length, berry pedicel length, berry detachment from pedicel, berry weight, berry height, berry width, berry height/width ratio, berry shape, berry skin color on plant, berry skin color after harvest, berry skin color after polishing, berry wax, berry flesh color, pedicel scar, berry firmness, berry flavor, berry texture, color of dried seeds, length of well-developed dried seed, width of well-developed dried seed, weight of well-developed dried seed, productivity and size of 1st (spring) vs 2nd (summer) crop.

TABLE 4

Characteristic	Echo
Berry cluster	Loose
Berry peduncle length	26.73 mm
Berry pedicel length	5.71 mm
Berry detachment from pedicel	Slight resistance to pull
Berry weight	0.75 g
Berry height	12.21 mm
Berry width	11.94 mm
Berry height/width ratio	1.12
Berry shape	Subglobose
First ripe (5%) fruit first (spring) crop	May 24
First ripe (5%) fruit first (spring) crop	Aug. 19
Berry skin color on plant when ripe	Violet-Blue group N92C
Berry skin color after harvest	Violet-Blue group N92C
Berry skin color after polishing harvested fruit	Blue group 103A
Berry wax	Present
Berry flesh color	Green group 139D
Pedicel scar	1.59 mm
Berry firmness	Moderately firm
Berry flavor	Mild
Berry texture	Firm, do not notice seeds
Color of dried seeds	Greyed-Orange group 177C
Length of well-developed dried seed	1.76 mm
Width of well-developed dried seed	0.84 mm
Weight of well-developed dried seed	0.40 mg
Productivity	Moderate; comparable to 'Perpetua'

TABLE 4-continued

Characteristic	Echo
Size of 1st (spring) vs 2nd (summer) crop	2nd crop is larger than first

Resistance to diseases and insects: In a minimal spray program with dormant sprays only, 'Echo' has had no fungal or bacterial disease symptoms and no indication of problems with gall midge. 'Echo' plants have not yet shown symptoms or tested positive for Blueberry shock virus; if it was an extremely susceptible (e.g. 'Bluegold' (unpatented)), it would be showing symptoms and testing positive after 5 years in the field, however, it is too soon to characterize it as slow to get the virus.

COMPARISON WITH PARENTAL AND COMMERCIAL CULTIVARS

'Echo' differs from the female parent 'Perpetua' (U.S. Plant Pat. No. 24,209), which is also the closest comparable commercial cultivar, in that its fruit appear longer as they have a greater length to width ratio, the fruit are a much darker violet-blue and the plants are broader and more spreading than upright with grey-red vs olive-green bark and the leaves are narrower and smaller.

'Echo' differs from the male parent ORUS 55-1 (unpatented) in that the fruit are much more blue than black and the plant is more compact with glossy leaves.

I claim:

1. A new and distinct cultivar of blueberry plant, substantially as illustrated and described, characterized by its ability to flower on new growth without any chilling; attractive combination of flowers and fruit in late summer and fall; dark green, glossy, attractive leaves; edible, dark blue fruit; and vigorous and compact, spreading, plant habit.

* * * * *



Fig. 1



Fig. 2



Fig. 3