

US00PP29827P2

# (12) United States Plant Patent Burton

# (10) Patent No.: US PP29,827 P2 (45) Date of Patent: Nov. 13, 2018

#### (54) COPROSMA PLANT NAMED 'NGCOP8'

(50) Latin Name: *Coprosma repens*Varietal Denomination: **NGCOP8** 

(71) Applicant: Stephen Burton, Cambridge (NZ)

(72) Inventor: Stephen Burton, Cambridge (NZ)

(\*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: 15/732,626

(22) Filed: Dec. 5, 2017

(51) Int. Cl. A01H 5/00

(2018.01)

(52)	U.S. Cl.	
	USPC	Plt./226
	CPC	<b>A01H 5/00</b> (2013.01)
(58)	Field of Classification Search	
, ,	USPC	Plt./226
	CPC	A01H 5/00
	See application file for complete search history.	

Primary Examiner — Kent L Bell

# (57) ABSTRACT

A new cultivar of *Coprosma* plant named 'NGCOP8' that is characterized by a compact habit, glossy leaves, young upper side leaves that are red in color with speckles of yellow-green, mature upper side leaves that are grey-green in color and resistance to leaf scorch.

# 1 Drawing Sheet

1

Botanical classification: *Coprosma repens*. Variety denomination: 'NGCOP8'.

#### BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Coprosma* plant botanically known as *Coprosma repens* and hereinafter referred to by the cultivar name 'NGCOP8'.

The new cultivar was discovered by the inventor in a cultivated area of Cambridge, New Zealand in 2012. <sup>10</sup> 'NGCOP8' was discovered as a naturally occurring branch mutation of *Coprosma* 'Pina Colada' (U.S. Plant Pat. No. 21,032).

Asexual reproduction by tip cuttings of the new cultivar 'NGCOP8' was first performed in 2013 in Cambridge, New Zealand. Since that time, under careful observation, the unique characteristics of the new cultivar have been uniform, stable and reproduced true to type in successive generations of asexual reproduction.

### SUMMARY OF THE INVENTION

The following represent the distinguishing characteristics of the new *Coprosma* cultivar 'NGCOP8'.

- 1. Coprosma 'NGCOP8' exhibits a compact habit.
- 2. Coprosma 'NGCOP8' exhibits glossy leaves.
- 3. Coprosma 'NGCOP8' exhibits young upper side leaves that are red in color with speckles of yellow-green.
- 4. Coprosma 'NGCOP8' exhibits mature upper side leaves that are grey-green in color.
- 5. Coprosma 'NGCOP8' exhibits resistance to leaf scorch. The closest comparison cultivars are Coprosma 'Tequila Sunrise' (U.S. Plant Pat. No. 18,392) and the parent plant Coprosma 'Pina Colada'. The new cultivar 'NGCOP8' is distinguishable from 'Tequila Sunrise' by the following characteristics:
  - 1. 'NGCOP8' has young upper side leaves that are red in color with speckles of yellow-green. In comparison, the young upper side leaves of 'Tequila Sunrise' have green centers surrounded by yellow areas with orange margins.

2

2. 'NGCOP8' has mature upper side leaves that are grey-green in color. In comparison, the mature upper side leaves of 'Tequila Sunrise' have green centers surrounded by red areas with red margins.

The new cultivar 'NGCOP8' is distinguishable from the parent plant 'Pina Colada' by the following characteristics:

- 1. 'NGCOP8' has young upper side leaves that are red in color with speckles of yellow-green. In comparison, the young upper side leaves of 'Pina Colada' are orange-yellow.
- 2. 'NGCOP8' exhibits resistance to leaf scorch. In comparison, the leaves of 'Pina Colada' are more susceptible to leaf scorch.

# BRIEF DESCRIPTION OF THE DRAWING

The accompanying photograph illustrates the distinguishing traits of *Coprosma* 'NGCOP8'. The plant in the photograph shows an overall view of a 12 month old plant.

The photograph was taken using conventional techniques and although colors may appear different from actual colors due to light reflectance it is as accurate as possible by conventional photographic techniques.

#### BOTANICAL DESCRIPTION OF THE PLANT

The following is a detailed description of the new *Coprosma* cultivar named 'NGCOP8'. Data was collected in Grand Haven, Mich. from 12 month old greenhouse grown plants in one gallon containers. The plants were grown under natural light. The time of year was Summer and the average temperature range was 18 to 27 degrees Centigrade during the day and 5 to 10 degrees Centigrade at night. No photoperiodic treatments or growth retardants were used. Color determinations are in accordance with The Royal Horticultural Society Colour Chart 2015 edition, except where general color terms of ordinary dictionary significance are used. The growing requirements are similar to the species. 'NGCOP8' has not been tested under all possible conditions and phenotypic differences may be observed with

**4** 

Foliage: variations in environmental, climatic, and cultural conditions, however, without any variance in genotype. Leaf arrangement.—Opposite. Botanical classification: Coprosma repens 'NGCOP8'. Compound or single.—Single. *Leaf shape.*—Ovate. Use: Ornamental perennial. Parentage: 'NGCOP8' is a naturally occurring branch muta- 5 Leaf apex.—Rounded. tion of *Coprosma* 'Pina Colada'. Leaf base.—Cuneate. Leaf length.—Average 2.0 cm. in length. Vigor: Moderate. Leaf width.—Average 1.5 cm. in width. Growth rate: Moderate. Growth habit: Freely branching, dense, bushy habit. Leaf texture.—Smooth, waxy both surfaces. Plant shape: Rounded, mounding. Leaf luster.—Glossy both surfaces. 10 Suitable container size: One gallon container. Leaf pubescence.—Absent. Height: Average 14 cm. in height. Leaf margin.—Entire, unevenly recurved, slightly Width: Average 24 cm. in width. wavy. Low temperature tolerance: -1 degrees Centigrade. Vein pattern.—Pinnate. High temperature tolerance: 39 degrees Centigrade. Young leaf color (upper surface).—46A, speckled with Propagation: Tip cuttings. 150A. Time to initiate roots: Approximately 30 days in the Summer Young leaf color (lower surface).—164B. at 22 degrees Centigrade and 35 days in the Winter at 22 Mature leaf color (upper surface).—189A. degrees Centigrade. Mature leaf color (lower surface).—199C. Time to produce a rooted cutting: Approximately 112 days 20 Vein color (upper surface).—Same as leaf color. in the Summer at 25 degrees Centigrade and 140 days in Vein color (lower surface).—Same as leaf color. Durability of foliage to stress.—Moderate to high. the Winter at 25 degrees Centigrade. Crop time: Approximately 12 months. Leaf attachment.—Petiolate. Root system: Fibrous, densely matted. Petiole: Root color: 165D. Petiole dimensions.—5.0 mm. in length and 1.5 mm. in Lateral branches: diameter. Branching habit.—Freely branching. *Petiole texture.*—Smooth. Basal branching.—Yes. Petiole pubescence.—Absent. Quantity.—Approximately 6. Petiole color.—179B. Dimensions.—12.0 cm. in length and 3.0 mm. in diam- 30 Flowers: Coprosma 'NGCOP8' has not produced flowers to date. eter. Internode length.—10.0 mm. Disease and insect resistance: Plants of the new *Coprosma* have not been observed for disease or insect resistance. Stem color.—143D with tinges of 39A. Stem shape.—Round in cross-section. The invention claimed is: Stem pubescence.—Absent. 1. A new and distinct variety of Coprosma plant named Stem aspect.—45 degrees from vertical. 'NGCOP8' as described and illustrated. Stem strength.—Strong.

Pinching.—Yes.

