



US00PP30133P2

(12) **United States Plant Patent**  
**Hanes**

(10) **Patent No.:** **US PP30,133 P2**  
(45) **Date of Patent:** **Jan. 22, 2019**

(54) **PELARGONIUM PLANT NAMED ‘PEQZ0030’**

(50) Latin Name: *Pelargonium interspecific*  
Varietal Denomination: **PEQZ0030**

(71) Applicant: **SYNGENTA PARTICIPATIONS AG,**  
Basel (CH)

(72) Inventor: **Mitchell E. Hanes,** Gilroy, CA (US)

(73) Assignee: **Syngenta Participations AG,** Basel  
(CH)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 13 days.

(21) Appl. No.: **15/731,600**

(22) Filed: **Jul. 6, 2017**

(51) **Int. Cl.**  
*A01H 5/02* (2018.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./324**  
CPC ..... *A01H 5/02* (2013.01)

(58) **Field of Classification Search**  
USPC ..... **Plt./324**  
See application file for complete search history.

*Primary Examiner* — Anne Marie Grunberg

(74) *Attorney, Agent, or Firm* — Dale Skalla

(57) **ABSTRACT**

A new *Pelargonium* plant named ‘PEQZ0030’ particularly distinguished by the dark rose colored flower center surrounded by a deep rose petal edge. The plant is compact in the greenhouse with good heat and drought tolerance with continuous color through the summer.

**1 Drawing Sheet**

**1**

Latin name of the genus and species of the plant claimed:  
*Pelargonium interspecific*.

Varietal denomination: ‘PEQZ0030’.

**BACKGROUND OF THE NEW PLANT**

The present invention comprises a new *Pelargonium*, botanically known as *Pelargonium interspecific*, and hereinafter referred to by the variety name ‘PEQZ0030’.

‘PEQZ0030’ is a product of a planned breeding program. The new cultivar ‘PEQZ0030’ has rose bicolored inflorescences held above the medium-green foliage, very heat and drought tolerant with continuous color through the most extreme summer heat, and exceptional edema tolerance on a well-branched plant habit.

‘PEQZ0030’ originates from a hybridization in a controlled breeding program made in September 2010 in a greenhouse in Guatemala. The female parent was an unpatented, proprietary plant of *P. interspecific* parentage, identified as ‘10816-3’ with rose color florets and a splash or spot in the center of the floret rose. When compared to ‘PEQZ0030’ the female has a less pronounced splash.

The male parent of ‘PEQZ0030’ was an unpatented, proprietary plant of *P. interspecific* parentage, identified as ‘10997-1’ with a rose base color and a very large splash pattern. The male has a more open plant habit when compared to ‘PEQZ0030’. The resultant seed was sown in June 2011.

‘PEQZ0030’ was selected as one flowering plant within the progeny of the stated cross in October 2011 in a greenhouse in Gilroy, Calif.

The first act of asexual reproduction of ‘PEQZ0030’ was accomplished when vegetative cuttings were propagated from the initial selection in the December 2011 in a greenhouse in Gilroy, Calif.

**BRIEF SUMMARY OF INVENTION**

Horticultural examination of plants grown from cuttings of the plant initiated in January 2012 in Gilroy, Calif., and

**2**

continuing thereafter, has demonstrated that the combination of characteristics as herein disclosed for ‘PEQZ0030’ are firmly fixed and are retained through successive generations of asexual reproduction.

5 ‘PEQZ0030’ has not been observed under all possible environmental conditions. The phenotype may vary significantly with variations in environment such as temperature, light intensity, and day length.

10 A Plant Breeder’s Right for this cultivar has not been applied for. ‘PEQZ0030’ has not been made publicly available prior to the effective filing date of this application, notwithstanding any disclosure that may have been made less than one year prior to the effective filing date of this application by the inventor or another who obtained  
15 ‘PEQZ0030’ directly from the inventor.

The following traits have been repeatedly observed and are determined to be the basic characteristics of the new variety. The combination of these characteristics distinguishes this *Pelargonium* as a new and distinct variety.  
20

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

25 The accompanying photographic drawing shows typical flower and foliage characteristics of ‘PEQZ0030’ with colors being as true as possible with an illustration of this type. The photographic drawing shows in FIG. 1, a flowering plant of the new variety and  
30 in FIG. 2, a close-up of an inflorescence.

**DETAILED BOTANICAL DESCRIPTION**

35 The plant descriptions and measurements were taken in Gilroy, Calif. in late May 2017 under natural light. These plants were approximately 12-14 weeks old and were grown in a 4.5 inch pots, in a greenhouse trial. The plants shown in the photographs were about 14 weeks old. The photographs were taken in June 2017. Color references are made to The Royal Horticultural Society Colour Chart (R.H.S.) 2001.

TABLE 1

DIFFERENCES BETWEEN THE NEW VARIETY 'PEQZ0030' AND A MOST SIMILAR VARIETY		
	'PEQZ0030'	'PEQZ0006' (U.S. Plant Pat. No. 25,175)
Floret size:	Same	Same
Floret petal color:	Rose base color smaller splash	Red base color larger splash
Umbel diameter size:	Smaller	Larger
Peduncle length:	Shorter	Longer

## Plant:

*Form, growth and habit.*—Upright, outwardly spreading and rounded growth habit, heat and drought tolerant with continuous color through the most extreme summer heat, edema tolerance, well-branched plant habit.

*Plant height.*—14-19 cm.

*Plant height (inflorescence included).*—16-24 cm.

*Plant width.*—25-28 cm.

## Roots:

*Number of days to initiate roots.*—15-18 days at about 22 degrees C.

*Number of days to produce a rooted cutting.*—21-23 days at 22 degrees C.

*Type.*—Fine, fibrous, free branching.

*Color.*—RHS N155B but whiter.

## Foliage:

*Immature leaf, color upper surface.*—Between RHS 144A and RHS 144B.

*Immature leaf, lower surface.*—Between RHS 144A and RHS 144B.

*Mature leaf, color upper surface.*—Closest to RHS 137A with slightly lighter mottling.

*Mature leaf, color lower surface.*—Closest to RHS 137D.

*Length.*—9-11 cm.

*Width.*—10.5-12.5 cm.

*Shape.*—Cordate.

*Base shape.*—Cordate.

*Apex shape.*—Acute.

*Margin.*—Slightly dentate.

*Texture upper side.*—Hirsute.

*Texture lower side.*—Hirsute.

*Leaf zonation color.*—None.

*Color of veins, upper surface.*—RHS 137C becoming indistinct.

*Color of veins, lower surface.*—RHS 137C becoming indistinct.

*Pattern of veining.*—Palmate.

*Petiole color.*—RHS 137C.

*Petiole length.*—6-9.0 cm.

*Diameter of petiole.*—0.2 cm.

*Texture.*—Pilose, hirsute, glandular hairs.

## Stem:

*Quantity of branches.*—3.

*Color of stem.*—Between RHS 144A and RHS 144B.

*Length of stem.*—6-10 cm.

*Diameter.*—0.5-0.7 cm.

*Length of internodes.*—2.0-2.5 cm.

*Texture.*—Sparsely hirsute, pilose, glandular hairs.

## Peduncle:

*Color of peduncle.*—RHS 137D.

*Length of peduncle.*—10-14 cm.

*Peduncle diameter.*—0.3-0.35 cm.

*Texture.*—Hirsute, glandular hairs.

## Pedicel:

*Color of pedicel.*—RHS 137D turning to RHS 187D.

*Length of pedicel.*—3.5-4.5 cm.

*Diameter of pedicel.*—0.15-0.2 cm.

*Texture.*—Sparsely pilose, glandular hairs.

## Bud (just before opening):

*Color.*—Between RHS 57C and 57D.

*Length.*—1.5-2.0 cm.

*Width.*—0.7-0.8 cm.

*Shape.*—Elliptical.

## Inflorescence:

*Type.*—Umbel; semi-spherical or nearly semi-spherical.

*Lastingness of individual flowers.*—7-9 days at 18° C. temperature.

*Number of inflorescences per plant.*—2, with 2 immature umbels in various stages.

*Fragrance.*—None.

*Umbel diameter.*—9-12 cm.

*Umbel depth.*—6.5-7.5 cm.

## Corolla:

*Form.*—Semi-double.

*Number of petals.*—7.

*Diameter of flower.*—About 5.0-5.5 cm.

*Depth of flower.*—2-2.5 cm.

*Color upper petals, upper surface.*—Closest to RHS 67C with RHS 45B splash and veining.

*Color upper petals, lower surface.*—Closest to RHS 67C with RHS 45B spots.

*Length of upper petals.*—3.0 cm.

*Width of upper petals.*—2.3 cm.

*Color lower petals, upper surface.*—Closest to RHS 67C with a spot or RHS 45B.

*Color lower petals, lower surface.*—Closest to RHS 67C.

*Length of lower petals.*—2.5-2.7 cm.

*Width of lower petals.*—2.0-2.5 cm.

*Petal shape.*—Obovate to spatulate.

*Apex shape.*—Rounded.

*Margin.*—Entire.

*Base.*—Attenuate.

*Petal texture.*—Papillose on both surfaces.

## Calyx:

*Number of sepals.*—5.

*Color of sepals.*—RHS 144A and RHS 144B.

*Length of sepals.*—1.2-1.4 cm.

*Width of sepals.*—0.3-0.4 cm.

*Sepal shape.*—Lanceolate to linear.

*Apex shape.*—Acute.

*Margins.*—Mostly fused.

*Texture, upper surface.*—Glabrous.

*Lower surface.*—Glandular hairs, hirsute.

## Reproductive organs:

## Gynoecium:

*Pistil.*—1.

*Length.*—0.8 cm.

*Style color.*—Closest to RHS 53C.

*Style length.*—0.4 cm.

*Stigma color.*—Closest to RHS 53C.

*Ovary color.*—Between RHS 144A and RHS 144B.

*Ovary length.*—0.3-0.4 cm.

*Ovary diameter.*—0.2 cm.

Androecium:

*Number of stamens.*—9.

*Color of filaments.*—RHS 67C with RHS 155C basally.

*Length filaments.*—0.8-1.0 cm.

*Anther color.*—Between RHS 25B and RHS 25C with 5  
a light mottling of RHS N57B.

*Length of anthers.*—0.2 cm.

*Color of pollen.*—RHS 25B.

*Pollen amount.*—Normal.

*Fertility/seed set.*—Has not been determined to date.

*Disease/pest resistance.*—Has not been determined to date.

What is claimed is:

1. A new and distinct variety of *Pelargonium* plant named 'PEQZ0030' substantially as illustrated and described herein.

\* \* \* \* \*



FIG. 1



FIG. 2