



US00PP30584P2

(12) **United States Plant Patent**
Clark

(10) **Patent No.:** **US PP30,584 P2**
(45) **Date of Patent:** **Jun. 18, 2019**

(54) **COLEUS PLANT NAMED ‘UF16-64-1’**

(50) Latin Name: *Plectranthus scutellarioides*
Varietal Denomination: **UF16-64-1**

(71) Applicant: **Florida Foundation Seed Producers, Inc.**, Marianna, FL (US)

(72) Inventor: **David G. Clark**, Gainesville, FL (US)

(73) Assignee: **Florida Foundation Seed Producers, Inc.**, Marianna, FL (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/732,776**

(22) Filed: **Dec. 27, 2017**

(51) **Int. Cl.**
A01H 5/12 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./469**
CPC *A01H 5/12* (2013.01)

(58) **Field of Classification Search**
USPC **Plt./469**
See application file for complete search history.

Primary Examiner — Anne Marie Grunberg
(74) *Attorney, Agent, or Firm* — Dentons US LLP

(57) **ABSTRACT**

‘UF16-64-1’ is a new coleus plant with novel characteristics that include a vigorous compact mounded growth habit, an excellent heat tolerance, and consistent brightly colored leaves. It has superior stability in foliage color in both sun and shade conditions, maintaining stable color in all conditions. ‘UF16-64-1’ has not been observed to produce flowers in any trial we have conducted to date, and thus it has long season performance as an annual plant in the landscape until late Fall.

3 Drawing Sheets

1

Latin name of the genus and species of the plant claimed: *Plectranthus scutellarioides*.
Cultivar denomination: ‘UF16-64-1’.

BACKGROUND OF THE INVENTION

The invention relates to a new and distinct cultivar of coleus plant named ‘UF16-64-1’. ‘UF16-64-1’ originated from an open-pollination conducted in May-November 2015 in Gainesville, Fla. between the female coleus plant ‘UF15-56-1’ (unpatented) and an unknown male coleus plant. A single seedling was chosen in May 2016 for further asexual propagation in Gainesville, Fla.

‘UF16-64-1’ has been reproduced asexually for over 18 months through vegetative cuttings and has been found to retain its distinctive characteristics through successive asexual propagations. ‘UF16-64-1’ was first propagated asexually by meristem tip cuttings in May 2016 in Gainesville, Fla., and has remained true-to-type since that time.

‘UF16-64-1’ has large green leaves with distinct hot-pink centers and maroon venation that are much longer than they are wide; whereas, ‘UF15-56-1’, the female parent, has much smaller leaves that are dark green with maroon centers and no pink coloration. ‘UF16-64-1’ has an extremely compact well-branched mounded habit; whereas, ‘UF15-56-1’ is more vigorous and upright in habit with less lateral branching.

‘UF16-64-1’ and commercial cultivar ‘Mainstreet Fifth Avenue’ (‘UF12-86-9’, unpatented) both have tri-colored green, maroon, and pink leaves and purple stems; however, ‘UF16-64-1’ has a more vigorous growth habit and its leaves are larger and more brightly colored in both sun and shade conditions. ‘UF16-64-1’ has not been observed to flower; whereas, ‘Mainstreet Fifth Avenue’ flowers profusely.

SUMMARY OF THE INVENTION

The following are characteristics of ‘UF16-64-1’ when grown under normal horticultural practices in Gainesville,

2

Fla. ‘UF16-64-1’ has a novel, compact upright growth habit, an excellent heat tolerance, and consistent bright colored multi-colored leaves that are significantly different than other coleus plants. It has superior stability in foliage color in both sun and shade conditions, maintaining stable color in all conditions. When ‘UF16-64-1’ is grown as a stock plant, it has a vigorous but compact upright growth habit with excellent lateral branching, and thus provides ample vegetative propagules for producers. ‘UF16-64-1’ has not been observed to produce flowers in any trial we have conducted to date in Gainesville, Fla., and thus it is desirable for long-season performance in the landscape.

BRIEF DESCRIPTION OF THE DRAWINGS

This new coleus plant is illustrated by the accompanying photographs, which show the plant’s form and foliage. The colors shown are as true as can be reasonably obtained by conventional photographic procedures. The photographs in FIGS. 2 and 3 are of plants grown for 11 weeks from unrooted cuttings in May-July 2017 in a poly-covered plastic greenhouse in Gainesville, Fla.

FIG. 1—Shows the pedigree of ‘UF16-64-1’.

FIG. 2—Shows the growth habit, form, and foliage of ‘UF16-64-1’.

FIG. 3—Shows a close-up of the foliage of ‘UF16-64-1’.

DETAILED BOTANICAL DESCRIPTION

The following detailed description sets forth the distinctive characteristics of ‘UF16-64-1’. The detailed description was obtained using 10-week-old plants grown from unrooted cuttings in May-July 2017 in a poly-covered plastic greenhouse in Gainesville, Fla. The plants were propagated in mist for 10 days after cuttings were stuck and then grown in 1-gallon pots for approximately 8.5 additional

weeks. Color references are to The R.H.S. Colour Chart of The Royal Horticultural Society of London (R.H.S.), 2007 5th Edition.

Classification:

Family.—Lamiaceae.

Botanical.—*Plectranthus scutellarioides*.

Common name.—Coleus.

Cultivar name.—‘UF16-64-1’.

Plant description:

Form.—Spreading.

Habit.—Upright.

Height (from top of soil).—25-30 cm.

Width (horizontal plant diameter).—45-50 cm.

Propagation:

Type cuttings.—Vegetative meristems having at least 1 node.

Time to initiate roots.—3-4 days.

Time to produce a rooted cutting.—7-10 days.

Root habit.—Fibrous.

Root description.—Callus forms in 2 to 3 days, roots initiate in 3-4 days and become a highly branched cutting in 7-10 days.

Branches:

Quantity per plant.—6-7.

Branch color.—RHS N77B.

Texture.—Smooth.

Pubescence.—Not present.

Stem description.—Square-shaped stem, 0.8 cm in diameter at the soil line.

Branch diameter.—0.4-0.5 cm at the base of a 24 cm long branch.

Branch length.—22-25 cm.

Internode length.—4-5 cm.

Anthocyanin.—Not present.

Leaves:

Quantity of leaves per branch.—14-16. Arrangement: Opposite.

Fragrance.—Not fragrant.

Shape.—Ovate.

Length.—11-13 cm.

Width.—7-8 cm.

Apex.—Broadly acute.

Base.—Rounded.

Margin.—Crenate.

Leaf texture (both surfaces).—Smooth.

Pubescence color (both surfaces).—Not present.

Venation color.—Upper surface: Center=RHS 64A; Mid-Center=RHS N79A; Margin=RHS 138A. Lower surface: RHS N77B.

Venation pattern.—Upper surface: Arcuate. Lower surface: Reticulate.

Color.—Immature leaf: Upper surface: Center=RHS 72A; Margin=RHS 146B. Lower surface: RHS N186C.

Color.—Mature leaf: Upper surface: Center=RHS 64A; Mid-Center=RHS N79A; Margin=RHS 138A. Lower surface: Center=RHS 64B; Mid-Center to Margin=RHS N79A.

Petiole length.—3-4 cm.

Petiole diameter.—0.2-0.3 cm.

Petiole color.—RHS 64A.

Petiole texture.—Smooth, no pubescence.

Flowers and seeds: Flowers and seeds have not been observed to date during formal trials in Gainesville, Fla.

Fruit/seed set: Fruit/seed not observed.

Disease and insect resistance: Disease and insect resistance is typical of the species, and thus no claims are made of any superior disease or insect resistance with this cultivar. The most common insect pests observed on this plant in Gainesville, Fla. have been long-tailed or citrus mealybugs (*Pseudococcus* sp.), which occur on older stock plant material held in the greenhouse for over 3-4 months. Impatiens Necrotic Spot Virus (*Bunyaviridae*) has also been observed in plants confined in greenhouses with mixed crops (peppers) infected with Western flower thrips (*Frankliniella occidentalis*). The most common pathogen of this species in the U.S. is downy mildew (*Peronospora lamii*). This pathogen has been observed in stock materials grown closely together in cooler growing seasons.

What is claimed is:

1. A new and distinct *Plectranthus scutellarioides* plant called ‘UF16-64-1’ as shown and described herein.

* * * * *



FIG. 1




FIG. 2



FIG. 3