



US00PP30854P3

(12) **United States Plant Patent**
Wain

(10) **Patent No.:** **US PP30,854 P3**
(45) **Date of Patent:** **Aug. 27, 2019**

- (54) **CHRYSANTHEMUM PLANT NAMED ‘FICHRYSUNAU’**
- (50) Latin Name: *Chrysanthemum X morifolium*
Varietal Denomination: **Fichrysunaub**
- (71) Applicant: **DUMMEN GROUP B.V.**, De Lier (NL)
- (72) Inventor: **Peter Wain**, Locks Heath (GB)
- (73) Assignee: **Dümmen Group B.V.**, De Lier (NL)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **15/732,517**
- (22) Filed: **Nov. 21, 2017**
- (65) **Prior Publication Data**
US 2018/0146592 P1 May 24, 2018
- Related U.S. Application Data**
- (60) Provisional application No. 62/497,574, filed on Nov. 22, 2016.
- (51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/14 (2018.01)
- (52) **U.S. Cl.**
USPC **Plt./287**
CPC *A01H 6/1424* (2018.05); *A01H 5/02* (2013.01)

(58) **Field of Classification Search**
USPC Plt./263.1, 284, 286, 287
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

Germania Seed Company plant listing (https://www.germaniaseed.com/pdf/2016/761_mast_mums.pdf). 2016. Citation for ‘Sunbeam Aubergine’. 5 pages. (Year: 2016).*
Dummen Orange 2018 collection (https://www.dummenorange.com/flippingbooks/DummenOrange-Garden-Mums-2018/files/assets/common/downloads/Catalog_Garden%20mums%202018_FLIPPING%20BOOK.pdf). Citation for ‘Sunbeam Aubergine’. 6 pages.*

* cited by examiner

Primary Examiner — Susan McCormick Ewoldt

Assistant Examiner — Karen M Redden

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named ‘Fichrysunaub’, characterized by its upright to outwardly spreading and uniformly and broadly mounded plant habit; moderately vigorous to vigorous growth habit; freely branching habit; dense and full plant form; uniform and freely flowering habit; decorative-type inflorescences with deep red purple-colored ray florets; early to mid-season flowering habit, grown under natural season conditions, plants begin flowering in mid to late August in the United Kingdom; and good garden performance.

1 Drawing Sheet

Botanical designation: *Chrysanthemum X morifolium*.
Cultivar denomination: ‘FICHRYSUNAU’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct *Chrysanthemum* plant, botanically known as *Chrysanthemum X morifolium*, commercially grown as a garden *Chrysanthemum* plant, referred to as code number 65799 in U.S. Provisional Patent Application Ser. No. 62/497,574 and hereinafter referred to by the name ‘Fichrysunaub’.

The new *Chrysanthemum* plant is a product of a planned breeding program conducted by the Inventor in Fareham, Hampshire, United Kingdom. The objective of the breeding program is to create new early to mid-season flowering garden *Chrysanthemum* plants with numerous attractive inflorescences.

The new *Chrysanthemum* plant originated from a cross-pollination made in January, 2012 by the Inventor in Fareham, Hampshire, United Kingdom of a proprietary selection of *Chrysanthemum X morifolium* identified as code number 83453, not patented, as the female, or seed, parent with a proprietary selection of *Chrysanthemum X morifolium* identified as code number 802403, not patented, as the male, or

pollen, parent. The new *Chrysanthemum* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Fareham, Hampshire, United Kingdom in September, 2012.

Asexual reproduction of the new *Chrysanthemum* plant by terminal vegetative cuttings was first conducted in Fareham, Hampshire, United Kingdom in December, 2012. Asexual reproduction by terminal vegetative cuttings has shown that the unique features of this new *Chrysanthemum* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Chrysanthemum* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature, daylength and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Fichry-

sunaub'. These characteristics in combination distinguish 'Fichrysunaub' as a new and distinct *Chrysanthemum* plant:

1. Upright to outwardly spreading and uniformly and broadly mounded plant habit.
2. Moderately vigorous to vigorous growth habit.
3. Freely branching habit; dense and full plant form.
4. Uniform and freely flowering habit.
5. Decorative-type inflorescences with deep red purple-colored ray florets.
6. Early to mid-season flowering habit, grown under natural season conditions, plants begin flowering in mid to late August in the United Kingdom.
7. Good garden performance.

Plants of the new *Chrysanthemum* can be compared to plants of the female parent selection. Plants of the new *Chrysanthemum* differ primarily from plants of the female parent selection in time to flower as plants of the new *Chrysanthemum* flower earlier than plants of the female parent selection.

Plants of the new *Chrysanthemum* can be compared to plants of the male parent selection. Plants of the new *Chrysanthemum* differ primarily from plants of the male parent selection in the following characteristics:

1. Plants of the new *Chrysanthemum* flower later than plants of the male parent selection.
2. Plants of the new *Chrysanthemum* and the male parent selection differ in ray floret color as plants of the male parent selection have inflorescences with dark pink-colored ray florets.

Plants of the new *Chrysanthemum* can be compared to plants of *Chrysanthemum X morifolium* 'Fichryevenpur', disclosed in U.S. Plant patent application Ser. No. 15/732, 504. In side-by-side comparisons, plants of the new *Chrysanthemum* differ primarily from plants of 'Fichryevenpur' in time to flower as plants of the new *Chrysanthemum* flower earlier than plants of 'Fichryevenpur'. In addition, plants of the new *Chrysanthemum* have larger inflorescences than plants of 'Fichryevenpur'.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying photograph illustrates the overall appearance of the new *Chrysanthemum* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Chrysanthemum* plant.

The photograph is a top perspective view of a typical flowering plant of 'Fichrysunaub' grown in a 14-cm container.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations and measurements describe plants grown during the winter in 14-cm containers in a glass-covered greenhouse in Fareham, Hampshire, United Kingdom and under cultural practices typical of commercial garden *Chrysanthemum* production. During the production of the plants, day and night temperatures ranged from 17° C. to 21° C. and light levels averaged 6,000 lux. Plants were grown under long day/short night conditions for about five weeks (including propagation period) and then grown under short day/long night conditions to induce inflorescence initiation and development. Plants were twelve weeks old when the photograph

and detailed description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, Fifth Edition, except where general terms of ordinary dictionary significance are used.

- 5 Botanical classification: *Chrysanthemum X morifolium* 'Fichrysunaub'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Chrysanthemum X morifolium* identified as code number 83453, not patented.

Male, or pollen, parent.—Proprietary selection of *Chrysanthemum X morifolium* identified as code number 802403, not patented.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate roots, summer.—About ten days at temperatures about 21° C.

Time to initiate roots, winter.—About twelve days at temperatures about 21° C.

Time to produce a rooted young plant, summer.—About three weeks at temperatures about 21° C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures about 21° C.

Root description.—Medium in thickness, fibrous; typically light brown in color, actual color of the roots is dependent on substrate composition, water quality, fertilizer type and formulation, substrate temperature and physiological age of roots.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Herbaceous decorative-type garden *Chrysanthemum*; stems upright to outwardly spreading giving a uniformly and broadly mounded appearance to the plant; numerous lateral branches and relatively short internodes, dense and full plant form; moderately vigorous to vigorous growth habit.

Plant height.—About 16 cm.

Plant width.—About 27 cm.

Branching habit.—Freely branching habit; about ten lateral branches develop after removal of terminal apex (pinching).

Lateral branches.—Length: About 9.5 cm. Diameter: About 3 mm. Internode length: About 1.9 cm. Strength: Strong. Aspect: About 50° from vertical and then bending upwardly. Texture: Fine pubescence. Color: Close to 138A.

Leaf description:

Arrangement.—Alternate, simple.

Length.—About 3.2 cm.

Width.—About 2.2 cm.

Shape.—Palmately-lobed; roughly ovate with three to five lobes.

Apex.—Mucronate.

Base.—Attenuate to truncate.

Margin.—Slightly dentate and palmately lobed; sinuses between lateral lobes mostly divergent.

Texture, upper and lower surfaces.—Fine pubescence; slightly rough; veins prominent on lower surface.

Color.—Developing leaves, upper surface: Close to 137A. Developing leaves, lower surface: Close to 137B. Fully expanded leaves, upper surface: Close to N137C; venation, close to 138B. Fully expanded leaves, lower surface: Close to 137C; venation, close to 138B.

Petioles.—Length: About 1.5 cm. Diameter: About 2 mm. Texture, upper and lower surfaces: Fine pubescence; slightly rough. Color, upper and lower surfaces: Close to 138B.

Inflorescence description:

Form and flowering habit.—Decorative-type inflorescence form with ligulate-shaped ray florets; inflorescences borne on terminals above and beyond the foliar plane; disc and ray florets arranged acropetally on a capitulum; freely flowering habit with about 70 inflorescences developing per plant during the flowering season.

Fragrance.—Mildly fragrant; pungent, herbaceous.

Flowering response.—Early to mid-season flowering habit, plants exposed to natural season conditions begin flowering in mid to late August in the United Kingdom; plants flower uniformly and continuously during the flowering season.

Inflorescence longevity.—Inflorescences maintain good color and substance for about three to five weeks on the plant; inflorescences persistent.

Inflorescence buds.—Height: About 4 mm. Diameter: About 5 mm. Shape: Oblate. Color: Close to 138A.

Inflorescence diameter.—About 4 cm.

Inflorescence height.—About 1.5 cm.

Receptacles.—Height: About 3 mm. Diameter: About 3 mm. Shape: Conical. Color: Close to 145C.

Ray florets.—Number of ray florets per inflorescence: About 147 arranged in about seven whorls. Orientation: Initially upright, then about 40° from vertical. Length: About 1.9 cm. Width: About 5 mm. Shape: Ligulate. Apex: Emarginate. Base: Fused into a short tube. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; double-keeled. Color: When opening, upper surface: Close to 59A. When opening, lower surface: Close to 61A. Fully opened, upper surface: Close to N59B; with development,

color becoming closer to 71B. Fully opened, lower surface: Close to 60C; with development, color becoming closer to 79D.

Disc florets.—Disc floret initiation and development has not been observed on plants of the new *Chrysanthemum* to date.

Phyllaries.—Number of phyllaries per inflorescence: About 18 arranged in about three whorls. Length: About 6 mm. Width: About 2 mm. Shape: Lanceolate. Apex: Acute. Base: Obtuse. Margin: Entire. Texture, upper surface: Smooth, glabrous; waxy. Texture, lower surface: Fine pubescence; waxy. Color, upper surface: Close to 138B. Color, lower surface: Close to 138A.

Peduncles.—Length, terminal peduncle: About 2.1 cm. Diameter, terminal peduncle: About 2 mm. Angle: Erect to about 10° from vertical. Strength: Moderately strong. Texture: Densely pubescent. Color: Close to 138A.

Reproductive organs.—Androecium: None observed. Gynoecium: Present only on ray florets. Pistil length: About 5 mm. Stigma shape: Bi-parted. Stigma color: Close to 6B. Style length: About 3 mm. Style color: Close to 1C. Ovary color: Close to NN155C.

Seeds and fruits.—Seed and fruit production has not been observed on plants of the new *Chrysanthemum* to date.

Disease & pest resistance: Resistance to pathogens and pests common to *Chrysanthemum* plants has not been observed on plants of the new *Chrysanthemum* to date.

Garden performance: Plants of the new *Chrysanthemum* have demonstrated good garden performance and to tolerate temperatures from about 0° C. to about 35° C.

It is claimed:

1. A new and distinct *Chrysanthemum* plant named 'Fichrysunaub' as illustrated and described.

* * * * *

