

US00PP30879P2

(12) **United States Plant Patent**
Finn

(10) **Patent No.:** **US PP30,879 P2**
(45) **Date of Patent:** **Sep. 10, 2019**

(54) **BLACKBERRY PLANT NAMED ‘TWILIGHT’**

(50) Latin Name: *Rubus* subg. *Rubus* Watson
Varietal Denomination: **Twilight**

(71) Applicant: **The United States of America, as Represented by the Secretary of Agriculture**, Washington, DC (US)

(72) Inventor: **Chad E. Finn**, Corvallis, OR (US)

(73) Assignee: **The United States of America, as Represented by the Secretary of Agriculture**, Washington, DC (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/998,301**

(22) Filed: **Aug. 2, 2018**

(51) **Int. Cl.**
A01H 5/08 (2018.01)
A01H 6/74 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./203**

(58) **Field of Classification Search**
USPC Plt./156, 203
See application file for complete search history.

Primary Examiner — Susan McCormick Ewoldt

Assistant Examiner — Karen M Redden

(74) *Attorney, Agent, or Firm* — John Fado; Ariel Atkinson

(57) **ABSTRACT**

A new and distinct blackberry cultivar that originated from seed produced from a cross between the thornless female blackberry plant ‘Ouachita’ (U.S. Plant Pat. No. 17,762) and the thornless, male parent blackberry plant ‘ORUS 2867-4’ (unpatented). This new blackberry cultivar can be distinguished by being fairly early ripening for a semi-erect blackberry, by its large sized, very firm berries with tough skin and a crisp texture when eaten and excellent flavor, by its high yields borne on a vigorous plant with a semi-erect type growth habit and by its completely thornless canes.

4 Drawing Sheets

1

Latin name of the genus and species of the plant claimed: ‘Twilight’ is a blackberry plant that is *Rubus* subg. *Rubus* Watson.

Variety denomination: The new blackberry plant claimed is of the variety denominated ‘Twilight’ *Rubus* subg. *Rubus* Watson.

BACKGROUND OF THE NEW PLANT

The present invention relates to a new and distinct blackberry cultivar designated ‘Twilight’ and botanically known as *Rubus* subg. *Rubus* Watson. This new blackberry cultivar was discovered in Corvallis, Oreg. in August 2012 and originated from a cross between the thornless female blackberry plant ‘Ouachita’ (U.S. Plant Pat. No. 17,762) and the thornless, male parent blackberry plant ‘ORUS 2867-4’ (unpatented). ‘ORUS 2867-4’ (unpatented) was a selection from a cross of the thornless semi-erect blackberry ‘Triple Crown’ (unpatented) and a thorny trailing blackberry selection ‘ORUS 1393-1’ (unpatented). ‘Twilight’s spinelessness was originally derived from ‘Merton Thornless’ (U.S. Plant Pat. No. 571). The original seedling of the new cultivar was asexually propagated at a nursery in Benton County, Oreg. The new cultivar was established in vitro from a cane cutting and microcuttings have been taken and rooted from this sort of culture. The present invention has been found to be stable and reproduce true to type through successive asexual propagations.

SUMMARY OF THE NEW PLANT

The following are the most outstanding and distinguishing characteristics of this new cultivar when grown under normal horticultural practices in Oregon. First, the new

2

cultivar has a high plant vigor as compared to semi-erect blackberry ‘Loch Ness’ (U.S. Plant Pat. No. 6,782); second, the new cultivar has a semi-erect growth habit with thornless canes; third, the new cultivar has an early date for 50% of ripe fruit compared to ‘Chester Thornless’ (unpatented) and ‘Triple Crown’ (unpatented) but not as early as ‘Eclipse’ (U.S. patent application Ser. No. 15/731,503) or ‘Galaxy’ (U.S. Plant Pat. No. 30,062); fourth, the new cultivar has a very firm fruit with abrasion-resistant, tough skin compared to ‘Triple Crown (unpatented) and ‘Chester Thornless’ (unpatented); fifth, the new cultivar has excellent flavor compared to ‘Chester Thornless’ (unpatented) and ‘Loch Ness’ (U.S. Plant Pat. No. 6,782); and sixth, the new cultivar has larger fruit than those of ‘Chester Thornless’ (unpatented) and similar to those of ‘Triple Crown’ (unpatented).

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

This new blackberry plant is illustrated by the accompanying photographs that show the flowers, fruit and entire plants; the colors shown are as true as can be reasonably obtained by conventional photographic procedures.

FIG. 1 shows typical fruiting cluster with ripe fruit ‘Twilight’.

FIG. 2 shows a flat of harvested ‘Twilight’ fruit (left) compared to a flat of ‘Chester Thornless’ (unpatented) fruit (right).

FIG. 3 shows three clamshells of ‘Twilight’ fruit (left column) compared to ‘Chester Thornless’ (unpatented) fruit (right column) after 21 days of refrigerated storage at ~1° C.

FIG. 4 shows entire flowering 3-year old ‘Twilight’ plants. As is typical for commercial production, semi-erect blackberry primocanes are topped at ~1 m and the primocanes and laterals that develop after topping are tied to a two wire

trellis with the lower wire approximately 1.0 m above the ground and the upper wire approximately 1.5 m above the ground.

DETAILED DESCRIPTION OF THE NEW CULTIVAR

The following description of 'Twilight' is based on observations taken from 2012 to 2017 growing seasons in trials in Corvallis and Aurora, Oreg. This description is in accordance with UPOV terminology. Color designations, color descriptions and other phenotypical descriptions may deviate from the stated values and descriptions depending upon variation in environmental, seasonal, climatic and cultural conditions. 'Twilight' has not been observed under all possible environmental conditions. Color terminology follows The Royal Horticultural Society Colour Chart. London (R.H.S.) (5th edition, 2007).

Table 1 shows important plant characteristics of the new cultivar. Characteristics include plant vigor, growth habit, date 50% of fruit were ripe, weight of primary fruit, firmness of fruit flavor of fruit and winter tolerance in Aurora, Oreg. (45° 16' 49" N/122° 44' 50" W) and Lynden, Wash. (48° 56' 48" N/122° 27' 2" W).

TABLE 1

Characteristic	'Twilight'
Plant vigor	High vigor, similar to 'Eclipse', less vigorous than 'Chester Thornless' and 'Triple Crown' and more vigorous than 'Loch Ness'
Growth habit	Semi-erect
Date 50% of fruit were ripe	21 July, early compared to 'Triple Crown' and 'Chester Thornless'; Later than 'Eclipse'
Weight of primary fruit (g)	8.7 g, larger than 'Chester Thornless', similar in size to 'Triple Crown'
Firmness of fruit	Excellent, much firmer than 'Triple Crown' and firmer than 'Chester Thornless'
Skin toughness of fruit	Excellent, much better than 'Triple Crown' and better than 'Chester Thornless'
Flavor of fruit	Very good, excellent compared to 'Chester Thornless', similar to 'Triple Crown' and 'Eclipse'
Winter tolerance in Aurora, Oregon (45 16' 49" N/122 44' 50" W)	Excellent, comparable to 'Triple Crown', 'Chester Thornless', and 'Navaho' (U.S. Plant Pat. No. 6,679)
Winter tolerance in Lynden, Washington (48 56' 48" N/122 27' 2" W)	Excellent, comparable to 'Triple Crown' and 'Chester Thornless'

Table 2 shows floricanes and mature primocane characteristics of the new cultivar. Characteristics include diameter at base, diameter at midpoint, diameter at terminus, internode length at base, internode length at midpoint, internode length at terminus, presence of spines further than 0.6 m from the soil surface, presence of spines less than 0.6 m from the soil surface, floricanes color at base, floricanes color at midpoint, floricanes lateral length, floricanes lateral strength, primocane color at base, primocane color at midpoint, and primocane color at terminus. The characteristics of floricanes color at terminus, floricanes length, and floricanes length (range) are not available for the new cultivar as the standard practice for the cultivar is to remove the tips of primocanes (which become floricanes after receiving winter chilling) to encourage fruiting.

TABLE 2

Characteristic	'Twilight'
Diameter at base (cm)	2.00
Diameter at midpoint (cm)	1.23
Diameter at terminus (cm)	0.37
Internode length at base (cm)	11.93
Internode length at midpoint (cm)	2.87
Internode length at terminus (cm)	2.87
Presence of spines further than 0.6 m from the soil surface	Absent
Presence of spines less than 0.6 m from the soil surface	Absent
Floricanes color at base	146A
Floricanes color at midpoint	144A w/175A tint
Floricanes lateral length	Medium
Floricanes lateral strength	Medium
Primocane color at base	144A
Primocane color at midpoint	144A w/176A streaks
Primocane color at terminus	144A w/185A tint

Table 3 shows floricanes foliage characteristics of the new cultivar. Floricanes characteristics include mature compound leaf width, mature compound leaf length, number of leaflets per floricanes compound leaf, mature leaflet shape, mature leaflet apex, mature leaflet base, mature terminal leaflet width, mature terminal leaflet length, mature first lateral leaflet width, mature first lateral leaflet length, leaflet margin, leaflet serration teeth length, leaflet serration teeth width at base, pubescence on floricanes leaflet: upper surface, pubescence on floricanes leaflet: undersurface, floricanes leaf color abaxial, floricanes leaf color adaxial, petiole length, petiolule length: terminal leaflet, petiolule length: first distal leaflet, petiolule color: abaxial, petiolule color: adaxial, stipule length, and stipule width.

TABLE 3

Characteristic	'Twilight'
Mature compound leaf width (cm)	16.65
Mature compound leaf length (cm)	13.44
Number of leaflets per floricanes compound leaf	3.00
Mature leaflet shape	Ovate
Mature leaflet apex	Acute
Mature leaflet base	Roundly truncate
Mature terminal leaflet width (cm)	7.25
Mature terminal leaflet length (cm)	9.25
Mature first lateral leaflet width (cm)	6.43
Mature first lateral leaflet length (cm)	8.52
Leaflet margin	Serrated
Leaflet serration teeth length	0.33
Leaflet serration teeth width at base	0.38
Pubescence on floricanes leaflet: upper surface	Light pubescence
Pubescence on floricanes leaflet undersurface	Heavy, dense pubescence
Floricanes leaf color abaxial	137A
Floricanes leaf color adaxial	138B
Petiole length (cm)	5.72
Petiole color adaxial	165A
Petiole color abaxial	N144D
Petiolule length terminal leaflet (cm)	1.52
Petiolule length first distal leaflet (cm)	2.11
Petiolule color abaxial	165A
Petiolule color adaxial	N144D
Stipule length (cm)	1.00
Stipule width (cm)	0.24
Stipule attitude	Erect

Table 4 shows flower and flowering characteristics of the new cultivar. Flower and flowering characteristics include date 1st bloom, date full bloom, date last bloom, petal color, the number flowers per cluster, the number of petals per flower, flower diameter, petal length, petal width, the num-

ber of sepals per flower, peduncle length, rachis length, peduncle color, and cyme type.

TABLE 4

Characteristic	'Twilight'
Date 1 st bloom	May 13
Date full bloom	May 20
Date last bloom	May 30
Petal color adaxial and abaxial surfaces	69C
Number flowers per cluster	8.83
Number of petals per flower	5.33
Flower diameter (cm)	32.45
Petal length (cm)	14.28
Petal width (cm)	11.33
Petal texture adaxial and abaxial	No pubescence
Number of sepals per flower	4.83
Peduncle length (cm)	3.67
Rachis length (cm)	10.85
Peduncle color	N144A streaked w/185A
Cyme type	Elongate simple cyme

Table 5 shows fruit and fruiting characteristics of the new cultivar. Fruit and fruiting characteristics include date 5% of fruit were ripe, date 50% of fruit were ripe, date 95% of fruit were ripe, weight of primary fruit, weight of secondary fruit, diameter of primary fruit at equator, diameter of 2° fruit at equator, diameter of 1° fruit at poles: tip, diameter of 1° fruit at poles: base, diameter of 2° fruit at poles: tip, diameter of 2° fruit at poles: base, berry length 1° fruit, berry length 2° fruit, ratio of primary fruit length to width, shape description, uniformity of berry shape, color when full ripe, number of drupelets per fruit, individual seed weight, glossiness, firmness, flavor, texture of fruit when chewed, drupelet skin resistance to abrasion, ease of separation of fruit from pedicel, machine harvestability, resistance to heat damage of fruit, berries per inflorescence—mean, berries per inflorescence range, soluble solids (%; in Brix), pH, titratable acidity (% as citric acid), yield (actual kg·plt⁻¹), disease response, and red berry mite response.

TABLE 5

Characteristic	'Twilight'
Date 5% of fruit were ripe	Jul. 10
Date 50% of fruit were ripe	Jul. 21
Date 95% of fruit were ripe	Aug. 9
Weight of primary fruit (g)	8.03
Weight of secondary fruit (g)	5.91
Weight of tertiary fruit (g)	6.00
Diameter of primary fruit at equator (cm)	2.16
Diameter of 2 fruit at equator (cm)	2.22
Diameter of 3 fruit at equator (cm)	2.03
Diameter of 1 fruit at poles: tip (cm)	1.70
Diameter of 1 fruit at poles: base (cm)	2.18
Diameter of 2 fruit at poles: tip (cm)	1.70
Diameter of 2 fruit at poles: base (cm)	2.08
Diameter of 3 fruit at poles: tip (cm)	1.61
Diameter of 3 fruit at poles: base (cm)	1.96
Berry length primary fruit (cm)	2.85
Berry length 2 fruit (cm)	2.51
Berry length 3 fruit (cm)	2.47
Ratio of primary fruit length to width	1.32
Shape description	Blocky
Uniformity of berry shape	Consistent shape
Color	when full ripe 203A
Number of drupelets per fruit	68.50
Total seed weight per fruit (mg)	199.30
Individual seed weight (mg)	2.96
Glossiness	Very glossy

TABLE 5-continued

Characteristic	'Twilight'
Firmness	Very firm
5 Flavor	Very good flavor
Texture of fruit when chewed	Good
Drupelet skin resistance to abrasion	Excellent
Ease of separation of fruit from pedicel	Medium
Machine harvestability	Good
10 Resistance to heat damage of fruit	Good
Berries per inflorescence - mean	7.50
Berries per inflorescence range	7-8
Soluble solids (%; in Brix)	13.07
pH	3.51
Titratable acidity (% as citric acid)	8.58
15 Yield (actual kg · plt ⁻¹)	9.78
Disease response	Under a typical, minimal, disease management program does not exhibit any particular disease problems
Red berry mite response	Moderate susceptibility

COMPARISON WITH PARENTAL AND COMMERCIAL CULTIVARS

'Twilight' differs from the female parent blackberry plant 'Ouachita' (U.S. Plant Pat. No. 17,762) in that it has a semi-erect type plant growth habit, is more productive, with larger (8.7 g) and sweeter (13% Brix) fruit that have a pleasant crispness when eaten, while 'Ouachita' (U.S. Plant Pat. No. 17,762) has an erect type plant growth habit, is less productive, has smaller (7.0 g) and less sweet (11% Brix) fruit that, while firm, are not crisp when eaten.

'Twilight' differs from the male parent blackberry plant 'ORUS 2867-4' (unpatented) in that it is 7 days earlier ripening with large sized (8.7 g) fruit that are firm with a tough skin and have a lower incidence of fruit defects due to excessive heat and UV light damage, while 'ORUS 2867-4' (unpatented) is 7 days later ripening with medium-large (7.2 g) fruit that are soft with tender skin and are prone to heat and UV light damage.

'Twilight' is 7-14 days earlier ripening than other commercial semi-erect fresh market blackberries such as 'Triple Crown' (unpatented) or 'Chester Thornless' (unpatented). 'Twilight' ripens 4-5 d later than the semi-erect blackberry 'Loch Ness' (U.S. Plant Pat. No. 6,782), 'Eclipse' (U.S. patent application Ser. No. 15/731,503), 'Galaxy' (U.S. Plant Pat. No. 30,062), and 'Von' (U.S. Plant Pat. No. 27,299). 'Twilight' is a fresh market blackberry and has had excellent fruit quality after 7 days in refrigerated storage in plastic clam shell packaging at 1° C. compared to fruit from 'Eclipse' (U.S. patent application Ser. No. 15/731,503), 'Triple Crown' (unpatented), and 'Von' (U.S. Plant Pat. No. 27,299). 'Twilight' has a pleasantly crisp, firm texture when eaten, firmer than 'Triple Crown' (unpatented), and with fewer noticeable seeds than 'Chester Thornless' (unpatented). 'Twilight' was as high yielding as 'Chester Thornless' (unpatented) and 'Triple Crown' (unpatented), and higher yielding than 'Eclipse' (U.S. patent application Ser. No. 15/731,503). 'Twilight' fruit are sweeter than those of 'Chester Thornless' (unpatented) but not as sweet as those of 'Triple Crown' (unpatented) and comparable to 'Eclipse' (U.S. patent application Ser. No. 15/731,503).

I claim:

1. A new and distinct cultivar of thornless blackberry plant, substantially as illustrated and described, characterized by its large sized fruit that are sweet, crisp, firm, and have a very tough skin.

5

* * * * *




FIG. 1




FIG 2




FIG. 3



FIG. 4