



US00PP31035P3

(12) **United States Plant Patent**  
**Bell**

(10) **Patent No.:** **US PP31,035 P3**  
(45) **Date of Patent:** **Nov. 12, 2019**

(54) **BLUEBERRY PLANT NAMED ‘RIDLEY 1607’**

(50) Latin Name: *Vaccinium* hybrid  
Varietal Denomination: **Ridley 1607**

(71) Applicant: **MOUNTAIN BLUE ORCHARDS, PTY LTD**, Lindendale, New South Wales (AU)

(72) Inventor: **Ridley Bell**, Lindendale (AU)

(73) Assignee: **Mountain Blue Orchards, Pty Ltd**, Lindendale (AU)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/998,145**

(22) Filed: **Jul. 10, 2018**

(65) **Prior Publication Data**

US 2019/0037747 P1 Jan. 31, 2019

**Related U.S. Application Data**

(60) Provisional application No. 62/605,029, filed on Jul. 28, 2017.

(51) **Int. Cl.**

*A01H 5/08* (2018.01)  
*A01H 6/36* (2018.01)

(52) **U.S. Cl.**

USPC ..... **Plt./157**  
CPC ..... *A01H 6/368* (2018.05)

(58) **Field of Classification Search**

USPC ..... Plt./205, 157  
CPC ..... *A01H 5/0812*  
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP22,778 P3 6/2012 Wright et al.  
PP25,432 P3 4/2015 Bell  
PP29,469 P3 7/2018 Bell

*Primary Examiner* — Kent L Bell

(74) *Attorney, Agent, or Firm* — Plant & Planet Law Firm

(57) **ABSTRACT**

The new blueberry plant variety ‘Ridley 1607’ is provided. ‘Ridley 1607’ is a commercial variety intended for use as fresh fruit for shipping, hand pick, customer pick, machine harvest and processing markets and as a home garden plant. The variety is produced from a cross of ‘Ridley 1403’ (U.S. Plant Pat. No. 25,432) and ‘Ridley 4609’ (U.S. Plant Pat. No. 29,469), which can be distinguished by its outstanding features.

**7 Drawing Sheets**

Latin name of the genus and species:

Genus—*Vaccinium*.

Species—hybrid.

Variety denomination: The new blueberry plant claimed is of the variety denominated ‘RIDLEY 1607’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct perennial variety of *Vaccinium* hybrid (Southern Highbush Blueberry), which has been given the variety denomination of ‘Ridley 1607’. Its market class is that of a fruiting plant. ‘Ridley 1607’ is intended for use as fresh fruit for shipping, customer pick and processing markets and as a home garden plant.

The new *Vaccinium* hybrid cultivar is a selection resulting from seedlings produced in a breeding program of *Vaccinium* at Lindendale, NSW, Australia in 2011 from the controlled pollination of seed parent ‘Ridley 1403’ (U.S. Plant Pat. No. 25,432) with pollen parent ‘Ridley 4609’ (U.S. Plant Pat. No. 29,469). The new cultivar was discovered and selected as a single plant within a population of 100 resulting *Vaccinium* hybrid plants from this controlled pollination in 2014 in a commercial field plantation environment at Lindendale, New South Wales, Australia. Selection criteria were a combination of strong plant growth vigor, upright to semi-upright plant growth habit, low chilling requirement, late to very late time of flowering and fruit

ripening, medium to large, sweet, firm, crisp fruit suited to handling, desirable tasting berries and small picking scar.

The selection was subsequently evaluated for a number of years at the commercial farms at Lindendale, New South Wales, Australia and Tabulam, New South Wales, Australia.

Asexual reproduction of the new cultivar by softwood cutting propagation since 2014 at Lindendale, New South Wales, Australia has demonstrated that the new cultivar reproduces true to type with all of the characteristics, as herein described, firmly fixed and retained through successive generations of such asexual propagation.

**SUMMARY OF THE INVENTION**

The new *Vaccinium* hybrid cultivar is a selection resulting from seedlings produced in a breeding program of *Vaccinium* at Lindendale, NSW, Australia in 2011 from the controlled pollination of seed parent ‘Ridley 1403’ with pollen parent ‘Ridley 4609’. The seed parent was produced from a seedling selection of ‘S02-25-05’ (not patented) and pollen parent ‘S03-08-02’ (not patented). The pollen parent was produced from a seedling selection of ‘C95-152’ (not patented) and pollen parent ‘C00-09’ (U.S. Plant Pat. No. 22,778).

Plants of the new cultivar differ from plants of the seed parent ‘Ridley 1403’ primarily in ripening two weeks later than ‘Ridley 1403’ and is easier to shake and harvest than ‘Ridley 1403’. Plants of the new cultivar differ from plants

of the pollen parent 'Ridley 4609' primarily in being more upright in growth habit than 'Ridley 4609' and has firmer, sweeter fruit that is easier to shake and harvest than 'Ridley 4609'.

The new blueberry variety was designated M14-16-07, and has been planted in replicated trials since 2014.

The new *Vaccinium* hybrid cultivar is a selection resulting from seedlings produced in a controlled breeding program of *Vaccinium* at Lindendale, NSW, in 2014 from the controlled pollination of seed parent 'Ridley 1403' with pollen parent 'Ridley 4609'.

The following characteristics of the new cultivar have been repeatedly observed and can be used to distinguish M14-16-07 ('Ridley 1607') as a new and distinct cultivar of *Vaccinium* hybrid plant:

1. Upright to semi-upright bushy plant growth habit.
2. Strong plant growth vigor.
3. Large leaf size.
4. Late to very late timing of flowering beginning.
5. Late to very late timing of fruit ripening.
6. Very firm and crisp berry.
7. Dry picking scar.
8. Medium to large berry size.
9. Berry sweetness is high to very high.

Asexual reproduction of the new cultivar by cutting propagation since 2014 at Lindendale, New South Wales, Australia has demonstrated that the new cultivar reproduces true to type with all of the characteristics, as herein described, firmly fixed and retained through successive generations of such asexual propagation.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying photographic illustrations shows typical specimens in full color of the foliage and fruit of the new variety 'Ridley 1607' The colors are as nearly true as is reasonably possible in a color representation of this type.

FIG. 1 is a photograph of the new variety 'Ridley 1607', demonstrating the plant's upright growth habit.

FIG. 2 is a photograph of the fruit of the new variety 'Ridley 1607'.

FIG. 3 is a photograph of the flowers of the new variety 'Ridley 1607'.

FIG. 4 is a photograph of the leaves of the new variety 'Ridley 1607'.

FIG. 5 is a photograph of the leaves, flowers, fruit, new shoot and fruit cluster of the new variety 'Ridley 1607'.

FIG. 6 is a photograph of fruit of the variety 'Ridley 1607'.

FIG. 7 is a photograph of fruit cross section of the variety 'Ridley 1607'.

The colors in the photographs are as close as possible with the photographic and printing technology utilized. The color values cited in the detailed botanical description accurately describe the colors of the new blueberry.

#### DETAILED BOTANICAL DESCRIPTION

The following detailed description sets forth the distinctive characteristics of 'Ridley 1607'. The data which define these characteristics were collected from asexual reproductions of the original selection. Dimensions, sizes, colors, and other characteristics are approximations and averages set forth as accurately as possible. The plant history was taken on plants approximately 3 years of age, and the descriptions relate to plants grown in the field in Tabulam, New South

Wales, Australia. Descriptions of fruit characteristics were made on fruit grown in Tabulam, New South Wales, Australia. Color designations are from Tabulam, New South Wales, Australia.

Except where noted, color notations are based on *The Royal Horticultural Society Colour Chart*, of The Royal Horticultural Society, London, 2015 edition.

Classification:

- a. *Family*.—Ericaceae.
- b. *Genus*.—*Vaccinium*.
- c. *Species*.—Hybrid.
- d. *Common name*.—Southern Highbush Blueberry.

Parentage:

*Female parent*.—'Ridley 1403' (U.S. Plant Pat. No. 25,432).

*Male parent*.—'Ridley 4609' (U.S. Plant Pat. No. 29,469).

Market class: A fruiting plant intended for use as fresh fruit for shipping, hand pick, customer pick, machine harvest and processing markets and as a home garden plant.

#### PLANT

General:

*Parentage*.—Female Parent — 'Ridley 1403' (U.S. Plant Pat. No. 25,432); Male Parent — 'Ridley 4609' (U.S. Plant Pat. No. 29,469).

*Plant height*.—1.7 m.

*Plant width*.—1.5 m.

*Growth habit*.—Upright to semi-upright.

*Growth*.—Strong to very strong vigor.

*Productivity*.—6 to 7 kg per season from 3-4 year old plants planted at 3.0 m×0.9 m density.

*Cold hardiness*.—Has not been grown in all environments including harsh winter environments.

*Cold tolerance*.—Cold tolerance is expected to be low.

*Chilling requirement*.—Has not been grown in all environments and is typically grown as an evergreen crop where chilling hours are not important. 'Ridley 1607' is classed as "low chill," typical of Southern Highbush Blueberry varieties with an estimated chilling requirement of 250 hours (not tested).

*Tolerance to disease*.—Moderate resistance to root disease (*Phytophthora* spp) and good resistance to blueberry rust.

*Leafing*.—Vegetative bud burst is early when grown as an evergreen in Australia (August in Tabulam, NSW, Australia).

*Twigginess*.—Low.

#### STEM

General:

*Suckering tendency*.—Plants typically have 5-7 major canes per plant from a base 30 cm in diameter on 6 year old plants.

*Mature cane color*.—Greyed orange 198C.

*Mature cane length*.—0.8-1.0 m.

*Mature cane width*.—1.2 cm.

*Bark texture*.—Medium roughness (typical of species).

*Fall color on new shoots*.—Top side (7419C/7420C) Under side (7488CP) (Pantone color bridge plus).

*Surface texture of new wood*.—Smooth.

*Internode length on strong, new shoots*.—20-30 mm.

*Fruiting wood*.—To 15 cm in length.

## FOLIAGE

## General:

*Time of beginning of leaf bud burst.*—Early (August in Tabulam, NSW, Australia).

*Leaf color (top side).*—Yellow green 147A.

*Leaf color (under side).*—Yellow green 147C.

*Leaf arrangement.*—Alternate.

*Leaf shape.*—Elliptic.

*Leaf margins.*—Entire.

*Undulation of margin.*—Weak.

*Leaf venation.*—Reticulate.

*Leaf apices.*—Acuminate.

*Leaf bases.*—Obtuse.

*Leaf length.*—63-69 mm.

*Leaf width.*—31-38 mm.

*Leaf length/width ratio.*—Medium (1.92:1).

*Leaf nectaries.*—Absent.

*Pubescence of upper side.*—Absent.

*Pubescence of lower side.*—Absent.

*Cross sectional profile.*—Flat.

*Longitudinal profile.*—Straight.

*Attitude.*—Semi-upright to horizontal.

## Petioles:

*Length.*—1.0 to 3.0 mm.

*Width.*—2.0 mm.

*Color.*—Yellow green 146C-147C.

## FLOWERS

## General:

*Time of beginning of flowering.*—Late to very late (early October to-early November at Tabulam, NSW Australia).

*Time of 50% anthesis.*—Late October at Tabulam, NSW Australia.

*Flower shape.*—Urceolate.

*Flower bud density.*—Medium density.

*Flower fragrance.*—Weak.

## Corolla:

*Color.*—White NN155D.

*Length.*—9-12 mm.

*Width.*—7-9 mm.

*Aperture width.*—3-5 mm.

*Anthocyanin coloration of corolla.*—Absent, occasional in bud stage.

*Corolla ridges.*—Present.

*Protrusion of stigma.*—Present.

## Inflorescence:

*Length.*—40 to 70 mm.

*Diameter.*—20-40 mm.

*Length of peduncle.*—15 mm.

*Surface texture of peduncle.*—Smooth.

*Color of peduncle.*—Yellow green 146C.

*Length of pedicel.*—7-10 mm.

*Surface texture of pedicel.*—Smooth.

*Color of pedicel.*—Yellow green 147C.

*Number of flowers per cluster.*—9-12.

*Flower cluster density.*—Medium.

## Calyx (with sepals):

*Diameter.*—9-11 mm.

*Color (sepals).*—Green 138C to yellow green 147C.

## Stamen:

*Length.*—4-5 mm.

*Number per flower.*—Approximately 10.

*Filament color.*—Yellow green 144D.

## Pistil:

*Length.*—10-13 mm.

*Style.*—Length — 5-6 mm.

*Ovary color (exterior).*—Yellow green 144D-147C.

## Anther:

*Length.*—3.0-3.5 mm.

*Number.*—Approximately 10.

*Color.*—Greyed orange 167C-D.

## Pollen:

*Abundance.*—Medium.

*Color.*—Greyed orange 167C-D.

*Self-compatibility.*—60% fruit set with own pollen in tests at Lindendale, NSW Australia.

## FRUIT

## General:

*Time of fruit ripening.*—Late to very late (late October to-mid December at Tabulam, NSW Australia).

*Time of 50% maturity.*—Mid November.

*Fruit development period.*—65 days.

*Cluster density.*—Medium, 7-10 berries per cluster.

*Unripe fruit color.*—Green 138A.

*Ripe berry color.*—Black 202A.

*Berry surface wax abundance.*—Strong.

*Berry flesh color.*—Greyed green 194C, lightening to 194D, then 193D.

*Berry weight.*—3.0-3.6 g.

*Berry height from calyx to scar.*—14-18 mm.

*Berry diameter.*—20 mm.

*Berry shape.*—Oblate.

*Fruit stem scar.*—Small (dry).

*Sweetness when ripe.*—High to very high as compared to market standard.

*Firmness when ripe.*—Very firm.

*Acidity when ripe.*—Medium as compared to market standard.

*Storage quality.*—Good, lasted 10 weeks at 2 degrees Celsius in tests. Not tested with modified atmosphere storage yet.

*Suitability for mechanical harvesting.*—Very firm fruit firmness, strong blush and fruit shape suited to mechanical harvesting. Shake requirement and season compactness not yet tested.

*Self-fruitfulness.*—60% fruit set with own pollen in tests at Lindendale, NSW Australia.

*Use.*—Fresh fruit for shipping, customer pick and processing markets and as a home garden plant.

## SEED

## General:

*Seed abundance in fruit.*—Abundant.

*Seed color.*—Greyed orange 164A.

*Seed dry weight.*—Not recorded.

*Seed length.*—1.5-2.0 mm.

COMPARISON BETWEEN COMMERCIAL CULTIVARS

TABLE 1

Organ	Context	'Ridley 1607'	'Ridley 1602' (not patented)	'Ridley 1105' (not patented)
Plant	Vigour	strong to very strong	strong to very strong	strong
Plant	growth habit	upright to semi-upright	upright to semi-upright	upright
One-year-old shoot	Color	green	green	green
One-year-old shoot	length of internode	medium	medium	medium
Leaf	Length	long	long	long to very long
Leaf	Width	broad	broad	very broad
Leaf	Shape	elliptic	elliptic	elliptic
Leaf	color of upper side	green	green	green
Leaf	intensity of green color on upper side (varieties with green leaf color only)	medium	medium	medium
Leaf	margin	entire	entire	entire
Flower bud	anthocyanin coloration	weak	weak	weak
Flower	shape of corolla	urceolate	urceolate	urceolate
Flower	size of corolla tube	medium	medium	medium
Flower	anthocyanin coloration of corolla tube	absent or very weak	absent or very weak	absent or very weak
Flower	ridges on corolla tube	present	present	present

TABLE 1-continued

Organ	Context	'Ridley 1607'	'Ridley 1602' (not patented)	'Ridley 1105' (not patented)
Fruit cluster	density	medium	medium	medium
Unripe fruit	intensity of green color	light	light	light
Fruit	size	medium to large	large to very large	large to very large
Fruit	shape in longitudinal section	oblate	oblate	oblate
Fruit	attitude of sepals	erect	erect	erect
Fruit	diameter of calyx basin	large to very large	large to very large	medium to large
Fruit	depth of calyx basin	deep	deep	medium
Fruit	intensity of bloom	strong	strong	strong
Fruit	color of skin	dark blue	dark blue	dark blue
Fruit	firmness	very firm	very firm	firm to very firm
Fruit	sweetness	high to very high	high to very high	high
Fruit	acidity	medium	medium to high	medium to high
Time of vegetative bud burst	vegetative bud burst	early	very early	very early
Time of beginning of flowering	beginning of flowering	late to very late	very early to early	very early
Time of beginning of fruit ripening	beginning of fruit ripening	late to very late	very early to early	very early

The invention claimed is:

1. A new and distinct variety of blueberry plant named 'Ridley 1607', substantially as illustrated and described herein.

\* \* \* \* \*



Fig. 1




Fig. 2



Fig. 3




Fig. 4






Fig. 5



Fig. 6



Fig. 7