

**(12) United States Plant Patent  
Bergman****(10) Patent No.: US PP31,048 P2****(45) Date of Patent: Nov. 12, 2019****(54) CHRYSANTHEMUM PLANT NAMED  
'CIDZ0093'****(50) Latin Name: *Chrysanthemum x morifolium*  
Varietal Denomination: CIDZ0093****(71) Applicant: SYNGENTA PARTICIPATIONS AG,  
Basel (CH)****(72) Inventor: Wendy R. Bergman, Gilroy, CA (US)****(73) Assignee: Syngenta Participations AG, Basel  
(CH)****(\*) Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.**(21) Appl. No.: 16/350,517****(22) Filed: Nov. 28, 2018****(51) Int. Cl.**  
*A01H 5/02* (2018.01)  
*A01H 6/14* (2018.01)**(52) U.S. Cl.**  
USPC ..... **Plt./295****(58) Field of Classification Search**  
USPC ..... **Plt./263.1, 284, 286, 295**  
See application file for complete search history.*Primary Examiner* — Susan McCormick Ewoldt*Assistant Examiner* — Karen M Redden**(74) Attorney, Agent, or Firm** — Dale Skalla**(57) ABSTRACT**A new *Chrysanthemum* plant named 'CIDZ0093' having yellow daisy-type inflorescences, excellent flower production, very vigorous growth and spreading plant habit, and very uniform flowering with response time of 7.5 weeks when grown under short days.**1 Drawing Sheet****1**Latin name of the genus and species of the plant claimed:  
*Chrysanthemum x morifolium*.

Varietal denomination: 'CIDZ0093'.

**BACKGROUND OF THE NEW PLANT**The present invention comprises a new *Chrysanthemum*, botanically known as *Chrysanthemum x morifolium*, and hereinafter referred to by the variety name 'CIDZ0093'.

'CIDZ0093' is a product of a planned breeding program. The new cultivar has yellow daisy-type inflorescences, excellent flower production, very vigorous growth and spreading plant habit, and very uniform flowering with response time of 7.5 weeks when grown under short days.

'CIDZ0093' originates as a natural whole plant mutation discovered among flowering plants of 'C0135' growing in a greenhouse in Gilroy, Calif. in November 2015.

The parent 'C0135' is an unpatented proprietary breeding line and having inflorescences of a pale pink hue under usual conditions, differing from 'CIDZ0093' which has yellow inflorescences.

The first act of asexual reproduction of 'CIDZ0093' was accomplished when vegetative cuttings were propagated from the initial selection in February 2016 in a greenhouse in Gilroy, Calif.

**BRIEF SUMMARY OF INVENTION**

Horticultural examination of plants grown from cuttings of the plant initiated February, 2016, and continuing thereafter, has demonstrated that the combination of characteristics as herein disclosed for 'CIDZ0093' are firmly fixed and are retained through successive generations of asexual reproduction.

'CIDZ0093' has not been observed under all possible environmental conditions. The phenotype may vary significantly with variations in environment such as temperature, light intensity and day length.

**2**

A Plant Breeder's Right for this cultivar was applied for in Canada on Nov. 19, 2018, No. 15582. 'CIDZ0093' has not been made publicly available prior to the effective filing date of this application, notwithstanding any disclosure that may have been made less than one year prior to the effective filing date of this application by the inventor or another who obtained 'CIDZ0093' directly from the inventor.

The following traits have been repeatedly observed and are determined to be basic characteristics of the new variety. The combination of these characteristics distinguishes this *Chrysanthemum* as a new and distinct variety.**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographic drawing shows typical flower and foliage characteristics of 'CIDZ0093' with colors being as true as possible with an illustration of this type. The photographic drawing shows in FIG. 1 a close-up view of the inflorescence of the new variety and in FIG. 2 a whole flowering plant grown in a greenhouse.

**DETAILED BOTANICAL DESCRIPTION**

The plant descriptions and measurements were taken indoors in Gilroy, Calif. in November 2018 on about 12 week old plants. Unrooted cuttings had been planted, 4 plants per pot, in 6 inch pots on Aug. 22, 2018. The plants were pinched and moved to short days on Sep. 19, 2018 to induce flowering.

The aforementioned photographs: FIGS. 1 and 2 was taken in Gilroy, Calif. in November 2018, and showing a plant from a trial, in which flowering had been induced by applying black cloth.

These plants were about 12 weeks of age. Unrooted cuttings had been planted, 4 plants per pot, in 6 inch pots. The plants were pinched and moved to short days to induce flowering.

Color references are made to The Royal Horticultural Society Colour Chart (R.H.S.) 2001.

TABLE 1

DIFFERENCES BETWEEN THE NEW VARIETY 'CIDZ0093' AND A MOST SIMILAR VARIETY:		
	'CIDZ0093'	'Yochesapeake' (U.S. Plant Pat. No. 12,535)
Plant height:	Shorter	Taller
Flower color:	Medium yellow	Stronger yellow
Flowering response:	Faster	Slower

## Plant:

*Form, growth and habit.*—Herbaceous pot-type, stems upright, freely branching, strong and moderately vigorous growth habit.

*Plant height.*—24-26 cm.

*Plant height (inflorescence included).*—29-30 cm.

*Plant width.*—25-27 cm.

## Roots:

*Number of days to initiate roots.*—4 days at about 22 degrees C.

*Number of days to produce a rooted cutting.*—10-12 days at 22 degrees C.

*Type.*—Fine, fibrous, free branching.

*Color.*—RHS N155B but whiter.

## Foliage:

*Arrangement.*—Alternate.

*Immature, leaf color, upper surface.*—RHS NN137A.

*Lower surface.*—RHS NN137C.

*Mature, leaf color, upper surface.*—RHS 137B.

*Lower surface.*—RHS 137C.

*Length.*—7.0-8.0 cm.

*Width.*—6.0-8.0 cm.

*Shape.*—Ovate

*Base shape.*—Attenuate to Divergent.

*Apex shape.*—Obtuse.

*Margin.*—Lobed and some serrated.

*Texture, upper surface.*—Small hairs.

*Lower surface.*—Small hairs.

*Color of veins, upper surface.*—RHS 138A.

*Color of veins, lower surface.*—RHS 138A.

*Pattern of veining.*—Palmate to Pinnate.

*Petiole color.*—RHS 138B.

*Length.*—1.6-1.8 cm.

*Diameter.*—0.2 cm.

*Texture.*—Small hairs.

## Stem:

*Quantity of main branches per plant.*—4-5.

*Color of stem.*—RHS 137A.

*Length of stem.*—20-22 cm.

*Diameter.*—0.3-0.5 cm.

*Length of internodes.*—1.5 cm.

*Texture.*—Small hairs.

*Color of peduncle.*—RHS 137A.

*Length of peduncle.*—1 cm.

*Peduncle diameter.*—0.2 cm.

*Texture.*—Hairs.

## Inflorescence:

*Type.*—Compositae type, solitary daisy-type inflorescences borne terminally above foliage, ray florets arranged acropetally on a capitulum.

*Quantity of short days to flowering (response time).*—7.5 weeks.

*Natural season flowering.*—Not determined on this variety.

*Quantity of inflorescences per plant.*—16-18 flowers.

*Lastingness of individual blooms on the plant.*—4 weeks.

*Fragrance.*—Daisy like.

## Bud (just when opening/showing color):

*Color.*—RHS 1B.

*Length.*—1.6 cm.

*Width.*—1.1-1.3 cm.

*Shape.*—Oblate.

## Immature inflorescence:

*Diameter.*—5.0-7.0 cm.

*Color of ray florets, upper surface.*—RHS 5A.

*Lower surface.*—RHS 5D.

## Mature inflorescence: Not fully mature.

*Diameter.*—7.0-8.2 cm.

*Depth.*—2.5-3.1 cm.

*Total diameter of disc.*—1.7-2.1 cm.

*Receptacle color.*—RHS 149C.

*Receptacle height.*—0.3 cm.

*Receptacle diameter.*—0.7 cm.

## Ray florets:

*Average quantity of florets.*—20-24.

*Color of ray florets, upper surface.*—RHS 2A.

*Lower surface.*—RHS 3D.

*Length.*—3.6 cm.

*Width.*—1.1-1.2 cm.

*Shape.*—Eliptical.

*Apex shape.*—Acute.

*Margin.*—Entire.

*Texture, upper surface.*—Smooth.

*Lower surface.*—Smooth.

## Disc florets:

*Average quantity of florets.*—180-210.

*Color of florets.*—RHS 6A to RHS 145B lower end.

*Length.*—0.4-0.6 cm.

*Width.*—0.2 cm.

*Shape.*—Tubular, elongated.

*Apex shape.*—Acute, 5 pointed.

## Phyllaries:

*Quantity.*—20-25.

*Color, upper surface.*—RHS 137A.

*Lower surface.*—RHS 137B.

*Length.*—0.7-1.1 cm.

*Width.*—0.2-0.3 cm.

*Shape.*—Lanceolate.

*Apex shape.*—Acute.

*Base.*—Fused.

*Margins.*—Entire; slightly papery.

*Texture, upper surface.*—Hair.

*Lower surface.*—Smooth.

## Reproductive organs:

*Pistil.*—1, found on both types of florets.

*Length.*—0.7 cm.

*Style color.*—RHS 1D.

*Style length.*—0.5 cm.

*Stigma color.*—RHS 5A.

*Stigma shape.*—Bi-parted.

*Ovary color.*—RHS 155C but more translucent.

*Stamens.*—4, found only on the disc florets.

*Color of filaments.*—RHS 155C.

*Length filaments.*—0.4-0.5 cm.

*Anther color.*—RHS 15A.

*Anther length.*—0.15 cm.

*Anther shape.*—Oval.  
*Color of pollen.*—RHS 17B.  
*Pollen amount.*—Sparse.  
*Fertility/seed set.*—Has not been observed to date.  
*Disease/pest resistance.*—Has not been observed to date.

What is claimed is:

1. A new and distinct variety of *Chrysanthemum* plant named 'CIDZ0093' substantially as illustrated and described herein.

\* \* \* \* \*



FIG. 1




FIG. 2