

(12) **United States Plant Patent**
Pineau

(10) **Patent No.:** **US PP31,305 P2**
(45) **Date of Patent:** **Dec. 31, 2019**

(54) **HIBISCUS PLANT NAMED ‘MINSYRE10’**
(50) Latin Name: *Hibiscus syriacus*
Varietal Denomination: **MINSYRE10**
(71) Applicant: **Patrick Pineau**, Saint Mathurin sur
Loire (FR)
(72) Inventor: **Patrick Pineau**, Saint Mathurin sur
Loire (FR)
(73) Assignee: **Hortival Diffusion SAS** (FR)
(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.
(21) Appl. No.: **16/350,722**
(22) Filed: **Dec. 20, 2018**
(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/60 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./257**
(58) **Field of Classification Search**
USPC Plt./257
CPC A01H 5/02; A01H 6/60
See application file for complete search history.

Primary Examiner — Annette H Para
(74) Attorney, Agent, or Firm — Cassandra Bright

(57) **ABSTRACT**
A new and distinct *Hibiscus* cultivar named ‘MINSYRE10’
is disclosed, characterized by large, single, dark magenta
flowers with a darker eye, up to 11 cm in diameter. Plants are
well-branched, upright and vigorous and have been
observed to have some resistance to Sclerotinia. The new
variety is a *Hibiscus*, normally produced as an outdoor
garden or container plant.

3 Drawing Sheets

1

Latin name of the genus and species: *Hibiscus syriacus*.
Variety denomination: ‘MINSYRE10’.

BACKGROUND OF THE INVENTION

The new *Hibiscus* cultivar is a product of a planned
breeding program conducted by the inventor, Patrick Pineau.
The objective of the breeding program was to vigorous
Hibiscus syriacus cultivars with healthy foliage. The cross
resulting in this new variety was made during 2008.

The seed parent is the unpatented *Hibiscus syriacus*
‘Flogu’. The pollen parent is the unpatented variety *Hibiscus*
syriacus ‘Freedom’. The new variety was identified and
selected as a potentially interesting selection in September
2012. Selection was made at a nursery in La Menitre,
France.

Asexual reproduction of the new cultivar ‘MINSYRE10’
was first performed in January 2013 in La Menitre, Maine Et
Loire, France, by grafting. Subsequent propagation has
shown that the unique features of this cultivar are stable and
reproduced true to type in 6 successive generations.

SUMMARY OF THE INVENTION

The cultivar ‘MINSYRE10’ has not been observed under
all possible environmental conditions. The phenotype may
vary somewhat with variations in environment such as
temperature, day length, and light intensity, without, how-
ever, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of
‘MINSYRE10’ These characteristics in combination distin-
guish ‘MINSYRE10’ as a new and distinct *Hibiscus* culti-
var:

1. Dark magenta flower.
2. Large single flower.
3. Erect plant form.
4. Well branched.

2

5. Vigorous.
6. Some observed resistance to Sclerotinia.

PARENT COMPARISON

Plants of the new cultivar ‘MINSYRE10’ are similar to
plants of the seed parent variety in most horticultural char-
acteristics, however, plants of the new cultivar
‘MINSYRE10’ differ in the following;

1. Minsyre10 is more vigorous and has more branches
than the seed parent.
2. Minsyre10 has longer and larger leaves than the seed
parent.
3. Minsyre10 has less leaf blade lobing than the seed
parent.
4. Minsyre10 has longer and larger petals than the seed
parent.
5. Minsyre10 has a dark red purple petal color, the seed
parent has a more purple petal color.
6. Minsyre10 has a darker red basal blotch than the seed
parent.

Plants of the new cultivar ‘MINSYRE10’ are similar to
plants of the pollen parent variety in most horticultural
characteristics, however, plants of the new cultivar
‘MINSYRE10’ differ in the following;

1. Minsyre10 is more vigorous with more branches than
pollen parent.
2. Minsyre10 has more lobing on the leaf blade than the
pollen parent.
3. Minsyre10 is a single flower, the pollen parent has a
double flower.
4. Minsyre10 has better pest resistance to Sclerotinia than
the pollen parent.

COMMERCIAL COMPARISON

Plants of the new cultivar ‘MINSYRE10’ can be com-
pared to the variety *Hibiscus* HAWAII ‘Woodbridge’,
unpatented. These varieties are similar in most horticultural
characteristics however, ‘MINSYRE10’ differs in the fol-
lowing:

1. Minsyre10 is more vigorous and with more branches than 'Woodbridge'.
2. Minsyre10 has darker red-purple petals than 'Woodbridge'.
3. Minsyre10 has more overlapping petals than 'Woodbridge'.
4. Minsyre10 has longer and larger petals than 'Woodbridge'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photograph in FIG. 1 illustrates in full color a typical plant of 'MINSYRE10' grown in the ground at approximately 3 years old in La Minetre, France.

FIG. 2 illustrates in full color a typical flower of 'MINSYRE10'.

FIG. 3 compares a flower of 'MINSYRE10', on the top of the figure, to a flower of 'Woodridge', on the bottom of the photo. The photographs were taken using conventional techniques and although colors may appear different from actual colors due to light reflectance it is as accurate as possible by conventional photographic techniques.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart 2015, 6th edition, except where general terms of ordinary dictionary significance are used. The following observations and measurements describe 'MINSYRE10' plants grown in an unheated polyhouse in Menitre, France under natural lighting. Measurements were taken during the August of 2018. The plants were about 3 years old in 4.5 liter containers. The growing temperature ranged from approximately -5° C. to 27° C. Measurements and numerical values represent averages of typical plant types.

Botanical Classification: *Hibiscus syriacus* 'MINSYRE10'.

PROPAGATION

Time to Initiate Roots: 20 days at approximately 25° C.
 Root Description: Fibrous, medium root thickness. Tan/brown in color, not effectively measured with a color chart.
 Time to Produce a Rooted Young Plant: About 90 days at 25° C.

PLANT

Plant Type: Perennial, hardy deciduous flowering shrub.
 Age of Plant Described: Approximately 3 years old.
 Growth Habit: Upright and vigorous.
 Height: 100 cm.
 Branching Characteristics: Well branched.
 Length of Primary Lateral Branches: 90 cm.
 Quantity of Lateral Branches: 8.
 Characteristics of Primary Lateral Branches:
 Diameter.—0.4 cm.
 Color.—RHS Yellow-Green 147C.
 Internode Length: 5 cm.

FOLIAGE

Leaf:

Arrangement.—Alternate.

Quantity.—Approximately 25 per branch.

Average Length.—11 cm.

Average Width.—6 cm.

Shape of blade.—Ovate, 3-lobed.

Apex.—Acute.

Base.—Obtuse.

Margin.—Lobed, crenate.

Texture of top surface.—Glabrous.

Texture of bottom surface.—Rough, with sparse adpressed hairs along veins.

Color.—Young foliage upper side: RHS Green 137R.

Young foliage under side: RHS Yellow-Green 144C.

Mature foliage upper side: RHS Green 137A. Mature

foliage under side: RHS Yellow-Green 144C.

Fall or Winter coloration variance.—Mature foliage upper side: None. Mature foliage under side: None.

Venation.—Type: Pinnate. Venation color upper side: RHS Yellow-Green 145A. Venation color under side: RHS Green 137B.

Petiole.—Length: 1.8 cm. Diameter: 0.2 cm. Color: RHS Yellow-Green 138A.

FLOWER

Natural Flowering Season: July to September.

Flower Type and Habit: Solitary, rotate.

Single, double, multiplex.—Single.

Petals overlapping.—Overlapping.

Rate of Flower Opening: 1 to 2 days from bud to fully opened flower.

Flower Longevity on Plant: 1 to 2 days.

Persistent or Self-Cleaning: Self-cleaning.

Bud:

Shape.—Ovate.

Length.—2 cm.

Diameter.—1.1 cm.

Color.—RHS Red-Purple 59A.

Flower Size:

Diameter.—11 cm.

Depth.—3 cm.

Petals:

Quantity.—5.

Arrangement.—Rotate.

Length.—6 cm.

Width.—5.5 cm.

Shape.—Rounded deltoid.

Apex.—Rounded, irregular shallow indentations.

Base.—Obovate.

Margin.—Entire.

Texture, upper and lower surfaces.—Color: When opening, upper surface: RHS Red-Purple 63A with darker edge blush near 64A. When opening, lower surface: RHS Red-Purple 63A with darker edge blush near 64A. Veins near 63D. Fully opened, upper surface: RHS Red-Purple 63B with darker edge blush near 64A. Fully opened, lower surface: RHS Red-Purple 63A with darker edge blush near 64A. Streaked near 63D. Color basal blotch: RHS Red-Purple 59A. Ratio of basal blotch/petal size: About 25% of petal blade blotched. Feathering color: RHS Red-Purple 59A. Feathering size: 0.4 to 1 cm.

Sepals:

Quantity.—5 to 6.
Length.—2 cm.
Width.—0.8 cm.
Shape.—Ovate.
Margin.—Entire.
Color.—RHS Green 143C.

Peduncles:

Length.—1 cm.
Diameter.—0.3 cm.
Angle.—About 45° to the lateral branch.
Strength.—Strong.
Texture.—Glabrous.
Color.—RHS Green 143C.

Fragrance: None.

REPRODUCTIVE ORGANS

Stamens:

Number.—50.

Anthers:

Shape.—Irregular sphere.
Length.—0.3 cm.
Color.—Near RHS White 155D.
Pollen color.—RHS Yellow 4D.
Pollen amount.—Abundant.

Pistil:

Number.—1.
Length.—3.5 cm.
Style.—Length: 3 cm. Color: Near RHS White 155C.
Stigma.—Shape: Rounded disc. Color: Near RHS White 155C.
Ovary color.—Near RHS Yellow-Green 145C.

OTHER CHARACTERISTICS

Seeds: Capsule, typically about 1 cm in diameter and length. Pubescent, colored near Grey-Brown 199B. Moderately abundant lightly pubescent seed.

Disease/Pest Resistance: Improved resistance to Sclerotinia. No other resistance nor susceptibility to normal diseases and pests of *Hibiscus syriacus* have been observed.

Temperature Tolerance: Low temperature tolerance to at least USDA zone 4, high temperature tolerance to at least 40° C.

What is claimed is:

1. A new and distinct cultivar of *Hibiscus* plant named ‘MINSYRE10’ as herein illustrated and described.

* * * * *



FIG. 1



FIG. 2



FIG. 3