



(12) **United States Plant Patent**
Magee

(10) **Patent No.:** **US PP31,556 P2**
(45) **Date of Patent:** **Mar. 17, 2020**

- (54) **CRAPEMYRTLE PLANT NAMED ‘JM6’**
- (50) Latin Name: *Lagerstroemia indica* ‘JM6’
Varietal Denomination: **JM6**
- (71) Applicant: **Jack Mitchell Magee**, Poplarville, MS (US)
- (72) Inventor: **Jack Mitchell Magee**, Poplarville, MS (US)
- (73) Assignees: **Greenleaf Nursery Company**, Park Hill, OK (US); **Evergreen Nursery**, Poplarville, MS (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **16/350,176**
- (22) Filed: **Oct. 10, 2018**

- (51) **Int. Cl.**
A01H 5/00 (2018.01)
- (52) **U.S. Cl.**
USPC **Plt./252**
- (58) **Field of Classification Search**
USPC Plt./252
See application file for complete search history.

Primary Examiner — Annette H Para
(74) Attorney, Agent, or Firm — C. A. Whealy

(57) **ABSTRACT**
A new and distinct cultivar of Crapemyrtle plant named ‘JM6’, characterized by its upright to somewhat outwardly spreading plant habit; freely branching habit; vigorous and sturdy growth habit; dark greyed purple-colored leaves; numerous inflorescences with white-colored flowers; and good garden performance and pathogen resistance.

2 Drawing Sheets

1

Botanical designation: *Lagerstroemia indica* ‘JM6’.
Cultivar denomination: ‘JM6’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of Crapemyrtle plant, botanically known as *Lagerstroemia indica* ‘JM6’ and hereinafter referred to by the name ‘JM6’.

The new Crapemyrtle plant is a product of a planned breeding program conducted by the Inventor in Poplarville, Miss. The objective of the breeding program was to develop new vigorous and freely-branching Crapemyrtle plants with dark-colored leaves and attractive flower colors.

The new Crapemyrtle plant originated from a cross-pollination conducted by the Inventor during the summer of 2015 of *Lagerstroemia indica* ‘Ebony and Ivory’, not patented, as the female, or seed, parent with *Lagerstroemia indica* ‘Natchez’, not patented, as the male, or pollen, parent. The new Crapemyrtle plant was discovered and selected by the Inventor in May, 2016 as a single flowering plant from within the progeny of the stated cross-pollination in a controlled nursery environment in Poplarville, Miss.

Asexual reproduction of the new Crapemyrtle plant by vegetative softwood cuttings in a controlled greenhouse environment in Poplarville, Miss. since August, 2016 has shown that the unique features of the new Crapemyrtle plant are stable and reproduced true to type in successive generations of asexual reproduction.

SUMMARY OF THE INVENTION

Plants of the new Crapemyrtle have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

2

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘JM6’. These characteristics in combination distinguish ‘JM6’ as a new and distinct Crapemyrtle plant:

1. Upright to somewhat outwardly spreading plant habit.
2. Freely branching habit.
3. Vigorous and sturdy growth habit.
4. Dark greyed purple-colored leaves.
5. Numerous inflorescences with white-colored flowers.
6. Good garden performance and pathogen resistance.

Plants of the new Crapemyrtle can be compared to plants of the female parent, ‘Ebony and Ivory’. Plants of the new Crapemyrtle differ primarily from plants of ‘Ebony and Ivory’ in the following characteristics:

1. Plants of the new Crapemyrtle are larger than plants of ‘Ebony and Ivory’.
2. Plants of the new Crapemyrtle are more vigorous than plants of ‘Ebony and Ivory’.

Plants of the new Crapemyrtle can be compared to plants of the male parent, ‘Natchez’. Plants of the new Crapemyrtle differ primarily from plants of Natchez in leaf color as plants of the new Crapemyrtle have dark greyed purple-colored leaves whereas plants of ‘Natchez’ have light green-colored leaves.

Plants of the new Crapemyrtle can be compared to plants of the *Lagerstroemia* L. ‘PIILAG-I’, disclosed in U.S. Plant Pat. No. 23,168. In side-by-side comparisons, plants of the new Crapemyrtle differ primarily from plants of ‘PIILAG-I’ in the following characteristics:

1. Plants of the new Crapemyrtle are taller and are more vigorous than plants of ‘PIILAG-I’.
2. Plants of the new Crapemyrtle have dark greyed purple-colored leaves whereas plants of ‘PIILAG-I’ have dark green-colored leaves.

Plants of the new Crapemyrtle can be compared to plants of the *Lagerstroemia indica* ‘Ebony Glow’, not patented. In

side-by-side comparisons, plants of the new Crapemyrtle differ primarily from plants of 'Ebony Glow' in the following characteristics:

1. Leaves of plants of the new Crapemyrtle grow faster than plants of 'Ebony Glow'.
2. Plants of the new Crapemyrtle have larger leaves than plants of 'Ebony Glow'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new Crapemyrtle plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Crapemyrtle plant.

The photograph on the first sheet (FIG. 1 of 2) is a side perspective view of a typical plant of 'JM6' grown in a container in an outdoor nursery.

The photograph on the second sheet (FIG. 2 of 2) is a close-up view of a typical flowering plant of 'JM6' grown in a container in an outdoor nursery.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown in 7-gallon containers in Park Hill, Okla. and Fort Worth, Tex. during the summer in outdoor nurseries and under cultural conditions which closely approximate commercial Crapemyrtle production. During the production of the plants, day temperatures averaged 33° C. and night temperatures averaged 14° C. Plants were three years when the photographs and the description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2015 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Lagerstroemia indica* 'JM6'.

Parentage:

Female, or seed, parent.—*Lagerstroemia indica* 'Ebony and Ivory', not patented.

Male, or pollen, parent.—*Lagerstroemia indica* 'Natchez', not patented.

Propagation:

Type.—By vegetative softwood and hardwood cuttings.

Time to initiate roots, summer.—About ten days at temperatures about 21° C. to 33° C.

Time to initiate roots, winter.—About 25 days at temperatures about 21° C. to 33° C.

Time to produce a rooted young plant, summer.—About one month at temperatures about 21° C. to 33° C.

Time to produce a rooted young plant, winter.—About one month for softwood cuttings and about two months for hardwood cuttings at temperatures about 21° C. to 33° C.

Root description.—Medium in thickness, fibrous; typically brownish white in color, actual color is dependent on substrate composition, water quality, fertilizer type and formulation, substrate temperature and physiological age of roots.

Rooting habit.—Freely branching; dense.

Plant description:

Plant form and growth habit.—Perennial shrub; upright to somewhat outwardly spreading plant habit; vigorous and sturdy growth habit.

Branching habit.—Freely branching habit with about four to six primary branches each with numerous secondary and tertiary lateral branches developing per plant.

Plant height.—About 88 cm.

Plant diameter (area of spread).—About 76 cm.

Lateral branch description:

Length.—About 87 cm.

Diameter, proximally.—About 1.5 cm.

Internode length.—About 1.2 cm.

Strength.—Strong.

Aspect.—About 45° from vertical.

Texture and luster.—Pubescent; woody and glabrous with age; matte.

Color, immature.—Close to 185A.

Color, mature.—Close to 177A.

Leaf description:

Arrangement.—Alternate; simple.

Length.—About 4.7 cm.

Width.—About 2.5 cm.

Shape.—Ovate.

Apex.—Acute.

Base.—Obtuse.

Margin.—Entire.

Texture and luster, upper surface.—Smooth, glabrous; dull.

Texture and luster, lower surface.—Smooth, glabrous; semi-glossy.

Venation pattern.—Pinnate.

Color.—Developing leaves, upper surface: Close to 147A. Developing leaves, lower surface: Close to 146A. Fully expanded leaves, upper surface: Close to N186A tinged with close to 200A; venation, close to 183A. Fully expanded leaves, lower surface: Close to 146A; venation, close to 144A.

Petioles.—Length: About 2 mm. Diameter: About 1 mm. Texture and luster, upper and lower surfaces: Smooth, glabrous; matte. Color, upper and lower surfaces: Close to 183A.

Flower description:

Flower type, arrangement and habit.—Showy single ruffled flowers arranged in terminal panicles; freely flowering habit with usually about twelve flowers per inflorescence and numerous inflorescences developing during the flowering season; flowers face upright and outwardly; flowers not persistent.

Natural flowering season.—Plants of the new Crapemyrtle flower during the late summer and autumn in Fort Worth, Tex.

Fragrance.—None detected.

Inflorescence height.—About 8 cm.

Inflorescence diameter.—About 4.7 cm.

Flower length.—About 3.5 cm.

Flower diameter.—About 3.5 cm.

Flower depth.—About 1.5 cm.

Flower buds.—Length: About 7 mm. Diameter: About 7 mm. Shape: Obovate to spherical. Texture and luster: Smooth, glabrous; slightly glossy. Color: Proximally, close to 144A; distally and sutures, close to 59A to 59B.

Petals.—Quantity per flower and arrangement: Six arranged in a single whorl. Length: About 1 cm.

Width: About 1.1 cm. Shape: Roughly orbicular. Apex: Rounded, undulate, ruffled appearance. Base: Rounded. Margin: Entire, undulate, ruffled appearance. Texture and luster, upper and lower surfaces: Smooth, glabrous; soft; delicate; matte. Color: When opening, upper and lower surfaces: Close to NN155D. Fully opened, upper and lower surfaces: Close to NN155D; flower stalk, close to 59A to 59B; colors do not change with development.

Sepals.—Quantity per flower and arrangement: Six arranged in a single whorl. Length: About 7 mm. Width: About 5 mm. Shape: Deltoid. Apex: Acute. Base: Fused. Margin: Entire. Texture and luster, upper surface: Smooth, glabrous; semi-glossy. Texture and luster, lower surface: Smooth, glabrous; slightly glossy. Color: When opening and fully opened, upper surface: Close to 144B to 144C. When opening and fully opened, lower surface: Close to 144A; towards the margins, close to 59A to 59B.

Pedicels.—Length: About 8 mm. Diameter: About 1.5 mm. Strength: Strong, flexible. Aspect: About 45° from stem axis. Texture and luster: Smooth, glabrous; slightly glossy. Color: Proximally, close to 59A to 59B; distally, close to 144A.

Reproductive organs.—Androecium: Quantity per flower: About twelve. Filament length: About 7 mm. Filament color: Close to NN155B to NN155C.

Anther length: About 1.5 mm. Anther shape: Oblong. Anther color: Close to 7A. Amount of pollen: None observed. Gynoecium: Quantity per flower: One. Pistil length: About 1.5 cm. Style length: About 1.3 cm. Style color: Close to 58A to 58B. Stigma appearance: Spherical. Stigma color: Close to NN137A. Ovary: Close to 10B to 10C.

Fruits and seeds.—Fruit and seed development has not been observed on plants of the new Crapemyrtle.

Garden performance: Plants of the new Crapemyrtle have been observed to have good garden performance and to tolerate rain, wind and temperatures ranging from about -10° C. to about 38° C. and to be cold hardy to USDA Hardiness Zone 6.

Pathogen & pest resistance: Plants of the new Crapemyrtle have been observed to be resistant to leaf spot (*Cercospora lythracearum*) and powdery mildew (*Erysiphe lagerstroemia*). Plants of the new Crapemyrtle have not been observed to be resistant to pests and other pathogens common to Crapemyrtle plants.

It is claimed:

1. A new and distinct Crapemyrtle plant named 'JM6' as illustrated and described.

* * * * *



