



US00PP32787P2

(12) **United States Plant Patent**  
**Smith**

(10) **Patent No.:** **US PP32,787 P2**  
(45) **Date of Patent:** **Feb. 2, 2021**

(54) **CHRYSANTHEMUM PLANT NAMED**  
**'CIFZ0075'**

(50) Latin Name: *Chrysanthemum x morifolium*  
Varietal Denomination: **CIFZ0075**

(71) Applicant: **SYNGENTA CROP PROTECTION**  
**AG, Basel (CH)**

(72) Inventor: **Mark A. Smith**, Alva, FL (US)

(73) Assignee: **Syngenta Crop Protection AG**, Basel  
(CH)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/873,305**

(22) Filed: **Mar. 16, 2020**

(51) **Int. Cl.**  
**A01H 5/02** (2018.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./289**

(58) **Field of Classification Search**  
USPC ..... Plt./289  
See application file for complete search history.

*Primary Examiner* — Annette H Para

(74) *Attorney, Agent, or Firm* — Dale Skalla

(57) **ABSTRACT**

A new garden type *Chrysanthemum* plant named  
'CIFZ0075' particularly distinguished by its medium to  
large size plant with round mound plant habit, a medium size  
decorative flower (2.0 inch), yellow flower color and a  
natural season response of early September.

**1 Drawing Sheet**

**1**

Latin name of the genus and species of the plant claimed:  
*Chrysanthemum x morifolium*.

Varietal denomination: 'CIFZ0075'.

**BACKGROUND OF THE NEW PLANT**

The present invention comprises a new *Chrysanthemum*,  
botanically known as *Chrysanthemum x morifolium*, and  
hereinafter referred to by the variety name 'CIFZ0075'.

'CIFZ0075' is a product of a planned breeding program.  
The new cultivar has a medium to large size plant with round  
mound plant habit, a medium size decorative flower (2.0  
inch), yellow flower color and a natural season response of  
early September.

'CIFZ0075' originated from a cross made in January 2014  
in Tenjo, Colombia. Seeds were sown in June 2015 in  
Gilroy, Calif. and 'CIFZ0075' was selected from the result-  
ing population on September 28 in Monroeville, N.J.

The female parent was the variety 'Syelect Ambe', U.S.  
Plant Pat. No. 22,770.

**TABLE 1**

| Characteristics of the female parent compared to 'CIFZ0075': |               |                |
|--|---------------|----------------|
| Trait  | 'CIFZ0075'    | 'Syelect Ambe' |
| Natural response:  | 1 week faster | 1 week slower  |
| Blackcloth response:   | Similar       | Similar        |
| Flower size:   | Similar       | Similar        |
| Flower color:  | Clear yellow  | Yellowy bronze |
| Flower type:   | Similar       | Similar        |
| Plant size:  | Similar       | Similar        |
| Plant habit:   | Similar       | Similar        |
| Flower longevity:  | ½ week less   | ½ week more    |

The male parent was 'CIFZ0013', U.S. Plant Pat. No.  
25,878.

**2**

**TABLE 2**

| Characteristics of the male parent compared to 'CIFZ0075': |                     |                    |
|--|---------------------|--------------------|
|  | 'CIFZ0075'          | 'CIFZ0013'         |
| Natural response - North US:                               | Similar             | Similar            |
| Natural response - South US:                               | 1 week faster       | 1 week slower      |
| Blackcloth response:                                       | ½ week faster       | ½ week slower      |
| Flower size:   | Similar/bit smaller | Similar/bit larger |
| Flower color:  | Similar             | Similar            |
| Flower type:   | Similar             | Similar            |
| Plant size:  | Larger              | Smaller            |
| Plant habit:   | Similar             | Similar            |
| Flower longevity:  | ½ week less         | ½ week more        |

The first act of asexual reproduction of 'CIFZ0075' was  
accomplished when vegetative stem cuttings were propa-  
gated from the initial selection in January 2016 in Mebane,  
N.C.

**BRIEF SUMMARY OF INVENTION**

Horticultural examination of plants grown from cuttings  
of the plant initiated in January 2016 and continuing there-  
after has demonstrated that the combination of characteris-  
tics as herein disclosed for 'CIFZ0075' are firmly fixed and  
are retained through successive generations of asexual  
reproduction.

'CIFZ0075' has not been observed under all possible  
environmental conditions. The phenotype may vary signifi-  
cantly with variations in environment such as temperature,  
light intensity and day length.

A Plant Breeder's Right for this cultivar has not yet been  
applied for. 'CIFZ0075' has not been made publicly avail-  
able prior to the effective filing date of this application,  
notwithstanding any disclosure that may have been made  
less than one year prior to the effective filing date of this  
application by the inventor or another who obtained  
'CIFZ0075' directly from the inventor.

The following traits have been repeatedly observed and are determined to be basic characteristics of the new variety. The combination of these characteristics distinguishes this *Chrysanthemum* as a new and distinct variety.

## BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographic drawings show typical flower and foliage characteristics of 'CIFZ0075' with colors being as true as possible with an illustration of this type.

The photographic drawing shows in FIG. 1 a close view of flower of the new variety and in

FIG. 2 a flowering plant from an outdoor trial.

The aforementioned photographs, FIG. 1, as well as FIG. 2, were taken in September 2019, both showing a plant from the same black cloth indoor trial in Gilroy, Calif. These plants were about 10 weeks of age. One rooted cutting per pot had been planted in a one gallon pot, not pinched in week 24, and black clothed from week 25. Plants started flowering the end of August.

## DETAILED BOTANICAL DESCRIPTION

The measurements were taken in Gilroy, Calif. in September 2019 on the plants from the aforementioned trial.

Color references are made to The Royal Horticultural Society Colour Chart (R.H.S.) 2015.

## DIFFERENCES BETWEEN THE NEW VARIETY 'CIFZ0075' AND TWO MOST SIMILAR VARIETIES

TABLE 3

| Trait                | 'CIFZ0075'    | 'Sunrise Yellow',<br>Not patented |
|----------------------|---------------|-----------------------------------|
| Natural response:    | ½ week faster | ½ week slower                     |
| Blackcloth response: | ½ week faster | ½ week slower                     |
| Flower size:         | Similar       | Similar                           |
| Flower color:        | Similar       | Similar                           |
| Flower type:         | Similar       | Similar                           |
| Plant size:          | Larger        | Smaller                           |
| Plant habit:         | Similar       | Similar                           |
| Plant strength:      | Softer        | Sturdier                          |

TABLE 4

| Trait                     | 'CIFZ0075'          | 'Sinelli Yellow', U.S.<br>Patent No. PP23,763 |
|---------------------------|---------------------|---|
| Natural response - North: | Similar             | Similar                                       |
| Natural response - South: | 1 week faster       | 1 week slower                                 |
| Blackcloth response:      | Similar             | Similar                                       |
| Flower size:              | Similar/bit larger  | Similar/bit smaller                           |
| Flower color:             | Similar             | Similar                                       |
| Flower type:              | Similar/less bracts | Similar/more bracts                           |
| Plant size:               | Larger              | Smaller                                       |
| Plant habit:              | Similar             | Similar                                       |
| Plant strength:           | Softer              | Sturdier                                      |

## Plant:

*Form, growth and habit.*—Herbaceous garden-type, stems. Upright and outwardly spreading, freely branching, strong and moderately vigorous growth habit.

*Plant height (above soil).*—15-17 cm.

*Plant height (inflorescence included).*—18-20 cm.

*Plant width.*—37-38 cm.

## Roots:

5 *Number of days to initiate roots.*—About 4 days at about 22° C.

*Number of days to produce a rooted cutting.*—14-16 days at 22° C.

*Type.*—Fine, fibrous, free branching.

10 *Color.*—RHS N155B.

## Foliage:

*Arrangement.*—Alternate.

*Immature leaf, color upper surface.*—RHS 147A.

*Immature leaf, color lower surface.*—Between RHS 137B and RHS 137C.

*Mature leaf, color, upper surface.*—RHS 137A.

*Mature leaf, color lower surface.*—RHS 137C.

*Length.*—4.3 cm.

*Width.*—3.5 cm.

*Shape.*—Ovate, with distinct lobes.

*Base shape.*—Attenuate.

*Apex shape.*—Mucronate.

*Margin.*—5-lobed (palmate), edge dentate.

25 *Number of margin indentations.*—Few.

*Depth of margin indentations.*—Shallow.

*Leaf length terminal lobe relative to total leaf length.*—Terminal lobe is ⅓ the length of the total leaf=Medium.

30 *Leaf depth lower lateral sinus.*—Shallow.

*Texture, upper surface.*—Bifid hairs.

*Texture, lower surface.*—Bifid hairs.

*Color of veins, upper surface.*—RHS 138B.

35 *Color of veins, lower surface.*—RHS 138A.

*Pattern of veining.*—Palmate.

*Petiole color.*—RHS 138A.

*Petiole length.*—1.5-2.0 cm.

*Diameter.*—0.2-0.3 cm.

40 *Texture.*—Bifid hairs.

*Presence of stipules.*—Yes.

*If present, size of stipules.*—Small.

## Stem:

*Quantity of main branches per plant.*—6-8.

*Color of stem.*—RHS 138A.

*Length of stem.*—8.0-10.0 cm.

*Diameter.*—0.5 cm.

*Length of internodes.*—1.5-3.0 cm.

*Texture.*—Bifid hairs.

50 *Color of peduncle.*—RHS 138A.

*Length of peduncle.*—4.5-5.0 cm.

*Peduncle diameter.*—0.1-0.3 cm.

*Texture.*—Bifid hairs.

## Inflorescence:

55 *Type.*—Compositae, solitary, decorative type inflorescences borne terminally above foliage, ligulate ray florets arranged acropetally on a capitulum giving a double flower.

*Quantity of short days to flowering (response time).*—Approximately 5 weeks.

*Natural season flowering.*—Early September.

*Quantity of inflorescences per plant.*—150-160 with many small buds developing.

*Lastingness of individual blooms on the plant.*—About five weeks from first color.

*Fragrance.*—Slightly spicy.

- Bud (when showing color):  
*Color*.—RHS 1B, Tips are RHS 4A.  
*Length*.—1.1 cm.  
*Width*.—1.0 cm.  
*Shape*.—Oblate.
- Immature inflorescence (at moment of opening):  
*Diameter*.—2.0 cm.  
*Color of ray florets, upper surface*.—RHS 7A.  
*Color of ray florets, lower surface*.—RHS 7C.
- Mature inflorescence:  
*Diameter*.—4.0 cm.  
*Depth*.—1.3 cm.  
*Total diameter of disc*.—0.3 cm to no disc present.  
*Receptacle color*.—RHS 9A.  
*Receptacle height*.—0.4 cm.  
*Receptacle diameter*.—1.0 cm.  
*Length of corolla tube*.—Medium.
- Ray florets:  
*Average quantity of florets*.—150-160 in several whorls.  
*Color of florets, upper surface (abaxial)*.—RHS 9A.  
*Color inner surface (adaxial)*.—RHS 9C.  
*Length*.—1.8 cm.  
*Width/diameter*.—0.5 cm.  
*Shape*.—Elliptical.  
*Apex shape*.—Mostly obtuse and dentate.  
*Base shape*.—Tube.  
*Margin*.—Entire. Small incisions may be present at tip.  
*Margin*.—Type of rolling Weakly involute.  
*Texture, upper surface*.—Papillate.  
*Lower surface*.—Papillate.  
*Ribs present*.—Yes.  
*Number of keels*.—2.
- Disc florets:  
*Number of disc florets*.—Up to 200, may not be present in all flowers.  
*Width*.—Up to 0.5 cm.  
*Length*.—3.0 cm.  
*Color*.—RHS 1B; 144B at base.
- Inflorescence (at moment of senescence):  
*Color of ray florets, upper surface*.—RHS 9A.  
*Color of ray florets, lower surface*.—RHS 9C.

- Phyllaries:  
*Quantity*.—20-25.  
*Color, upper surface*.—RHS 143C.  
*Color, lower surface*.—RHS 143A.  
5 *Length*.—0.6 cm.  
*Width*.—0.1-0.2 cm.  
*Shape*.—Lanceolate.  
*Apex shape*.—Acute.  
*Base*.—Fused.  
10 *Margins*.—Entire.  
*Texture, upper surface*.—Glabrous.  
*Texture, lower surface*.—Canescent.
- Reproductive organs:  
*Pistil*.—One.  
15 *Length*.—0.5 cm.  
*Style color*.—RHS 150B.  
*Style length*.—0.3 cm.  
*Stigma color*.—RHS 7D.  
*Stigma shape*.—Bi-parted.  
20 *Ovary color*.—RHS 145C.  
*Ovary length*.—0.1-0.15 cm.  
*Ovary width*.—0.1 cm.
- Androecium:  
*Stamens*.—1, found on only disc florets.  
25 *Color of filaments*.—RHS 2B.  
*Length filaments*.—0.1 cm.  
*Anther color*.—RHS 13A.  
*Anther length*.—0.1 cm.  
*Anther shape*.—Oval to club shaped.  
30 *Color of pollen*.—RHS 13A.  
*Pollen amount*.—Moderate.  
*Fertility/seed set*.—Has not been determined to date.  
*Disease/pest resistance*.—Has not been determined to date.  
35 *Hardiness*.—Has not been determined to date.
- What is claimed is:  
1. A new and distinct variety of *Chrysanthemum* plant named 'CIFZ0075' substantially as illustrated and described herein.  
40

\* \* \* \* \*



FIG. 1




FIG. 2