



(12) **United States Plant Patent**
Koot

(10) **Patent No.:** **US PP33,744 P2**
(45) **Date of Patent:** **Dec. 14, 2021**

(54) **PETUNIA PLANT NAMED ‘DOPETSUSPAPU’**

(50) Latin Name: *Petunia X hybrida*
Varietal Denomination: **Dopetsuspapu**

(71) Applicant: **DUMMEN GROUP B.V.**, De Lier
(NL)

(72) Inventor: **Arjan Koot**, Oeffelt (NL)

(73) Assignee: **Dümmen Group B.V.**, De Lier (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **17/317,678**

(22) Filed: **May 11, 2021**

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/82 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./356.18**
CPC **A01H 6/824** (2018.05)

(58) **Field of Classification Search**

USPC Plt./356.18
CPC A01H 5/02
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

<https://redbudcreekfarm.com/the-farm-report/specialty-petunias-that-we-are-growing-for-2020>; Mar. 28, 2021; 1 page.*

* cited by examiner

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — C. Anne Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Petunia* plant named ‘Dopet-suspapu’, characterized by its relatively compact, upright and mounding to hanging plant habit; moderately vigorous growth habit; freely branching habit; early and freely flowering habit; large reddish purple-colored flowers with random white-colored spots; and good garden performance.

2 Drawing Sheets

1

Botanical designation: *Petunia X hybrida*.
Cultivar denomination: ‘DOPETSUSPAPU’.

**STATEMENT REGARDING PRIOR
DISCLOSURES BY INVENTOR/APPLICANT &
ASSIGNEE**

An European Community Plant Breeder’s Rights application for the instant plant was filed by the Applicant/Assignee, Dümmen Group B.V. of De Lier, The Netherlands on Aug. 19, 2020, application number 2020/1953. Foreign priority is not claimed to this application.

The Inventor and Applicant/Assignee assert that no publications nor advertisements relating to sales, offers for sale or public distribution occurred more than one year prior to the effective filing date of this application. Any information about the claimed plant would have been obtained from a direct or indirect disclosure from the Inventor and/or Applicant/Assignee. Inventor and Applicant/Assignee claim a prior art exception under 35 U.S.C. 102(b)(1) for disclosure and/or sales prior to the filing date but less than one year prior to the effective filing date.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Petunia* plant, botanically known as *Petunia X hybrida* and hereinafter referred to by the name ‘Dopetsuspapu’.

The new *Petunia* plant is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. The objective of the breeding program is to create new compact, freely branching and early-flowering *Petunia* plants with numerous attractive flowers.

2

The new *Petunia* plant originated from a cross-pollination made by the Inventor in July, 2017 in Rheinberg, Germany of a proprietary selection of *Petunia X hybrida* identified as code number TT-0955, not patented, as the female, or seed, parent with a proprietary selection of *Petunia X hybrida* identified as code number TT16-107103-005, not patented, as the male, or pollen, parent. The new *Petunia* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rheinberg, Germany in May, 2020.

Asexual reproduction of the new *Petunia* plant by terminal vegetative cuttings in a controlled greenhouse environment in Rheinberg, Germany since June, 2020 has shown that the unique features of this new *Petunia* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Petunia* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Dopet-suspapu’. These characteristics in combination distinguish ‘Dopetsuspapu’ as a new and distinct *Petunia* plant:

1. Relatively compact, upright and mounding to hanging plant habit.
2. Moderately vigorous growth habit.
3. Freely branching habit.

4. Early and freely flowering habit.
5. Large reddish purple-colored flowers with random white-colored spots.
6. Good garden performance.

Plants of the new *Petunia* can be compared to plants of the female parent selection. In side-by-side comparisons, plants of the new *Petunia* differ primarily from plants of the female parent selection in flower color as flowers of plants of the new *Petunia* are reddish purple in color with random white-colored spots whereas flowers of plants of the female parent selection are pink in color.

Plants of the new *Petunia* can be compared to plants of the male parent selection. In side-by-side comparisons, plants of the new *Petunia* differ primarily from plants of the male parent selection in flower color as flowers of plants of the new *Petunia* are reddish purple in color with random white-colored spots whereas flowers of plants of the male parent selection are reddish purple in color without spots.

Plants of the new *Petunia* can be compared to plants of *Petunia* X *hybrida* 'KLEPH15313', disclosed in U.S. Plant Pat. No. 27,237. In side-by-side comparisons, plants of the new *Petunia* and 'KLEPH15313' differ primarily in the following characteristics:

1. Plants of the new *Petunia* are not as broad as plants of 'KLEPH15313'.
2. Plants of the new *Petunia* have larger flowers than plants of 'KLEPH15313'.
3. Flowers of plants of the new *Petunia* are reddish purple in color with random white-colored spots whereas flowers of plants of 'KLEPH15313' are violet in color with random white-colored spots.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Petunia* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Petunia* plant.

The photograph on the first sheet (FIG. 1) is a side perspective view of a typical flowering plant of 'Dopetsuspapu' grown in a container.

The photograph on the second sheet (FIG. 2) is a close-up view of a typical flower of 'Dopetsuspapu'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown during the spring in 22-cm containers in a glass-covered greenhouse in Rheinberg, Germany and under cultural practices typical of commercial *Petunia* production. During the production of the plants, day and night temperatures averaged 18° C. and light levels averaged 4,500 lux. Plants were twelve weeks old when the photographs were taken and 25 weeks old when the description was taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, Fifth Edition, 2007, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Petunia* X *hybrida* 'Dopetsuspapu'.
Parentage:

Female, or seed, parent.—Proprietary selection of *Petunia* X *hybrida* identified as code number TT-0955, not patented.

Male, or pollen, parent.—Proprietary selection of *Petunia* X *hybrida* identified as code number TT16-107103-005, not patented.

Propagation:

Type.—By terminal vegetative cuttings.

Time to initiate roots, summer.—About five days at temperatures about 20° C.

Time to initiate roots, winter.—About seven days at temperatures about 20° C.

Time to produce a rooted young plant, summer.—About three weeks at temperatures about 20° C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures about 20° C.

Root description.—Fine, fibrous; close to 155B in color, actual color of the roots is dependent on substrate composition, water quality, fertilizers, substrate temperature and age of roots.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Relatively compact, upright and mounding to hanging plant habit; freely branching habit with about five primary lateral branches each with about twelve secondary branches developing after pinching; moderately vigorous growth habit; moderate growth rate.

Plant height, soil level to top of foliar plane.—About 23 cm.

Plant height, soil level to top of floral plane.—About 25 cm.

Plant diameter.—About 73 cm.

Lateral branch description:

Length.—About 60 cm.

Diameter.—About 4 mm.

Internode length.—About 1.5 cm.

Strength.—Moderately strong.

Aspect.—Initially upright to outwardly spreading.

Texture and luster.—Pubescent; semi-glossy.

Color, developing and developed.—Close to 144A.

Leaf description:

Arrangement.—Before flowering, alternate; after flowering, opposite; simple.

Length.—About 2.8 cm.

Width.—About 1.4 cm.

Shape.—Spatulate.

Apex.—Obtuse.

Base.—Attenuate.

Margin.—Entire.

Texture and luster, upper and lower surfaces.—Pubescent; leathery; semi-glossy.

Venation pattern.—Pinnate; arcuate.

Color.—Developing and fully expanded leaves, upper surface: Close to 144A; venation, close to 144B. Fully expanded leaves, lower surface: Close to 145A; venation, close to 144B.

Petioles.—Length: About 3 mm. Diameter: About 1.2 mm. Strength: Moderately strong; firm. Texture and luster, upper and lower surfaces: Smooth, glabrous; matte. Color, upper and lower surfaces: Close to 144B.

Flower description:

Flower type and flowering habit.—Single salverform flowers arising from leaf axils; freely flowering habit with usually about 450 flowers and flower buds developing per plant; flowers face mostly upright to outwardly. 5

Fragrance.—None detected.

Natural flowering season.—Plants flower continuously during the spring and summer in Germany; early flowering habit, plants typically beginning flowering about nine weeks after planting. 10

Flower longevity.—Individual flowers last about two to three days on the plant; flowers persistent.

Flower buds.—Length: About 4.2 cm. Diameter: About 5.5 mm. Shape: Ovoid. Texture and luster: Rippled; semi-glossy. Color: Close to 83B and 145B. 15

Flower diameter.—About 6.5 cm.

Flower depth (height).—About 6.2 cm.

Flower throat diameter.—About 1.1 cm.

Flower tube length.—About 3.3 cm. 20

Flower tube diameter, proximally.—About 2 mm.

Corolla.—Arrangement: Five petals fused at the base and opening into a flared trumpet. Petal lobe length (from throat): About 3.1 cm. Petal lobe width: About 3.3 cm. Petal shape: Roughly spatulate. Petal apex: Obtuse. Petal margin: Entire; slightly undulate. Petal texture and luster, upper and lower surfaces: Rippled, glabrous; semi-glossy. Throat texture and luster: Rippled, glabrous; semi-glossy. Tube texture and luster: Rippled, pubescent; semi-glossy. Color: Petal lobe, when opening, upper surface: Close to 79B with random spots, close to NN155D. Petal lobe, when opening, lower surface: Close to 79C with random spots, close to NN155D. Petal lobe, fully opened, upper surface: Close to N78A with random spots, close to NN155D; venation, close to N79A; main color becoming closer to N88A with subsequent development. Petal lobe, fully opened, lower surface: Close to 77A with random spots, close to NN155D; venation, close to N79A; main color becoming closer to N88B with subsequent development. Flower throat: Close to 83A; venation, close to N79A. Flower tube: Close to 83A; venation, close to N79A. 25 30 35 40

Sepals.—Arrangement: Five sepals fused at the base forming a tubular star-shaped calyx. Length: About 1.6 cm. Diameter: About 2 mm. Shape: Oblong. Apex: Rounded. Base: Decurrent. Margin: Entire. Texture and luster, upper and lower surfaces: Smooth, glabrous; semi-glossy. Color: When opening and fully opened, upper surface: Close to 146A. When opening and fully opened, lower surface: Close to 146B.

Peduncles.—Length: About 2.5 cm. Diameter: About 1 mm. Strength: Moderately strong. Texture and luster: Smooth, glabrous; semi-glossy. Color: Close to 144A.

Reproductive organs.—Stamens: Quantity per flower: Five. Filament length: About 2.1 cm. Filament color: Close to 157D. Anther length: About 1.5 mm. Anther shape: Ovate. Anther color: Close to 158D. Pollen amount: Abundant. Pollen color: Close to 158B. Pistils: Quantity per flower: One. Pistil length: About 2.6 cm. Style length: About 2.1 cm. Style color: Close to 144D. Stigma diameter: About 2 mm. Stigma shape: Rounded. Stigma color: Close to 144C. Ovary color: Close to 144B. Fruits: Quantity produced per plant: About 90 during the flowering season. Length: About 6 mm. Diameter: About 5 mm. Texture: Smooth, glabrous. Color: Close to 161C. Seeds: Quantity per flower: About 150. Length: About 0.2 mm. Diameter: About 0.2 mm. Texture: Smooth, glabrous. Color: Close to 200C.

Garden performance: Plants of the new *Petunia* have been observed to have good garden performance and tolerate wind, rain, temperatures ranging from about 5° C. to about 40° C. and to be suitable for USDA Hardiness Zone 11.

Pathogen & pest resistance: To date, plants of the new *Petunia* have not been observed to be resistant to pathogens and pests common to *Petunia* plants.

It is claimed:

1. A new and distinct *Petunia* plant named 'Dopetsus-papu' as illustrated and described.

* * * * *




FIG. 1




FIG. 2