



US00PP33942P2

(12) **United States Plant Patent**
Lannes

(10) **Patent No.:** **US PP33,942 P2**
(45) **Date of Patent:** **Feb. 8, 2022**

(54) **MANDEVILLA PLANT NAMED**
‘LANNEWYORK’

(50) Latin Name: *Mandevilla sanderi*
Varietal Denomination: **Lannewyork**

(71) Applicant: **Robert Lannes**, Malause (FR)

(72) Inventor: **Robert Lannes**, Malause (FR)

(73) Assignee: **D.H.M. INNOVATION S.A.S.**,
Malause (FR)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **17/357,279**

(22) Filed: **Jun. 24, 2021**

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/08 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./232**
CPC *A01H 6/088* (2018.05)

(58) **Field of Classification Search**
USPC Plt./232
CPC A01H 5/02
See application file for complete search history.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — C. Anne Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Mandevilla* plant named
‘Lannewyork’, characterized by its compact, broadly
upright and vining plant habit; moderately vigorous growth
habit; freely branching habit; glossy dark green-colored
leaves; early and freely flowering habit; and large flowers
with red-colored petals and greenish yellow-colored throats.

2 Drawing Sheets

1

Botanical designation: *Mandevilla sanderi*.
Cultivar denomination: ‘LANNEWYORK’.

STATEMENT REGARDING PRIOR
DISCLOSURES BY THE
INVENTOR/APPLICANT & ASSIGNEE

An European Community Plant Breeder’s Rights appli-
cation for the instant plant was filed by the D.H.M. Inno-
vations S.A.S. of Malause, France on Nov. 30, 2020, appli-
cation number 2020/3092. Foreign priority is not claimed to
this application.

The Inventor/Applicant and Assignee assert that no pub-
lications nor advertisements relating to sales, offers for sale
or public distribution occurred more than one year prior to
the effective filing date of this application. Any information
about the claimed plant would have been obtained from a
direct or indirect disclosure from the Inventor/Applicant
and/or the Assignee. Inventor/Applicant and Assignee claim
a prior art exception under 35 U.S.C. 102(b)(1) for disclo-
sure and/or sales prior to the filing date but less than one year
prior to the effective filing date.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Mandevilla* plant, botanically known as *Mandevilla san-
deri* and hereinafter referred to by the name ‘Lannewyork’.

The new *Mandevilla* plant is a product of a planned
breeding program conducted by the Inventor in Malause,
France. The objective of the breeding program is to create
new freely branching *Mandevilla* plants with numerous
unique and attractive flowers.

The new *Mandevilla* plant originated from a cross-poll-
ination conducted by the Inventor in Malause, France in July,
2016 of a proprietary selection of *Mandevilla sanderi* iden-
tified as code number si 08-0426-044, not patented, as the

2

female, or seed, parent with a proprietary selection of
Mandevilla sanderi identified as code number si s346-0, not
patented, as the male, or pollen, parent. The new *Mandevilla*
plant was discovered and selected by the Inventor as a single
flowering plant from within the progeny of the stated
cross-pollination in a controlled greenhouse environment in
Malause, France in June, 2017.

Asexual reproduction of the new *Mandevilla* plant by
stem cuttings in a controlled greenhouse environment in
Malause, France since August, 2019 has shown that the
unique features of this new *Mandevilla* plant are stable and
reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Mandevilla* have not been observed
under all possible combinations of environmental conditions
and cultural practices. The phenotype may vary somewhat
with variations in environmental conditions such as tem-
perature and light intensity without, however, any variance
in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Lannewy-
ork’. These characteristics in combination distinguish
‘Lannewyork’ as a new and distinct *Mandevilla* plant:

1. Compact, broadly upright and vining plant habit.
2. Moderately vigorous growth habit.
3. Freely branching habit.
4. Glossy dark green-colored leaves.
5. Early and freely flowering habit.
6. Large flowers with red-colored petals and greenish
yellow-colored throats.

Plants of the new *Mandevilla* can be compared to plants
of the female parent selection. Plants of the new *Mandevilla*
differ primarily from plants of the female parent selection in
the following characteristics:

1. Plants of the new *Mandevilla* are more compact and have shorter internodes than plants of the female parent selection.
2. Plants of the new *Mandevilla* are more freely flowering than plants of the female parent selection.

Plants of the new *Mandevilla* can be compared to plants of the male parent selection. Plants of the new *Mandevilla* differ primarily from plants of the male parent selection in the following characteristics:

1. Plants of the new *Mandevilla* are more compact than plants of the male parent selection.
2. Flowers of plants of the new *Mandevilla* have red-colored petals whereas flowers of plants of the male parent selection have rose-colored petals.

Plants of the new *Mandevilla* can also be compared to plants of *Mandevilla* x *amabilis* X *Mandevilla hybrida* 'Lankentucky', disclosed in U.S. Plant Pat. No. 29,849. In side-by-side comparisons, plants of the new *Mandevilla* differ primarily from plants 'Lankentucky' in the following characteristics:

1. Plants of the new *Mandevilla* have larger and more rounded leaves than plants of 'Lankentucky'.
2. Flowers of plants of the new *Mandevilla* have red-colored petals whereas flowers of plants of 'Lankentucky' have brighter red-colored petals.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Mandevilla* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the actual colors of the new *Mandevilla* plant.

The photograph on the first sheet (FIG. 1) comprises a side perspective view of a typical flowering plant of 'Lannewyork' grown in a container.

The photograph on the second sheet (FIG. 2) is a close-up view of a typical flower of 'Lannewyork'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown during the autumn in 11-cm containers in a polyethylene-covered greenhouse in Malause, France and under cultural practices typical of commercial *Mandevilla* production. During the production of the plants, day temperatures ranged from 18° C. to 30° C. and night temperatures ranged from 15° C. to 18° C. Plants were ten months old when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2015 Edition, except where general terms of ordinary dictionary significance are used. Botanical classification: *Mandevilla sanderi* 'Lannewyork'. Parentage:

Female, or seed, parent.—Proprietary selection of *Mandevilla sanderi* identified as code number si 08-0426-044, not patented.

Male, or pollen, parent.—Proprietary selection of *Mandevilla sanderi* identified as code number si s346-0, not patented.

Propagation:

Type.—By vegetative stem cuttings.

Time to initiate roots, summer.—About two weeks at temperatures about 23° C. to 30° C.

Time to initiate roots, winter.—About three weeks at temperatures about 18° C. to 21° C.

Time to produce a rooted young plant, summer.—About five to six weeks at temperatures about 23° C. to 30° C.

Time to produce a rooted young plant, winter.—About six to eight weeks at temperatures about 18° C. to 21° C.

Root description.—Fine, fibrous; typically light yellowish white in color, actual color of the roots is dependent on substrate composition, water quality, fertilizers, substrate temperature and physiological age of roots.

Rooting habit.—Freely branching; medium density.

Plant description:

Plant and growth habit.—Compact, broadly upright and vining plant habit; overall plant shape, broadly oblong to obovate; moderately vigorous growth habit and moderate to slow growth rate.

Plant height, soil level to top of foliar plane.—About 31.8 cm.

Plant height, soil level to top of floral plane.—About 26.2 cm.

Plant diameter (spread).—About 18.9 cm.

Lateral branch description.—Branching habit: Freely branching habit, typically two primary lateral branches each with about two secondary lateral branches developing per plant. Length: About 25.3 cm. Diameter: About 2.5 mm. Internode length: About 3.6 cm. Aspect: Primary lateral branches, mostly erect; secondary lateral branches, about 20° from primary branch axis. Strength: Strong. Texture and luster: Smooth, glabrous; moderately glossy; becoming woody with development. Color, developing: Close to between 146A and 152A. Color, developed: Close to between 143C and 144A; when woody, close to N199B.

Leaf description:

Arrangement.—Opposite, simple.

Length.—About 9.3 cm.

Width.—About 6.5 cm.

Shape.—Broadly ovate to broadly oblong.

Apex.—Short apiculate to abruptly acute.

Base.—Truncate.

Margin.—Entire; slightly undulate.

Texture and luster, upper and lower surfaces.—Smooth, glabrous; moderately coriaceous; slightly to moderately glossy.

Venation pattern.—Pinnate.

Color.—Developing leaves, upper surface: Slightly darker and more intense than 144A. Developing leaves, lower surface: Close to 144A. Full expanded leaves, upper surface: Darker than between 139A and N189A; venation, close to 144A. Fully expanded leaves, lower surface: Close to between 146B and 147B; venation, close to between 146D and 147D.

Petioles.—Length: About 1.2 cm. Diameter: About 3 mm by 3 mm. Strength: Moderately strong. Texture and luster, upper surface: Smooth, glabrous; slightly glossy. Texture and luster, lower surface: Smooth,

glabrous; glossy. Color, upper surface: Close to 146C. Color, lower surface: Close to 146D.

Flower description:

Flower type and flowering habit.—Single salverform flowers arranged singly or in terminal or axillary cymes; flowers rounded star-shaped and face mostly outwardly to slightly upright; freely flowering habit with about twelve flower buds and flowers developing per plant.

Natural flowering season.—Plants flower continuously from spring into the autumn in France; early flowering habit, plants in full flower about six to seven months after planting.

Flower longevity on the plant.—About ten days; flowers not persistent.

Fragrance.—None detected.

Inflorescence height.—About 7.4 cm.

Inflorescence diameter.—About 8.2 cm.

Flower buds.—Length: About 6.1 cm. Diameter: About 1.4 cm. Shape: Narrowly oblanceolate. Texture and luster: Smooth, glabrous; slightly glossy. Color: Close to 54C; towards the base, close to 145C; tube, close to 145A.

Flowers.—Appearance: Flared trumpet, corolla fused and five-parted. Diameter: Large, about 9 cm. Depth (length): About 5.7 cm. Throat diameter: About 2 cm. Tube length: About 3.8 cm. Tube diameter: Distally, about 2.1 cm; proximally, about 3.5 mm.

Petals.—Quantity and arrangement: Five petals arranged in a single whorl; lower 50% portion of the petals are fused into a funnellform tube. Petal length: About 7.8 cm. Petal width: About 4.5 cm. Petal shape: Unequal spatulate. Petal apex: Bluntly acute. Petal margin: Entire; moderately and coarsely undulate. Petal texture and luster, upper surface: Smooth, glabrous; not rugose; velvety; matte. Petal texture and luster, lower surface: Smooth, glabrous; not rugose; slightly velvety; slightly glossy. Throat texture: Smooth, glabrous; velvety. Tube texture: Smooth, glabrous; slightly velvety. Color: Petal, when opening, upper surface: Close to N57B and N57C. Petal, when opening, lower surface: Close to N57C to N57D. Petal, fully opened, upper surface: Close to between 53C and N57A; towards the throat, close to 58B; venation, similar to lamina; color becoming closer to 53A to 53B with subsequent development. Petal, fully opened, lower surface: Close to 61C; venation, similar to lamina color; color becoming closer to 53D with subsequent development. Throat: Distally, close to 58C and proximally, close to 5A and 5B; towards the base, close to 146D; venation, similar to lamina colors. Tube: Distally,

close to 61C to 61D and proximally, close to 145C; towards the base, close to 145A; venation, similar to lamina colors.

Sepals.—Quantity and arrangement: Five sepals arranged in a single whorl. Calyx length: About 8 mm. Calyx diameter: About 5 cm. Sepal length: About 8 mm. Sepal width: About 2 mm. Sepal shape: Lanceolate. Sepal apex: Narrowly acuminate. Sepal base: Broadly cuneate and fused at the base. Sepal margin: Entire. Sepal texture and luster, upper surface: Smooth, glabrous; slightly glossy. Sepal texture and luster, lower surface: Smooth, glabrous; matte. Sepal color: When opening and fully opened, upper surface: Close to 144B; towards the margins, close to 145D. When opening and fully opened, lower surface: Close to 144A to 144B; towards the margins, close to 145D.

Peduncles.—Length: About 5.1 cm. Diameter: About 2 mm. Strength: Strong. Aspect: About 20° from lateral branch axis. Texture and luster: Smooth, glabrous; slightly to moderately glossy. Color: Close to 143B.

Pedicels.—Length: About 1.4 cm. Diameter: About 2.5 mm. Strength: Strong. Aspect: About 30° from peduncle axis. Texture and luster: Smooth, glabrous; moderately glossy. Color: Close to 144B.

Reproductive organs.—Stamens: Quantity and arrangement: Typically five; basifixed; anthers connivent. Filament length: About 2 mm. Filament color: Close to 150C to 150D. Anther size: About 1.25 mm by 8 mm. Anther shape: Narrowly oblong. Anther color: Close to 162C. Pollen amount: None observed. Pistils: Quantity: Typically one. Pistil length: About 2.4 cm. Style length: About 2.1 cm. Style color: Close to 145C to 145D. Stigma diameter: About 2 mm. Stigma shape: Club-shaped, pointed. Stigma color: Close to 146D. Ovary color: Close to 143B.

Seeds and fruits.—To date, seed and fruit production have not been observed on plants of the new *Mandevilla*.

Pathogen & pest resistance: To date, plants of the new *Mandevilla* have not been noted to be resistant to pathogens and pests common to *Mandevilla* plants.

Temperature tolerance: Plants of the new *Mandevilla* have been observed to tolerate temperatures of about 5° C. to 40° C. and to be suitable for USDA Hardiness Zones 9 through 13.

It is claimed:

1. A new and distinct *Mandevilla* plant named 'Lannewyork' as illustrated and described.

* * * * *



FIG. 1



FIG. 2