



(12) **United States Plant Patent**  
**Masor**

(10) **Patent No.:** **US PP34,037 P2**  
(45) **Date of Patent:** **Mar. 15, 2022**

(54) **CALIBRACHOA PLANT NAMED**  
**‘BALCABRIWII’**

(50) Latin Name: *Calibrachoa x hybrida*  
Varietal Denomination: **Balcabriwii**

(71) Applicant: **Ball Horticultural Company**, West  
Chicago, IL (US)

(72) Inventor: **Laura L. Masor**, Grover Beach, CA  
(US)

(73) Assignee: **Ball Horticultural Company**, West  
Chicago, IL (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **17/501,689**

(22) Filed: **Oct. 14, 2021**

(51) **Int. Cl.**  
**A01H 5/02** (2018.01)

**A01H 6/82** (2018.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./413**

(58) **Field of Classification Search**  
USPC ..... Plt./413  
See application file for complete search history.

*Primary Examiner* — Susan McCormick Ewoldt

(74) *Attorney, Agent, or Firm* — Audrey Charles

(57) **ABSTRACT**

A new and distinct cultivar of *Calibrachoa* plant named  
‘Balcabriwii’, characterized by its white-colored flowers,  
medium green-colored foliage, and moderately vigorous,  
mounded-trailing growth habit, is disclosed.

**1 Drawing Sheet**

**1**

Latin name of genus and species of plant claimed: *Cali-  
brachoa x hybrida*.

Variety denomination: ‘Balcabriwii’.

#### BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar  
of *Calibrachoa* plant botanically known as *Calibrachoa x*  
*hybrida* and hereinafter referred to by the cultivar name  
‘Balcabriwii’.

The new cultivar originated in a controlled breeding  
program in Guadalupe, Calif. during March 2019. The  
objective of the breeding program was the development of  
*Calibrachoa* cultivars with attractive novelty flower color-  
ation, and a moderately vigorous, mounded-trailing growth  
habit.

The new *Calibrachoa* cultivar is the result of cross-  
pollination. The female (seed) parent of the new cultivar is  
the proprietary *Calibrachoa x hybrida* breeding selection  
coded CAL-16400-01, not patented, characterized by its  
white-colored, medium bright-green colored foliage, and  
moderately vigorous, mounded-trailing growth habit. The  
male (pollen) parent of the new cultivar is the proprietary  
*Calibrachoa x hybrida* breeding selection coded CAL-  
16428-01, not patented, characterized by its cream-colored  
flowers, medium green-colored foliage, and moderately vig-  
orous, mounded-trailing growth habit. The new cultivar was  
selected as a single flowering plant within the progeny of the  
above stated cross-pollination during August 2019 in a  
controlled environment in Arroyo Grande, Calif.

Asexual reproduction of the new cultivar by terminal stem  
cuttings since August 2019 in Arroyo Grande, Calif. and  
West Chicago, Ill. has demonstrated that the new cultivar  
reproduces true-to-type with all of the characteristics, as  
herein described, firmly fixed and retained through succes-  
sive generations of such asexual propagation.

#### SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have  
been repeatedly observed and can be used to distinguish  
‘Balcabriwii’ as a new and distinct cultivar of *Calibrachoa*  
plant:

**2**

1. White-colored flowers;
2. Medium green-colored foliage; and
3. Moderately vigorous, mounded-trailing growth habit.

Plants of the new cultivar differ from plants of the female  
parent primarily in having more flowers per plant. Plants of  
the new cultivar differ from plants of the male parent  
primarily in having white-colored flowers.

Of the many commercially available *Calibrachoa* culti-  
vars, the most similar in comparison to the new cultivar is  
CABARET Bright White ‘Balcabriu’, not patented. How-  
ever, in side-by-side comparisons, plants of the new cultivar  
differ from plants of ‘Balcabriu’ in at least the following  
characteristics:

1. Plants of the new cultivar flower earlier than plants of  
‘Balcabriu’, and
2. Plants of the new cultivar has a more mounded-trailing  
growth habit than plants of ‘Balcabriu’.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it  
is reasonably possible to make the same in color illustrations  
of this type, typical flower and foliage characteristics of the  
new cultivar. Colors in the photographs may differ slightly  
from the color values cited in the detailed description, which  
accurately describes the colors of ‘Balcabriwii’. The plants  
were approximately 5-months old. The plants were grown in  
3-gallon containers for approximately 10 weeks in an out-  
door nursery in West Chicago, Ill. Plants were pinched twice  
prior to transplant.

FIG. 1 illustrates a side view of the overall growth and  
flowering habit of ‘Balcabriwii’.

FIG. 2 illustrates a close-up view of an individual flower  
of ‘Balcabriwii’.

#### DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible  
environmental conditions to date. Accordingly, it is possible

that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2015 edition, except where general color terms of ordinary significance are used. The color values were determined in August 2021 under natural light conditions in Naperville, Ill.

The following descriptions and measurements describe approximately 5-month-old plants produced from cuttings from stock plants and grown under conditions comparable to those used in commercial practice. The plants were grown in 3-gallon containers for approximately 10 weeks in an outdoor nursery in West Chicago, Ill. Plants were given two pinches prior to transplant. Prior to transplant plants were grown in a polycarbonate greenhouse in West Chicago, Ill. Greenhouse temperatures were maintained at approximately 70° F. to 85° F. (21° C. to 29° C.) during the day and approximately 60° F. to 70° F. (16° C. to 21° C.) during the night. Supplemental lighting was used during propagation stage. Measurements and numerical values represent averages of typical plants.

Botanical classification: *Calibrachoa* x *hybrida* 'Balcabriwii'.

Parentage:

*Female parent*.—Proprietary *Calibrachoa* x *hybrida* breeding selection coded CAL-16400-01, not patented.

*Male parent*.—Proprietary *Calibrachoa* x *hybrida* breeding selection coded CAL-16428-01, not patented.

Propagation:

*Type cutting*.—Terminal stem.

*Time to initiate roots*.—Approximately 6 to 8 days.

*Time to produce a rooted cutting*.—Approximately 21 to 28 days.

*Root description*.—Fibrous.

*Rooting habit*.—Freely branching.

Plant description:

*Commercial crop time*.—Approximately 6 to 9 weeks from a rooted cutting to finish in a 10 cm pot.

*Growth habit and general appearance*.—Moderately vigorous, mounded-trailing.

*Size*.—Height from soil level to top of plant plane: Approximately 21.5 cm. Width: Approximately 66.0 cm.

*Branching habit*.—Freely branching, pinching enhances basal branching. Quantity of main branches per plant: Approximately 9.

*Branch*.—Strength: Moderate. Length: Approximately 32.0 cm. Diameter: Approximately 2.0 mm to 3.0 mm. Length of central internode: Approximately 1.7 cm. Texture: Densely pubescent with a mixture of non-glandular and glandular hairs. Gland color: Colorless, transparent. Color of young stems: 146C. Color of mature stems: 146D with age becoming woody 199C.

Foliage description:

*General description*.—Fragrance: None detected. Form: Simple. Arrangement on flowering stem: Opposite.

*Leaves*.—Aspect: At acute angle to stem with tip turning downward. Shape: Elliptic. Margin: Entire. Apex: Broadly acute. Base: Attenuate. Venation pat-

tern: Pinnate. Length of mature leaf: Approximately 3.1 cm. Width of mature leaf: Approximately 1.0 cm. Texture of upper and lower surfaces: Densely glandular pubescent. Gland color: Colorless, transparent. Color of upper surface of young and mature foliage: 137B with midvein of 146C. Color of lower surface of young and mature foliage: Closest to 138B with midvein of 146D.

*Petiole*.—Length: Approximately 3.0 mm. Diameter: Approximately 2.0 mm. Texture: Densely glandular pubescent. Gland color: Colorless, transparent. Color: 146D.

Flowering description:

*Flowering habit*.—'Balcabriwii' is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

*Lastingness of individual flower on the plant*.—Approximately 5 to 7 days.

Flower description:

*General description*.—Type: Single, salverform. Quantity per plant: Approximately 65. Fragrance: None detected.

*Bud*.—Rate of opening: Generally takes 3 to 4 days for bud to progress from first color to fully open flower. Quantity per plant: Approximately 20.

*Bud just before opening*.—Shape: Oblong. Length: Approximately 1.8 cm. Diameter: Approximately 5.0 mm. Texture: Densely glandular pubescent. Gland color: Colorless, transparent. Color: 150D with venation of 145A.

*Corolla*.—Diameter: Approximately 3.0 cm.

*Petals*.—Quantity: 5, fused to form a tube. Shape: Obovate. Margin: Entire. Apex: Obtuse to shallowly notched. Length from tube: Approximately 1.2 cm. Length of free portion: Approximately 5.0 mm. Width: Approximately 1.2 cm. Texture of upper surface: Glabrous. Texture of lower surface: Densely glandular pubescent. Gland color: Colorless, transparent. Color of upper surface when first and fully open: NN155D with faint midveins of 146D. Color of lower surface when first and fully open: NN155D with midveins of 145A.

*Corolla tube*.—Length: Approximately 1.8 cm. Diameter at distal end: Approximately 6.0 mm. Diameter at proximal end: Approximately 1.0 mm. Texture of inner surface: Glabrous. Texture of outer surface: Densely glandular pubescent. Gland color: Colorless, transparent. Color of inner surface: 154C with venation of 145B. Color of outer surface: 154D with venation of 145A.

*Sepals*.—Quantity per flower: 5, fused along lower half. Shape: Lanceolate. Apex: Acute. Length: Approximately 1.1 cm. Width: Approximately 3.0 mm. Texture of upper and lower surfaces: Densely glandular pubescent. Gland color: Colorless, transparent. Color of upper surface: 137A. Color of lower surface: 137B with base of 146D.

*Peduncle*.—Strength: Strong, flexible. Aspect: Acute angle to stem. Length: Approximately 1.5 cm to 2.0 cm. Diameter: Approximately 1.0 mm. Texture: Densely glandular pubescent. Gland color: Colorless, transparent. Color: 146D.

*Reproductive organs*.—Androecium: Stamen quantity: 5, basifixed. Stamen length: Approximately 1.1 cm.

Filament length of fixed portion: Approximately 5.0 mm. Filament color: 150D. Anther shape: Bilobed, ovoid. Anther length: Approximately 1.0 mm. Anther color: 8C. Pollen amount: Abundant. Pollen color: 8D. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 1.0 cm. Stigma shape: Funnel. Stigma length: Less than 1.0 mm. Stigma color: 146D. Style length: Approximately 8.0 mm. Style color: 145D. Ovary diameter: Approximately 2.0 mm. Ovary color: 145B.

Seed and fruit production: Neither seed nor fruit production has been observed.  
Disease and pest resistance: Resistance to pathogens and pests common to *Calibrachoa* has not been observed.

What is claimed is:  
1. A new and distinct cultivar of *Calibrachoa* plant named ‘Balcabriwii’, substantially as herein illustrated and described.

\* \* \* \* \*



FIG. 1




FIG. 2