



(12) **United States Plant Patent**
van Sambeek

(10) **Patent No.:** **US PP34,301 P2**
(45) **Date of Patent:** **Jun. 7, 2022**

(54) **ERYSIMUM PLANT NAMED ‘DOERYWAPE’**

(50) Latin Name: *Erysimum hybrida*
Varietal Denomination: **Doerywape**

(71) Applicant: **DUMMEN GROUP B.V.**, De Lier
(NL)

(72) Inventor: **Ellen van Sambeek**, Oegstgeest (NL)

(73) Assignee: **Dümmen Group B.V.**, De Lier (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **17/499,077**

(22) Filed: **Oct. 12, 2021**

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/20 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./263.1**
CPC **A01H 6/20** (2018.05)

(58) **Field of Classification Search**

USPC Plt./263.1
CPC A01H 5/02
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

PLUTO UPOVROM Citation for ‘Doerywape’ as per QZ PBR
20211409; Aug. 15, 2021; 1 page.*

* cited by examiner

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — C. Anne Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Erysimum* plant named
‘Doerywape’, characterized by its compact, upright and
mounding plant habit; moderately vigorous growth habit;
freely branching habit; dense and bushy habit; freely and
continuous flowering habit; long flowering period; large
white-colored flowers; tolerance to low and high tempera-
tures and good garden performance.

1 Drawing Sheet

1

Botanical designation: *Erysimum hybrida*.
Cultivar denomination: ‘DOERYWAPE’.

STATEMENT REGARDING PRIOR DISCLOSURES BY INVENTOR & APPLICANT/ASSIGNEE

An European Community Plant Breeder’s Rights appli-
cation for the instant plant was filed by the Applicant/
Assignee, Dümmen Group B.V. of De Lier, The Netherlands
on Jun. 1, 2021, application number 2021/1409. Foreign
priority is not claimed to this application.

The Inventor and Applicant/Assignee assert that no pub-
lications nor advertisements relating to sales, offers for sale
or public distribution occurred more than one year prior to
the effective filing date of this application. Any information
about the claimed plant would have been obtained from a
direct or indirect disclosure from the Inventor and/or Appli-
cant/Assignee. Inventor and Applicant/Assignee claim a
prior art exception under 35 U.S.C. 102(b)(1) for disclosure
and/or sales prior to the filing date but less than one year
prior to the effective filing date.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Erysimum* plant, botanically known as *Erysimum hybrida*,
commonly referred to as Wallflower and hereinafter referred
to by the name ‘Doerywape’.

The new *Erysimum* plant is a product of a planned
breeding program conducted by the Inventor in Aalsmeer,

2

The Netherlands. The objective of the breeding program was
to develop new compact *Erysimum* plants with numerous
large and attractive flowers.

The new *Erysimum* plant originated from a cross-polli-
nation in February, 2017 in Aalsmeer, The Netherlands of a
proprietary selection of *Erysimum hybrida* identified as code
number ER16-000001-023, not patented, as the female, or
seed, parent, with a proprietary selection of *Erysimum*
hybrida identified as code number ER16-000001-028, not
patented, as the male, or pollen, parent. The new *Erysimum*
plant was discovered and selected by the Inventor as a single
flowering plant within the progeny of the stated cross-
pollination in a controlled greenhouse environment in Aals-
meer, The Netherlands in April, 2018. Asexual reproduction
of the new *Erysimum* plant by vegetative terminal cuttings
in a controlled greenhouse environment in Aalsmeer, The
Netherlands since May, 2018 has shown that the unique
features of this new *Erysimum* plant are stable and repro-
duced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Erysimum* have not been observed under
all possible combinations of environmental conditions and
cultural practices. The phenotype may vary somewhat with
variations in environmental conditions such as temperature
and light intensity without, however, any variance in geno-
type.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Doery-
wape’. These characteristics in combination distinguish
‘Doerywape’ as a new and distinct *Erysimum* plant:

1. Compact, upright and mounding plant habit.
2. Moderately vigorous growth habit.

3. Freely branching habit; dense and bushy habit.
4. Freely and continuous flowering habit.
5. Long flowering period.
6. Large white-colored flowers.
7. Tolerant to low and high temperatures and good garden performance.

Plants of the new *Erysimum* differ primarily from plants of the female parent selection in flower color as plants of the new *Erysimum* have white-colored flowers whereas plants of the female parent selection have pastel pink and yellow-colored flowers.

Plants of the new *Erysimum* differ primarily from plants of the male parent selection in flower color as plants of the new *Erysimum* have white-colored flowers whereas plants of the male parent selection have yellow-colored flowers.

Plants of the new *Erysimum* can be compared to *Erysimum hybrida* 'Poem Pastel', not patented. In side-by-side comparisons, plants of the new *Erysimum* differ from plants of 'Poem Pastel' in the following characteristics:

1. Plants of the new *Erysimum* are more compact than plants of 'Poem Pastel'.
2. Plants of the new *Erysimum* have larger flowers than plants of 'Poem Pastel'.
3. Flowers of plants of the new *Erysimum* are white in color whereas flowers of plants of 'Poem Pastel' are pastel pink and yellow in color.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Erysimum* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Erysimum* plant. The photograph is a top perspective view of a typical flowering plant of 'Doerywape' grown in a container.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations and measurements describe plants grown in 17-cm containers in a glass-covered greenhouse in Aalsmeer, The Netherlands and under cultural practices typical of commercial *Erysimum* production. During the production of the plants, average daily temperatures were 21° C. and average night temperatures were 15° C. Plants were 22 weeks old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, Second Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Erysimum hybrida* 'Doerywape'.
Parentage:

Female, or seed, parent.—Proprietary selection of *Erysimum hybrida* identified as code number ER16-000001-023, not patented.

Male, or pollen, parent.—Proprietary selection of *Erysimum hybrida* identified as code number ER16-000001-028, not patented.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate roots, summer.—About twelve days at temperatures about 26° C.

Time to initiate roots, winter.—About 16 days at temperatures about 23° C.

Time to produce a rooted young plant, summer.—About 16 days at temperatures about 23° C.

Time to produce a rooted young plant, winter.—About 20 days at temperatures about 18° C.

Root description.—Medium in thickness, fibrous; typically white in color, actual color of the roots is dependent on substrate composition, water quality, fertilizer type and formulation, substrate temperature and physiological age of roots.

Rooting habit.—Moderately freely branching; medium density.

Plant description:

Plant form and growth habit.—Perennial garden or container plant; compact, upright to mounding plant habit; freely branching habit with about 43 lateral branches developing per plant; dense and bushy appearance; moderately vigorous growth habit and moderate growth rate.

Plant height.—About 26 cm.

Plant diameter (area of spread).—About 31 cm.

Lateral branches.—Length: About 17 cm. Diameter: About 4 mm. Internode length: About 5 mm. Strength: Strong. Aspect: Upright to about 45° from vertical. Texture and luster: Smooth, glabrous; semi-glossy. Color, developing and developed: Close to 144A.

Leaf description:

Arrangement.—Alternate; simple; sessile.

Length.—About 8 cm.

Width.—About 1 cm.

Shape.—Narrowly oblanceolate to lanceolate.

Apex.—Acute.

Base.—Cuneate.

Margin.—Entire with some serrations.

Texture and luster, upper and lower surfaces.—Smooth, glabrous; semi-glossy.

Venation pattern.—Single midvein discernible.

Color.—Developing leaves, upper surface: Close to 137A. Developing leaves, lower surface: Close to 137C. Fully developed leaves, upper surface: Close to 137A; venation, close to 144B. Fully developed leaves, lower surface: Close to 138B; venation, close to 138B.

Flower description:

Flower type and habit.—Large rounded cruciform flowers arranged in dense racemes; flowers face mostly upright; freely flowering habit, about twelve flowers per inflorescence and about 350 flowers developing per plant.

Natural flowering season.—Relatively long flowering period; plants flower continuously from spring until autumn in The Netherlands; flowers not persistent.

Fragrance.—Faintly to moderately fragrant; sweet, pleasant.

Inflorescence height.—About 5 cm.

Inflorescence diameter.—About 4 cm.

Flower diameter.—About 3 cm.

Flower depth (height).—About 1.2 cm.

Flower buds.—Length: About 1 cm. Diameter: About 4 mm. Shape: Elliptic. Texture and luster: Smooth, glabrous; matte. Color: Close to 1C.

Petals.—Quantity and arrangement: Four petals arranged in a single whorl. Length: About 1.3 cm.

Width: About 1.2 cm. Shape: Obovate. Apex: Obtuse with truncate tendencies. Base: Cuneate. Margin: Entire, slightly undulate. Texture and luster, upper and lower surfaces: Smooth, glabrous; matte. Color: When opening, upper and lower surfaces: Close to 6D. Fully opened, upper and lower surfaces: During development, flowers transition through the following colors, close to 6D, 8D and 155A; venation, similar to lamina colors.

Sepals.—Quantity and arrangement: Four sepals arranged in a single whorl; calyx, tubular. Length: About 1 cm. Width: About 2 mm. Shape: Subulate. Apex: Acuminate. Base: Cuneate. Margin: Entire. Texture and luster, upper and lower surfaces: Smooth, glabrous; semi-glossy. Color, upper and lower surfaces: Close to 144B.

Pedicels.—Length: About 5 mm. Width: About 1 mm. Strength: Flexible. Aspect: About 50° from peduncle axis. Texture and luster: Smooth, glabrous; matte. Color: Close to 144A.

Reproductive organs.—Stamens: Quantity per flower: Six. Filament length: About 9 mm. Filament color:

Close to 145A. Anther shape: Elliptic; dorsifixed. Anther size: About 2 mm by 1 mm. Anther color: Close to 145C. Pollen amount: Abundant. Pollen color: Close to 158A. Pistils: Quantity per flower: One. Pistil length: About 7 mm. Style length: About 1 mm. Style color: Close to 144A. Stigma diameter: About 1 mm. Stigma shape: Rounded, capitate. Stigma color: Close to 144A. Ovary color: Close to 155D. Seeds: About 20 seeds develop per flower.

Pathogen & pest resistance: To date, plants of the new *Erysimum* have not been noted to be resistant to pathogens or pests common to *Erysimum* plants.

Garden performance: Plants of the new *Erysimum* have been observed to have good garden performance and to tolerate temperatures ranging from about -20° C. to about 35° C.; suitable for U.S.D.A. Hardiness Zone 6.

It is claimed:

1. A new and distinct *Erysimum* plant named 'Doerywape' as illustrated and described.

* * * * *

