



US00PP34335P2

(12) **United States Plant Patent**
Bergman

(10) **Patent No.:** **US PP34,335 P2**
(45) **Date of Patent:** **Jun. 14, 2022**

(54) **CHRYSANTHEMUM PLANT NAMED**
‘CIDZ0110’

(50) Latin Name: *Chrysanthemum x morifolium*
Varietal Denomination: **CIDZ0110**

(71) Applicant: **SYNGENTA CROP PROTECTION**
AG, Basel (CH)

(72) Inventor: **Wendy R. Bergman, Gilroy, CA (US)**

(73) Assignee: **Syngenta Crop Protection AG, Basel**
(CH)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **17/361,502**

(22) Filed: **Jun. 29, 2021**

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/14 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./296**

(58) **Field of Classification Search**

USPC Plt./296, 286
CPC ... A01H 5/02; A01H 5/00; A01H 6/14; A01H
6/1424; A01H 6/385; A01H 5/08; A01H
6/7481

See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

Upov Pluto Plant Variety Database Sep. 4, 2021, retrieved on Sep.
7, 2021, retrieved from the Internet at [https://www3.wipo.int/pluto/](https://www3.wipo.int/pluto/user/en/index.jsp)
[user/en/index.jsp](https://www3.wipo.int/pluto/user/en/index.jsp), one page. (Year: 2021).*

* cited by examiner

Primary Examiner — June Hwu

(74) *Attorney, Agent, or Firm* — Dale Skalla

(57) **ABSTRACT**

A new *Chrysanthemum* plant named ‘CIDZ0110’ having
large sized orange bronze daisy-type inflorescences, com-
pact strong and uniform plant habit, and a flowering
response time of 7 weeks.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed:
Chrysanthemum x morifolium.
Varietal denomination: ‘CIDZ0110’.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new *Chrysanthemum*,
botanically known as *Chrysanthemum x morifolium*, and
hereinafter referred to by the variety name ‘CIDZ0110’.

‘CIDZ0110’ is a product of a planned breeding program.
The new cultivar has large sized orange bronze daisy-type
inflorescences, compact strong and uniform plant habit, and
a flowering response time of 7 weeks.

‘CIDZ0110’ originated from a hybridization made in
August 2013 in a greenhouse in Gilroy, Calif. The female
parent was the unpatented, proprietary plant designated as
‘B7017’, having light purple daisy-type inflorescences,
slightly smaller flower size and more vigorous growth when
compared to ‘CIDZ0110’.

The male parent of ‘CIDZ0110’ was the unpatented,
proprietary plant designated as ‘B8680’, having orange
bronze daisy-type inflorescences, abundant pollen produc-
tion, smaller flower size and slower flowering response
when compared to ‘CIDZ0110’.

The seed was sown in September 2014 and the resulting
plants were grown in pots in a greenhouse in Gilroy, Calif.
‘CIDZ0110’ was selected as one flowering plant within the
progeny of the stated cross in January 2015.

The first act of asexual reproduction of ‘CIDZ0110’ was
accomplished when vegetative cuttings were propagated
from the initial selection in April 2015 in Gilroy, Calif.

BRIEF SUMMARY OF INVENTION

Horticultural examination of plants grown from cuttings
of the plant initiated in April 2015 and continuing thereafter,

2

has demonstrated that the combination of characteristics as
herein disclosed for ‘CIDZ0110’ are firmly fixed and are
retained through successive generations of asexual repro-
duction.

‘CIDZ0110’ has not been observed under all possible
environmental conditions. The phenotype may vary signifi-
cantly with variations in environment such as temperature,
light intensity and day length.

A Plant Breeder’s Right for this cultivar was applied for
in Canada on Aug. 4, 2021, No. 21 -10679. ‘CIDZ0110’ has
not been made publicly available prior to the effective filing
date of this application, notwithstanding any disclosure that
may have been made less than one year prior to the effective
filing date of this application by the inventor or another who
obtained ‘CIDZ0110’ directly from the inventor.

The following traits have been repeatedly observed and
are determined to be basic characteristics of the new variety.
The combination of these characteristics distinguishes this
Chrysanthemum as a new and distinct variety.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographic drawings show typical
flower and foliage characteristics of ‘CIDZ0110’ with colors
being as true as possible with an illustration of this type.

The photographic drawings show in FIG. 1 a close-up
view of an inflorescence, and in

FIG. 2 a flowering plant of the new variety.

DETAILED BOTANICAL DESCRIPTION

The plant descriptions and measurements were taken
indoors in Hollister, Calif. in May 2021 on about 9 week old

plants. Unrooted cuttings had been planted, 4 plants per pot, in 6-inch pots on Mar. 3, 2021. The plants were pinched and moved to short days on Mar.24, 2021 to induce flowering. The aforementioned photographs: FIGS. 1 and 2 were taken of the same plants used for the descriptions and measurements in Hollister, Calif. in May 2021. Color references are made to The Royal Horticultural Society Colour Chart (RHS) 2001.

TABLE 1

DIFFERENCES BETWEEN THE NEW VARIETY 'CIDZ0110' AND A MOST SIMILAR VARIETY:		
	'CIDZ0110'	'Yoauburn', U.S. Plant Pat. No. 12,526
Plant size and habit:	Shorter	Taller
Flowering response:	Faster	Slower
Flower size:	Larger	Smaller

Plant:

Form, growth and habit.—Herbaceous pot-type, stems upright to moderately angled, freely branching, strong and compact growth habit.
Plant height.—15.0-17.0 cm.
Plant height (inflorescence included).—21.0 cm.
Plant width.—31.0 cm.

Roots:

Number of days to initiate roots.—4 days at about 22 degrees C.
Number of days to produce a rooted cutting.—10-12 days at 22 degrees C.
Type.—Fine, fibrous, free branching.
Color.—RHS N155B but whiter.

Foliage:

Arrangement.—Alternate.
Immature, leaf color, upper surface.—RHS 137A.
Lower surface.—RHS 137C.
Mature, leaf color, upper surface.—RHS 139A.
Lower surface.—RHS 138B.
Length.—5.5-6.3 cm.
Width.—4.1-5.3 cm.
Shape.—Ovate.
Base shape.—Attenuate.
Apex shape.—Mucronulate.
Margin.—Palmately lobed; irregularly incised, somewhat serrate.
Number of margin indentations.—Few.
Texture, upper surface.—Puberlulent.
Lower surface.—Puberlulent.
Color of veins, upper surface.—RHS 138C.
Color of veins, lower surface.—RHS 138B.
Pattern of veining.—Palmate.
Petiole color.—RHS 139A.
Length.—1.3-2.5 cm.
Diameter.—0.4-0.5 cm.
Stipule.—Irregularly present.
Stipule length.—0.3-1.2 cm.
Stipule color.—RHS 137A.
Texture.—Puberlulent.

Stem:

Quantity of main branches per plant.—4-6.
Color of stem.—RHS 138B.
Length of stem.—11.5-14.5 cm.
Diameter.—0.4-0.5 cm.
Length of internodes.—0.7-2.6 cm.

Texture.—Puberlulent.
Color of peduncle.—RHS 138B.
Length of peduncle.—3.0-6.5 cm.
Peduncle diameter.—0.2-0.25 cm.
Texture.—Puberlulent.

Inflorescence:

Type.—Compositae type, single-type inflorescences borne terminally above foliage, ray florets arranged acropetally on a capitulum.
Quantity of short days to flowering (response time).—7 weeks.
Natural season flowering.—Not determined for this variety.
Quantity of inflorescences per plant.—14-21.
Lastingness of individual blooms on the plant.—4 weeks.
Fragrance.—Slightly spicy.

Bud (just when opening/showing color):

Color.—RHS 161C.
Length.—0.9 cm.
Width.—1.1 cm.
Shape.—Oblate.

Immature inflorescence:

Diameter.—5.6 cm.
Color of ray florets, upper surface.—Closest to RHS 167A.
Lower surface.—Closest to RHS 162A with tones of 167C.

Mature inflorescence:

Diameter.—9.0-9.8 cm.
Depth.—3.0 cm.
Total diameter of disc.—1.8 cm.
Receptacle color.—RHS 145B.
Receptacle height.—0.4 cm.
Receptacle diameter.—0.8 cm.

Ray florets:

Average quantity of florets.—22.
Type.—Ligulate.
Color of florets, upper surface.—Closest to RHS 167A with tones of RHS 22A and RHS 172C.
Lower surface.—Closest to RHS 162A with tones of RHS 167C.
Length.—4.2-4.7 cm.
Width.—1.2-1.3 cm.
Longitudinal axis.—Straight.
Shape.—Elliptical.
Apex shape.—Mostly mucronate and a few asymmetrically cleft.
Corolla tube.—Short.
Margin.—Entire.
Texture, upper surface.—Papillose.
Lower surface.—Papillose.

Disc florets:

Average quantity of florets.—297.
Color of immature florets.—Light green.
Color of mature florets.—RHS 8A.
Length.—0.7 cm.
Width.—0.1 cm.
Shape.—Tubular, elongated.
Apex shape.—Acute, 5 pointed.

Phyllaries:

Quantity.—Average of 29.
Color, upper surface.—RHS 137A.
Lower surface.—RHS 137A.
Length.—0.7-1.2 cm.

Width.—0.3-0.4 cm.
Shape.—Lanceolate.
Apex shape.—Acute.
Base.—Fused.
Margins.—Entire; slightly papery.
Texture, upper surface.—Glabrous.
Lower surface.—Puberlulent.
Reproductive organs:
Pistil.—1, found on both types of florets.
Length.—0.7 cm.
Style color.—RHS 145D.
Style length.—0.5 cm.
Stigma color.—RHS 4B.
Stigma shape.—Bi-parted.
Ovary color.—RHS 155C but more translucent.
Stamens.—4, found on only on the disc florets.
Color of filaments.—RHS 155C.

Length filaments.—0.2 cm.
Anther color.—RHS 12A.
Anther length.—0.15 cm.
Anther shape.—Oval.
5 *Color of pollen.*—RHS 12A.
Pollen amount.—Scant.
Fertility/seed set.—Good.
Disease/pest resistance.—Has not been observed to
date.
10 *Heat and cold resistance.*—Has not been determined to
date.

What is claimed is:

15 1. A new and distinct variety of *Chrysanthemum* plant
named 'CIDZ0110' substantially as illustrated and described
herein.

* * * * *



FIG. 1



FIG. 2