

US00PP36761P3

# (12) United States Plant Patent Reynbery

#### US PP36,761 P3 (10) Patent No.:

## (45) Date of Patent:

Jun. 24, 2025

#### CANNABIS PLANT NAMED 'PTC8506V'

- Latin Name: Cannabis sativa Varietal Denomination: **PTC8506V**
- Applicant: Phylos Bioscience, Inc., Portland, OR (US)
- **Jared Reynbery**, Portland, OR (US)
- Assignee: **Phylos Bioscience, Inc.**, Portland, OR

(US)

Subject to any disclaimer, the term of this Notice:

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

Appl. No.: 18/643,930

Filed: Apr. 23, 2024 (22)

(65)**Prior Publication Data** 

> US 2024/0381800 P1 Nov. 14, 2024

Int. Cl. (51)

A01H 5/00 (2018.01)A01H 6/28 (2018.01)

U.S. Cl. (52)

Field of Classification Search (58)

See application file for complete search history.

**References Cited** (56)

#### **PUBLICATIONS**

UPOV-PLUTO citation for 'PTC8506V'. retrieved from UPOV on Oct. 20, 2024. https://pluto.upov.int/result. 1 page. (Year: 2023).\*

\* cited by examiner

Primary Examiner — Karen M Redden (74) Attorney, Agent, or Firm — Klarquist Sparkman,

LLP

#### (57)**ABSTRACT**

A new and distinct variety of Cannabis sativa designated 'PTC8506V' as herein described and illustrated, characterized by elevated levels of THCV.

1 Drawing Sheet

Genus/species: Cannabis sativa. Variety denomination: 'PTC8506V'.

#### BACKGROUND

Cannabis plants contain over a hundred known cannabinoids, which bind to endogenous endocannabinoid receptors. Varinolic cannabinoids, as known as varins, are a type of cannabinoid compounds having three carbon atoms in their alkyl side chain instead of the five carbon atom alkyl 10 side chains more commonly associated with cannabinoids. Two such varins are tetrahydrocannabivarin (THCV) and cannabidivarin (CBDV), which are homologues of tetrahymolecular targets.

THCV and CBDV have potential benefits across a broad set of applications. Cannabis plants or extracts with high THCV levels, for example, can be used as an agent for 20 anticonvulsant activity, obesity-associated glucose intolerance, appetite suppression, anxiety management for PTSD, diabetic neuropathy, and major neuropathic and pain related pathologies. Another THCV application is that of an appetite suppressing compound. CBDV has been shown to have 25 anti-epileptic and anticonvulsant activity.

Research and development as well as the sale of varin products has been limited due to low commonly occurring levels of varins in *Cannabis* flower. The ability to produce Cannabis with high varin levels will create a platform for a 30 new Cannabinoid category with differentiated, high margin products in both medical and recreational markets. The present application solves these problems by providing a

novel Cannabis plant having high THCV, which is referred to by the variety name 'PTC8506V'.

#### SUMMARY

The new variety of Cannabis sativa was created by sourcing Cannabis plants that produce THCV and THC. Crosses were conducted to elevate further THCV, THC and/or other varins. All breeding, cultivation, and chemotype analyses were conducted in Hillsboro, Oregon. 'PTC8506V' resulted from the cross of a high THC pollen receiver plant ('21GAQ-1271', unpatented) and a high THCV pollen donor plant ('21VLP5-1-102', unpatented). drocannabinol (THC) and cannabidiol (CBD), respectively.

15 22.2%, and a THCV content below the limit of detection (LOD). '21GAQ-1271' has greenish-purple flowers with heavy trichome coverage, moderate density, and strong aroma. '21GAQ-1271' has an upright plant growth habit with moderately long internodes and matures in 8-9 weeks. '21VLP5-1-102' had a THC content of 1.2%, and THCV content of 5.66%. '21VLP5-1-102' flowers are greenishpurple with low trichome coverage, very low density, and moderately weak aroma. '21VLP5-1-102' has a moderately wide plant growth habit with very long internodes and matures in 10-12 weeks. A clone of '21VLP5-1-102' was crossed onto a clone of '21GAQ-1271' in 2021. 32 of the resulting F1 seed were sown. 'PTC8506V' was discovered and selected as a single plant from this F1 seed and was subsequently propagated clonally in Hillsboro, OR. 'PTC8506V' was asexually reproduced by apical and subapical semi-herbaceous stem cuttings in Hillsboro, Oregon for eight cycles. Asexual reproduction has demonstrated that 'PTC8506V' reproduces true to type with all of the charac-

teristics, as herein described, and firmly fixed and retained through successive generations of asexual propagation.

The new variety is named 'PTC8506V' and is characterized as having elevated levels of THCV. In comparison to its parents, 'PTC8506V' has higher THCV (relative to either parent), and higher THC relative to the pollen donor plant. 'PTC8506V' can additionally be distinguished from either parent at least based on the characteristics provided in Table 1

TABLE 1

Comparison to Parent Varieties.						
Variety	Trichome Coverage	Aroma Strength	Aroma Type	Growth Habit	Inter- node	Maturity Time
'21GAQ- 1271'	Heavy	Strong	Sweet cookies, earth, garlic	Upright	Moder- ately long	8-9 weeks
'21VLP5-1- 102'	Low	Moder- ately weak	Grass, earth, citrus	Moder- ately wide	Long	10-12 weeks
'PTC8506V'	Moderate	Moder- ately strong	Pine, earth, fuel, and a slight fruiti- ness	Upright	Moder- ately long	8-10 weeks

#### BRIEF DESCRIPTION OF THE PHOTOGRAPH

FIG. 1 is a photograph of a flowering plant of 'PTC8506V' that is approximately 83 days old, growing in Hillsboro, Oregon.

### DETAILED BOTANICAL DESCRIPTION

The following is a detailed description of the new *Cannabis* variety 'PTC8506V'. Unless otherwise indicated, all 40 color designations provided herein are hex color codes.

All breeding, cultivation, and chemotype analyses were conducted in Hillsboro, Oregon.

Cannabis plants having elevated THCV or THC levels were sourced and obtained, and crosses were conducted to produce new varieties to further elevate THCV and THC levels. The plants were crossed to produce new varieties having elevated THCV and THC levels. Plant variety 'PTC8506V' was selected and asexually reproduced.

Table 2 describes the general characteristics of <sup>50</sup> 'PTC8506V' at 55 days after lighting was changed to 12 hours on/12 hours off.

TABLE 2

Plant life forms	An herbaceous plant (herb)
Plant growth habit	An upright, tap-rooted annual
	plant, forming fibrous
	roots when
	asexually propagated
Plant origin	A controlled cross between
	pollen acceptor
	('21GAQ-1271') and pollen donor
	plant ('21VLP5-1-102).
Plant propagation	Asexually propagated by apical
	and sub-apical semi-
	herbaceous stem cuttings

#### TABLE 2-continued

5	Propagation Ease Plant shape Height Width Time to harvest Resistance to pests or diseases GMO?	Easy Simple erect 117 cm to 121 cm 87 cm to 101 cm 10-12 weeks Unknown No
---	--	---

Table 3 describes the leaf/foliage of 'PCT8506V'. Color codes are hex color codes.

#### TABLE 3

15	Leaf Arrangement	Alternate
	Leaflet Type	Palmately compound
	Leaflet Structure	Serrated margins, lightly acicular
		to lanceolate leaflets,
		tapering to an acuminate apex
	Leaflet Margins	Jaggedly serrate with each tooth
•		apex angled toward
20		leaflet apex
	Leaf hairs	Present on both upper and
		lower surfaces
	Leaf length with petiole at	18.5 cm to 23 cm
	maturity	10.5 011 to 25 011
	Petiole length at maturity	6 cm to 9.5 cm
25	Petiole diameter at maturity	1.9 mm to 2.5 mm
	Petiole color	#6B4247
	Anthocyanin color and intensity	#F9CCC5. Medium intensity
	in Petioles	in vegetative stage;
	III I CHOICS	stronger in flowering stage
	Stipule length at maturity	6.2 mm to 6.8 mm
30	Stipule shape	Linear with slight curvaceous
50	Supure snape	
	Stimula color	growth #3C4D03
	Stipule color No. of leaflets	3 to 7
	Middle largest (longest) leaflet	12 cm to 17.5 cm
2.5	length Middle largest (langest) leaflet	2.5 cm to 3.1 cm
35	Middle largest (longest) leaflet width	2.5 cm to 5.1 cm
		2 0 to 1 0
	Middle largest (longest) leaflet	3.8 to 4.8
	length/width ratio  No. teeth of middle leaflet	20 to 24
		28 to 34 #35530A
	Leaf (upper side-adaxial) color	
<b>4</b> 0	Leaf (lower side-abaxial) color	#35530A
	Leaf glossiness	Matte Obliguely continuous
	Vein/midrib shape	Obliquely continuous
	Voin/midule calar	throughout leaflet
	Vein/midrib color	#E6DAA6
	Aroma	Earthy notes of pine, fuel, and
45		a slight fruitiness
1.5		

Table 4 describes the stem of 'PTC8506V'.

### TABLE 4

Stem shape	Hollow, ribbed, texture
Stem length	117 cm to 121 cm
Stem strength	Strong and upright
Stem diameter at base	13.6 mm to 14.5 mm
Stem color	#373E02
Depth of main stem ribs/grooves	Shallow
Internode length	3 cm to 9.5 cm

Table 5 describes the inflorescence of 'PTC8506V'.

TARI	$\mathbf{F}$

55

60

	Inflorescence type	Elongated compound cymes
	Flowering (blooming) habit	Natural flowering season is
		Autumn in Hillsboro, Oregon.
	Proportion of female plants	100% pistillate
5	Inflorescence position	Terminal and axillary

6

#### TABLE 5-continued

Number of inflorescences	100 to 150
	100 to 130
produced per plant Flower arrangement	Cymose
Length of inflorescences	3 cm to 3.8 cm
Numbers of flowers per plant	Hundreds to thousands flowers per
Numbers of nowers per plant	
	plant, multitudinous, congested
	with high concentrations on bushier wide inflorescence
T11	
Flower shape	Urceolate
Flower (individual pistillate)	5.1 mm to 5.3 mm
length	
Flower (individual pistillate)	2 mm
depth	
Flower (compound cyme)	5.5 cm to 8 cm
diameter	
Corolla	No defined corolla
Corolla color	N/A
Bract shape	Urceolate
Bract color	#4B5D16
Typical bract numbers	3 to 200 bracts per inflorescence.
Bract length	3 mm to 5mm
Bract width	2.3 mm to 2.7 mm
Bract apex and base description	Bracts have a rounded base and
1	caudate apex.
Bract Texture	Bract texture is smooth and
	covered with sticky trichomes.
Bracteole number, shape, length,	Bracteoles are not present
width, apex, margin, base	Diacteores are not present
descriptors, texture, and color	
Calyx shape	No defined calyx
Caryx snape	
Colve color	NI/A
Calyx color	N/A One
Pistil number per flower	One
•	One Pistil consists of the stigma, style,
Pistil number per flower	One Pistil consists of the stigma, style, and ovary and cannot be
Pistil number per flower Pistil color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color.
Pistil number per flower Pistil color Stigma shape	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length Peduncle diameter	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm 4.1 mm
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length Peduncle diameter Peduncle texture	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm 4.1 mm Lightly ribbed
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length Peduncle diameter Peduncle color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm 4.1 mm Lightly ribbed #9BB53C No staminate flowers produced
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length Peduncle diameter Peduncle color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm 4.1 mm Lightly ribbed #9BB53C No staminate flowers produced naturally; however, male
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length Peduncle diameter Peduncle color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm 4.1 mm Lightly ribbed #9BB53C No staminate flowers produced naturally; however, male flower (staminate) can be
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length Peduncle diameter Peduncle color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm 4.1 mm Lightly ribbed #9BB53C No staminate flowers produced naturally; however, male flower (staminate) can be induced with stress or ethylene-
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length Peduncle diameter Peduncle texture Peduncle color Staminate shape	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm 4.1 mm Lightly ribbed #9BB53C No staminate flowers produced naturally; however, male flower (staminate) can be induced with stress or ethylene- blocking compounds
Pistil number per flower Pistil color  Stigma shape Number of stigmas per flower Stigma length Stigma color Trichome shape Trichome color Terminal inflorescence shape Terminal inflorescence color Terminal inflorescence length Terminal inflorescence width Pedicel Peduncle length Peduncle diameter Peduncle color	One Pistil consists of the stigma, style, and ovary and cannot be given a single color. Acute Two 4.6 mm to 6.5 mm #373E02 Capitate-stalked glandular #F1E7E2 Oblong #B7E1A1 12.5 cm to 14 cm 5 cm to 6.5 cm Absent 6.8 mm 4.1 mm Lightly ribbed #9BB53C No staminate flowers produced naturally; however, male flower (staminate) can be induced with stress or ethylene-

Apetalous

Petal description

Table 6 describes other characteristics of 'PTC8506V'

TABLE 6
---------

5	Time period and condition of flowering/blooming	8 to 10 weeks
	Proportion of hermaphrodite plants	Lower (<5%) in clones
1.0	Hardiness of plant	Hardy to tolerate of warm and dry conditions without showing stress.
10		Not tolerant to soil oversaturated with water for long periods

Chemical analyses were conducted using High Performance Liquid Chromatography to detect the cannabinoids described herein. Table 7 describes a chemotype analysis of 'PTC8506V', which was chemotyped by taking a sample from the main cola at day 56 after initiation of the 12:12 light cycle, which initiates flowering. Trichomes were 70% cloudy and 5% amber, indicative of full maturity at sampling.

TABLE 7

	Cannabinoid and other chemicals	% wt
25	Total THC	4.613
	Total THCV	11.067
	Total CBG	0.047
	Total CBD	<lod< td=""></lod<>
	Total CBC	<lod< td=""></lod<>
20	Total CBGV	<lod< td=""></lod<>
30	Total CBDV	<lod< td=""></lod<>
	alpha-Pinene	0.057
	Camphene	0.011
	beta-Myrcene	1.557
	beta-Pinene	0.098
2.5	delta-3-Carene	<loq< td=""></loq<>
35	alpha-terpinene	<loq< td=""></loq<>
	D-Limonene	0.211
	beta-Ocimene	0.275
	p-Cymene	<loq< td=""></loq<>
	Eucalyptol	<loq< td=""></loq<>
	gamma-Terpinene	<loq< td=""></loq<>
40	Terpinolene	<loq< td=""></loq<>
	Linalool	0.003
	Isopulegol	<loq< td=""></loq<>
	Geraniol	<loq< td=""></loq<>
	beta-Caryophyllene	0.05
	alpha-Humulene	0.0227
45	Nerolidol 1	<loq< td=""></loq<>
	Nerolidol 2	0.011
	Guaiol	<loq< td=""></loq<>
	Caryophyllene Oxide	0.011
	alpha-Bisabolol	0.026
	Total Terpenes	2.315

What is claimed is:

50

1. A new and distinct variety of *Cannabis sativa* designated 'PTC8506V' as herein described and illustrated.

\* \* \* \* \*

