

US00PP36997P2

(12) **United States Plant Patent**
NeSmith et al.

(10) **Patent No.:** **US PP36,997 P2**
(45) **Date of Patent:** **Oct. 7, 2025**

(54) **BLUEBERRY PLANT NAMED ‘CBG-23-05’**

CPC . A01H 5/08; A01H 5/00; A01H 6/368; A01H 6/36

(50) Latin Name: *Vaccinium corymbosum*
Varietal Denomination: **CBG-23-05**

See application file for complete search history.

(71) Applicants: **Rio Vista Farms Inc.**, Selma, CA (US);
University of Georgia Research
Foundation, Inc., Athens, GA (US)

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP21,167 P2 * 7/2010 NeSmith A01H 6/368
Plt./157

(72) Inventors: **D. Scott NeSmith**, Griffin, GA (US);
Gregory Willems, Kingsbury, CA (US)

(73) Assignees: **Rio Vista Farms Inc.**, Selma, CA (US);
University of Georgia Research
Foundation, Inc., Athens, GA (US)

OTHER PUBLICATIONS

My Little Green Garden, How to Select Blueberry Varieties for Your Location 2022, retrieved on Jan. 8, 2025 at <https://mylittlegreengarden.com/blueberry-varieties/>, 14 pp. (Year: 2022).*

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

* cited by examiner

Primary Examiner — June Hwu

(21) Appl. No.: **18/885,323**

(74) Attorney, Agent, or Firm — Knobbe, Martens, Olson & Bear, LLP

(22) Filed: **Sep. 13, 2024**

(51) **Int. Cl.**
A01H 5/08 (2018.01)
A01H 6/36 (2018.01)

(57) **ABSTRACT**

The new variety of blueberry ‘CBG-23-05’ is characterized by having large, firm, berries with excellent color. The berries ripen early in the season and have a long shelf life. Further, the berries of the new variety ‘CBG-23-05’ are crunchy, with moderate sweetness and mild acidity.

(52) **U.S. Cl.**
USPC **Plt./157**

(58) **Field of Classification Search**
USPC **Plt./157**

11 Drawing Sheets

1

Latin name of the genus and species claimed: *Vaccinium corymbosum*.

Variety denomination: ‘CBG-23-05’.

REFERENCE TO RELATED APPLICATIONS

This application is related to U.S. application Ser. No. 18/884,792, filed Sep. 13, 2024, U.S. application Ser. No. 18/884,736, filed Sep. 13, 2024, U.S. application Ser. No. 18/885,153, filed Sep. 13, 2024, and U.S. application Ser. No. 18/885,323, filed Sep. 13, 2024, which are directed to blueberry varieties developed by the same breeding program.

BACKGROUND AND SUMMARY OF THE INVENTION

Disclosed herein is a new and distinct southern highbush blueberry variety (*Vaccinium corymbosum*). The varietal denomination of the new variety is ‘CBG-23-05’.

The new variety ‘CBG-23-05’ is a product of a hand-pollinated cross of seed parent ‘TH-1492’ (unpatented) and pollen parent ‘TH-1870’ (unpatented) made by Dr. Scott NeSmith in 2019 in Griffin, Georgia. The new variety ‘CBG-23-05’ was germinated from seed. The date of first planting of the new variety ‘CBG-23-05’ was November 2021. The new variety ‘CBG-23-05’ was selected by Gregory Willems in May 2023 in Reedley, California. Subsequent testing has demonstrated that the combination of

2

traits disclosed herein that characterize the new variety are stable and retained true to type through successive generations of vegetative, asexual reproduction in Fresno County, California.

5 The new variety of blueberry ‘CBG-23-05’ is characterized by having large, firm, berries with excellent color. The berries ripen early in the season and have a long shelf life. Further, the berries of the new variety ‘CBG-23-05’ are crunchy, with moderate sweetness and mild acidity. The new variety ‘CBG-23-05’ has not shown notable susceptibility to diseases or pest problems beyond those also common for other blueberry varieties. The new variety ‘CBG-23-05’ has a chilling requirement of about 150-200 hours below 45° F.

15 The phenotype of the new blueberry variety ‘CBG-23-05’ may vary somewhat with variations in environment, such as temperature and light intensity, and cultural practices, without, however, any variance in genotype.

COMPARISON TO THE PARENTS

20 The new variety ‘CBG-23-05’ is distinguished from its seed parent ‘TH-1492’ (unpatented) in that the fruit of the new variety is firm, compared to a medium firmness for the fruit of ‘TH-1492’. Further, the fruit of the new variety ‘CBG-23-05’ is sweeter and crunchier than the fruit of ‘TH-1492’. Additionally, the fruit of the new variety ‘CBG-23-05’ ripens later on about May 10th, compared to a mid-season ripening on about May 1st to May 5th for ‘TH-

1492'. Lastly, the new variety 'CBG-23-05' flowers around February 24th compared to around February 27th to March 3rd for 'TH-1492'.

The new variety 'CBG-23-05' is distinguished from its pollen parent 'TH-1870' (unpatented) by having very large to large-sized fruit, compared to the medium to the large-sized fruit of 'TH-1870'. Additionally, the fruit of the new variety 'CBG-23-05' ripens later, on about May 10th, compared to about April 23rd to April 28th for 'TH-1870'. Lastly, the new variety 'CBG-23-05' flowers around February 24th compared to around February 20th to February 23rd for 'TH-1870'.

COMPARISON TO CLOSEST VARIETY

The new variety 'CBG-23-05' resembles 'Suziblue' (U.S. Plant Pat. No. 21,167) and 'Emerald' (U.S. Plant Pat. No. 12,165) in that they are early season southern highbush blueberry cultivars with the same vigor. However, the new variety 'CBG-23-05' begins flowering earlier than 'Suziblue' and later 'Emerald' in Central California. Additionally, the new variety 'CBG-23-05' begins ripening later than 'Suziblue' and 'Emerald' in Central California, but the new variety 'CBG-23-05' has a similar harvest period as 'Suziblue' and 'Emerald'. Further the fruit of the new variety 'CBG-23-05' is firmer than that of 'Suziblue', but similar to the fruit of 'Emerald'.

TABLE 1

Table 1 shows a comparison of the characteristics of 'CBG-23-05', 'Suziblue' (Plant Pat. 21,167) and 'Emerald' (Plant Pat. No. 12,165). Rating scales are based on 1 to 10 score, with 1 being the least desirable and 10 being the most desirable.			
	'Suziblue'	'Emerald'	'CBG-23-05'
Berry Size	8.0	8.0	9.2
Berry Scar	7.0	7.5	7.5
Berry Color	7.0	8.0	8.0
Berry Firmness	7.3	8.5	8.5
Berry Flavor	7.0	7.0	7.5
Cropping	8.5	8.5	7
Plant Vigor	8.0	8.0	8.0
Date of 50% Flowering	Feb. 28	Feb. 15	Feb. 24
Date of 50% Ripening	May 7	May 1	May 10
Fruit Development Period	69 Days	60 Days	85 Days

TABLE 2

Table 2 shows a comparison of the characteristics of the new variety 'CBG-23-05' and related varieties 'CBG-23-01', 'CBG-23-02', 'CBG-23-03' and 'CBG-23-04'.					
	'CBG-23-05'	'CBG-23-01'	'CBG-23-02'	'CBG-23-03'	'CBG-23-04'
Season	Early to mid-season	Early	Early to mid-season	Early	Early to mid-season
Harvest date	May 3rd-June 1st	April 18th-May 12th	April 28th-May 12th	April 18th-May 12th	May 3rd-June 1st
Fruit size	Very large, 19 mm-22 mm	Large, 20 mm-24 mm	Large, 19 mm-24 mm	Large, 18 mm-21 mm	Large, 18 mm-21 mm
Flavor	Sweet with moderate acidity	Sweet, aromatic	Balanced sweetness and acidity	Balanced sweetness and acidity	Very sweet with mild acidity and floral undertones

TABLE 2-continued

Table 2 shows a comparison of the characteristics of the new variety 'CBG-23-05' and related varieties 'CBG-23-01', 'CBG-23-02', 'CBG-23-03' and 'CBG-23-04'.					
	'CBG-23-05'	'CBG-23-01'	'CBG-23-02'	'CBG-23-03'	'CBG-23-04'
Fruit Texture	Smooth, crisp	Smooth, crisp	Crisp, firm	Crisp	Crisp
Fruit Color	Medium blue	Deep blue	Medium blue	Light to medium blue	Medium blue

BRIEF DESCRIPTION OF THE ILLUSTRATIONS

The accompanying color photographic illustrations taken in Reedley, California in 2024 pertain to 2.5 year old plants and show the overall appearance and distinct characteristics of the new variety 'CBG-23-05'. The colors in the photographs are as true as possible but may differ slightly from the color values cited in the detailed botanical description, which accurately describes the colors of the new variety 'CBG-23-05'.

FIGS. 1-4 show the berries of the new variety 'CBG-23-05' on the plant ready for harvest.

FIG. 5 shows the growth pattern of the plants of the new variety 'CBG-23-05'.

FIG. 6 shows the plants of the new variety 'CBG-23-05' during bloom.

FIG. 7 shows the flowers of the new variety 'CBG-23-05'.

FIG. 8 shows the average berry size of the new variety 'CBG-23-05'.

FIG. 9 shows the average berry weight of the new variety 'CBG-23-05'.

FIG. 10 shows the fruit of the new variety 'CBG-23-05' cut in half.

FIG. 11 shows the fruit of the new variety 'CBG-23-05' as a whole.

DESCRIPTION OF THE NEW VARIETY

The following traits have been consistently observed in the original plant of the new variety 'CBG-23-05' and in asexually propagated progeny via vegetative cuttings in Fresno County, California.

The descriptive matter which follows pertains to 2.5-year-old 'CBG-23-05' plants grown in the vicinity of Reedley, California during 2023, with supplemental irrigation, and is believed to apply to plants of the variety grown under similar conditions of soil and climate elsewhere. The dimensions, sizes, colors, and other characteristics given are to be understood to be approximations and averages set forth as accurately as practicable.

Throughout this specification, color names beginning with a small letter signify that the name of that color as used in common speech is aptly descriptive. Color names beginning with a capital letter designate values based upon The R.H.S. Colour Chart published by The Royal Horticultural Society, London, England, Sixth Edition, 2019 reprint. The colors and other phenotypic attributes may deviate from the stated values and descriptions depending upon variation in environmental, seasonal, climatic and cultural conditions.

General: The new variety 'CBG-23-05' produces fruit from about May 3rd to June 1st. Measurements were taken on 2.5-year-old plants.

Plant:

Size.—About 3.0 feet tall, 8-inch plant base/crown. 5

Habit.—Vigorous vertical growth. Plants have 1-4 main canes with low spreading branches starting 30-40 cm from the soil. Vertical canes have vertically pushing branches and open broad leaf clusters alternating. 10

Productivity.—About 1.9 pounds per plant.

Cold hardiness.—Similar to other early ripening southern highbush varieties in USDA Plant Hardiness Zone 9b. 15

Chilling requirements.—Approximately 150 to 200 hours of temperatures at or below 45° F. to induce normal leafing and flowering during the spring.

Leafing.—Plants readily break numerous leaf buds simultaneously with anthesis. 20

Canes.—Main Cane: Diameter: Approximately 10 mm. Color: About Medium Grey-Brown 199B. Length: Approximately 1.1m to 1.5 m. Texture: Woody. First Major Branch: Diameter: Approximately 6.5 mm. Color: About Medium Yellow-Green N144C. Length: Approximately 0.8 m to 1 m. Texture: Woody. 2-year-Old Cane: Diameter: Approximately 8 mm. Color: About Medium Yellow-Green N144C. Length: Approximately 0.7m to 0.9 m. Texture: Woody. Current Season Wood: Diameter: Approximately 4 mm. Color: About Medium Yellow-Green 144C. Length: Approximately 0.5 m to 0.7 m. Texture: Smooth. 25

Fruiting wood.—Numerous twigs approximately 5 to 10 cm in length and internode lengths of approximately 25 mm are common. The diameter is approximately 3 mm to 4 mm and the texture is smooth. 30

Foliage:

Leaf color of mature leaves.—Top side color: About Dark Yellow-Green 147A. Under side color: About Medium Yellow-Green 148C. 40

Leaf arrangement.—Alternate simple.

Leaf shape.—Elliptic.

Leaf venation.—Pinnate with slight netting (or slightly reticulated). 45

Leaf apices.—Broadley acuminate.

Leaf base.—Cuneate to acute.

Leaf length.—Approximately 60 mm.

Leaf width.—Approximately 40 mm.

Leaf margin.—Entire. 50

Leaf surface.—Glaucous (waxy) on the top of the leaf, while the bottom of the leaf has a smooth texture without the glaucous coating.

Petiole size.—Small, approximately 4 mm long and 1 mm wide. 55

Petiole color.—About Light Yellow-Green 145C.

Petiole leaves margin.—The leaf margin is smooth or occasionally ciliate, with stalked glands present on the petiole or along the leaf margin. 60

Flower:

Date of 50% anthesis.—About February 24th in Central California. 65

Flower shape.—Urceolate.

Flower bud number.—Medium (5 to 7 buds per fruiting shoot). 65

Flowers per cluster.—Approximately 4 to 6.

Flower fragrance.—Yes, slight to moderate cut flower/rose fragrance.

Corolla.—Color: About Medium White NN155A to Medium White NN155B for both the inner and outer surfaces. Length: Approximately 7 to 9 mm. Width: Approximately 6 to 7.5 mm. Aperture Width: Approximately 3 to 4 mm.

Peduncle.—Length: About 15 to 20 mm. Color: About Medium Yellow-Green 144A. Diameter: Approximately 3 mm to 4 mm. Texture: Slightly woody.

Pedicel.—Length: Approximately 4 to 6 mm. Color: About Medium Green 138D. Diameter: Approximately 2 mm to 2.5 mm. Texture: Smooth.

Calyx.—Calyx (with sepals) Diameter: Approximately 5 to 6 mm. Color center of calyx: About Medium Green 138B. Color of sepals: About Medium Yellow-Green 144A. Average sepals per calyx: About 5. Sepal length: Approximately 4 mm. Sepal width: Approximately 3 mm.

Stamen.—Length: Approximately 5 to 6 mm. Number per flower: About 10. Filament color: About Dark Green-White 157A to Medium Green-White 157B.

Style.—Length: Approximately 6 to 8.5 mm. Color: About Medium Yellow-Green 145A.

Pistil.—Length: Approximately 8 to 9.5 mm. Ovary color: About Medium Green 138B.

Anther.—Length: Approximately 4 to 5 mm. Number: About 10. Color: About Medium Greyed-Orange 163B.

Pollen.—Abundance: Medium. Color: About Medium White NN155A.

Self-compatibility.—The cultivar has demonstrated a small degree of self-compatibility.

Fruit:

Date of 50% maturity.—About May 10th.

Fruit development period.—About 85 days.

Berry color.—About Light Violet-Blue 97B with wax, and about Dark Black 203C without wax.

Berry flesh color.—About Medium Green 141D.

Berry surface wax abundance.—Medium.

Berry weight.—1st Harvest: Approximately 4 g. 2nd Harvest: Approximately 4 g.

Berry size.—Height from Calyx to Scar: Approximately 16.4 mm. Diameter: Approximately 21 mm.

Berry shape.—Semi spherical to spherical.

Fruit stem scar.—Small to medium, dry and with no tears.

Calyx.—Approximately 3 mm deep and 6 mm wide.

Sepals.—Typically not visible; slightly turned inward and flat when present.

Berry firmness.—Firm, 8.5/10 on berry rating scale.

Berry flavor.—Moderate sweetness with mild acidity.

Berry texture.—Crisp with smooth texture.

Average brix.—Approximately 14.43.

Seed.—Seed Abundance in Fruit: High. Seed Color: About Medium Greyed-Orange 165B. Seed Dry Weight: Approximately 34.2 mg per 100 seeds. Seed Size: Approximately 1.2 to 1.8 mm. Seed Quantity Average per Berry: About 31. Seed Shape: The seeds of 'CBG-23-03' are generally ovate to slightly flattened, with a smooth surface.

Storage qualities.—Shelf life of 'CBG-23-05' is good to great. The fruit of 'CBG-23-05' was tested for storage longevity in a 39° F. refrigerated environment with low humidity for 21 days. The berries

maintained quality throughout the trial, retaining firmness with minimal shrivel or leakage.

Use/market.—The berries of ‘CBG-23-05’ can be used as fresh fruit for shipping, customer-pick and processing markets. Not suitable for machine harvest.

Disease resistance.—Resistant to fungal diseases, but otherwise no noticeable disease resistance or susceptibility has been observed for ‘CBG-23-05’.

We claim:

1. A new and distinct variety of blueberry plant named ‘CBG-23-05’ as illustrated and described herein.

* * * * *



FIG. 1



FIG. 2



FIG. 3



FIG. 4



FIG. 5



FIG. 6



FIG. 7




FIG. 8




FIG. 9



FIG. 10




FIG. 11