

[54] OXIDATION HAIR DYES BASED UPON TETRAAMINOPYRIMIDINE DEVELOPERS

[75] Inventors: David Rose, Düsseldorf-Holthausen; Ferdi Saygin, Erkrath; Erwin Weinrich, Düsseldorf-Holthausen, all of Fed. Rep. of Germany

[73] Assignee: Henkel Kommanditgesellschaft auf Aktien (Henkel KGaA), Düsseldorf-Holthausen, Fed. Rep. of Germany

[21] Appl. No.: 953,688

[22] Filed: Oct. 23, 1978

Related U.S. Patent Documents

Reissue of:

[64] Patent No.: 4,003,699
Issued: Jan. 18, 1977
Appl. No.: 526,232
Filed: Nov. 22, 1974

[30] Foreign Application Priority Data

Nov. 29, 1973 [DE] Fed. Rep. of Germany 2359399
Oct. 2, 1974 [DE] Fed. Rep. of Germany 2447017

[51] Int. Cl.² A61K 7/13
[52] U.S. Cl. 8/10.2; 8/11; 8/32

[58] Field of Search 8/10.2, 11, 32

[56] References Cited

U.S. PATENT DOCUMENTS

2,350,812 6/1944 Peterson 96/56.5
2,355,691 8/1944 Allen et al. 96/56.5
3,359,168 12/1967 Brechner et al. 8/10.2
3,536,436 10/1970 Lange 8/10.2

Primary Examiner—Joseph E. Evans
Assistant Examiner—Vera C. Clarke
Attorney, Agent, or Firm—Hammond & Littell

[57] ABSTRACT

An aqueous hair dye preparation comprising an oxidation dyestuff combination of a coupling component and a developer component consisting of a tetraaminopyrimidine derivative or a water-soluble acid addition salt thereof, as well as a process for dyeing hair by utilizing this oxidation dyestuff combination.

9 Claims, No Drawings

OXIDATION HAIR DYES BASED UPON TETRAAMINOPYRIMIDINE DEVELOPERS

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.

THE PRIOR ART

Of great importance for the dyeing of hair are the so-called oxidation dyestuffs because of their intensive colors and very good fastness. These dyestuffs are formed by the oxidative coupling of a developer component with a coupling component. The developers customarily used are nitrogenous bases, such as p-phenylenediamine derivatives, diaminopyridines, 4-aminopyrazolone derivatives or heterocyclic hydrazones. Useful as so-called coupling components are m-phenylenediamine derivatives, phenols, naphthols, resorcinol derivatives and pyrazolones.

Good oxidation dyestuff components for hair dyeing must fulfill all of the following requirements.

They have to be able to develop a sufficient intensity of the desired color shades when oxidatively coupled with the respective developer component or coupling component. Furthermore, they have to possess a capacity for being absorbed by human hair, which capacity ranges from sufficient to very good; and in addition, they should be unobjectionable from toxicological and dermatological viewpoints.

As developers, it is customary to use the class of compounds consisting of substituted or unsubstituted p-phenylenediamines. However, this class of compounds has the disadvantage that sensitivity reactions and subsequently severe allergies are caused in numerous persons. The developers which have been recently proposed for avoiding these dermatological disadvantages are not always fully satisfactory with respect to their technical application.

OBJECTS OF THE INVENTION


An object of the invention is to provide usable oxidation hair dyes containing suitable components which optimally satisfy the above requirements.

Another object of the present invention is to provide an oxidation dyestuff combination of a coupling component and a developer component, which is based on tetraaminopyrimidines as the developer component.

These and further objects of the present invention will become apparent as the description thereof proceeds.


DESCRIPTION OF THE INVENTION

The present invention provides a composition and process for dyeing hair based upon an oxidation dyestuff combination of a coupling component and a developer component with tetraaminopyrimidines being the developer component, as well as novel tetraaminopyrimidines. It has now been found that the abovespecified requirements can be fulfilled to an especially significant extent by the use of hair coloring preparations that are based on oxidation dyestuff combinations containing tetraaminopyrimidines of the formula



and their inorganic or organic water-soluble acid addition salts as developers, and containing the couplers customarily used in oxidation hair dyes, in which R_1 to R_6 are each selected from the group consisting of hydrogen, alkyl having 1 to 4 carbon atoms, aryl, substituted aryl and $-(CH_2)_n-X$, in which n is an integer from 1 to 4, and X is selected from the group consisting of hydroxy, halogen and $-NR_7R_8$ wherein R_7 and R_8 are selected from the group consisting of hydrogen and alkyl having 1 to 4 carbon atoms and can form with the nitrogen atom a heterocyclic ring which may contain one additional nitrogen atom or an oxygen atom. In the above formula, R_1 to R_6 likewise can designate an optionally substituted five-membered or six-membered heterocyclic ring containing one or two nitrogen atoms, or one nitrogen atom and one oxygen atom.

More particularly, the present invention is directed to an aqueous preparation for the dyeing of hair consisting essentially of (1) from 0.2% to 5% by weight of an oxidation dyestuff combination of a developer component, and a coupling component in substantially equimolar amounts, said developer component consisting essentially of (A) a tetraaminopyrimidine of the formula



wherein R_1 , R_2 , R_3 , R_4 , R_5 and R_6 are each selected from the group consisting of hydrogen, phenyl, alkyl having 1 to 4 carbon atoms, phenylalkyl having 7 to 10 atoms, phenylalkenyl having 7 to 10 carbon atoms,



wherein n is an integer from 1 to 4, and X is selected from the group consisting of hydroxyl, halogen and NR_7R_8- in which R_7 and R_8 are each hydrogen or alkyl having 1 to 4 carbon atoms, and together with the nitrogen atom R_7 and R_8 from a member selected from the group consisting of a 5 to 6 membered heterocyclic ring optionally containing an additional nitrogen atom or oxygen atom,

and wherein R_1 and R_2 , or R_3 and R_4 , or R_5 and R_6 , together with the nitrogen atom form a five to six membered heterocyclic ring optionally containing another nitrogen or oxygen atom in the ring and (B) a water-soluble acid addition salt of (A); (2) from 0% to 5% by weight of a direct dyestuff; (3) from 0% to 30% by weight of a surfactant; (4) from 0% to 25% by weight of thickeners; and (5) the balance up to 100% by weight of water.

A particularly preferred subgenus of the above-mentioned developer component is wherein R_1 , R_2 , R_3 , R_4 , R_5 and R_6 are each selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, n-propyl, butyl, phenyl, benzyl and benzylidene, or $-(CH_2)_n-X$

wherein R_1 and R_2 , or R_3 and R_4 , or R_5 and R_6 , together with the nitrogen atom form a substituent selected from the group consisting of hydroxyl, halogen and $-NR_7R_8$ in which R_7 and R_8 are each hydrogen or alkyl having 1 to 4 carbon atoms.

When the compounds according to the invention are used as developer components, they react with the known couplers generally used in oxidation hair dye-stuffs to give very intensive, varying shades which previously heretofore could not be effected with these known couplers and these developers known so far. Thus, the compounds of the invention considerably increase the possibilities for utilizing oxidation hair dyes. Furthermore, the tetraaminopyrimidines according to the invention are distinguished by very good fastness of the dyeings effects with them, by good water-solubility, by good storage stability, and by excellent toxicological as well as dermatological unobjectionableness.

The tetraaminopyrimidines which are to be used as developer components according to the invention can be used either as such or in form of their water-soluble acid addition salts with non-toxic inorganic acids or organic acids, such as for example, hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, propionic acid, lactic acid or citric acid.

The preparation of most tetraaminopyrimidines to be used as developer components according to the invention is already known in the literature and can be taken from the monograph by D. J. Brown, in the series "Heterocyclic Compounds", Interscience Publishers, 1962, Vols. I and II, "The Pyrimidines". Only a few of the compounds used are novel substances, the preparation of which is separately described below.

Specific examples of the novel tetraaminopyrimidines according to the invention are 5-amino-2,4,6-tris(methylamino)-pyrimidine, 2,4,5-triamino-6-(di-n-propylamino)-pyrimidine and 2,4,5-triamino-6-morpholino-pyrimidine, or a water-solution acid addition salt thereof.


To synthesize the compounds to be used according to the invention, the starting material generally is a 2,4,6-aminopyrimidine, into which the 5-amino group is introduced by nitrosation and subsequent reduction. It is also possible to start from the correspondingly substituted triaminoalkylmercaptopyrimidines and to replace the alkylmercapto group with an amino group. The latter method is especially suitable for the introduction of substituted amino groups into the 2-, 4-, or 6-positions of the pyrimidine ring. Suitable examples of developer components to be used according to the invention are, for example: 2,4,5,6-tetraamino-pyrimidine, 4,5-diamino-2,6-bis-(methylamino)-pyrimidine,

2,5-diamino-4,6-bis-(methylamino)-pyrimidine,
4,5-diamino-6-(butylamino)-2-(dimethylamino)-6-pyrimidine,
2,5-diamino-4-(diethylamino)-6-(methylamino)-pyrimidine,
4,5-diamino-6-(diethylamino)-(2-dimethylamino)-pyrimidine,
4,5-diamino-2-(diethylamino)-6-(methylamino)-pyrimidine,

4,5-diamino-2-(dimethylamino)-6-(ethylamino)-pyrimidine,
4,5-diamino-2-(dimethylamino)-6-(isopropylamino)-pyrimidine,
4,5-diamino-2-(dimethylamino)-6-(methylamino)-pyrimidine,
4,5-diamino-6-(dimethylamino)-2-(methylamino)-pyrimidine,
4,5-diamino-2-(dimethylamino)-6-(propylamino)-pyrimidine,
2,4,5-triamino-6-(dimethylamino)-pyrimidine,
4,5,6-triamino-2-(dimethylamino)-pyrimidine,
2,4,5-triamino-6-(methylamino)-pyrimidine,
4,5,6-triamino-2-(methylamino)-pyrimidine,
4,5-diamino-2-(dimethylamino)-6-piperidino-pyrimidine,
4,5-diamino-6-(methylamino)-2-piperidino-pyrimidine,
2,4,5-triamino-6-piperidino-pyrimidine,
2,4,5-triamino-6-anilino-pyrimidine,
2,4,5-triamino-6-(benzylamino)-pyrimidine,
2,4,5-triamino-6-(benzylideneamino)-pyrimidine,
4,5,6-triamino-2-piperidino-pyrimidine,
5-amino-2,4,6-tris-(methylamino)-pyrimidine,
2,4,5-triamino-6-(di-n-propylamino)-pyrimidine,
2,4,5-triamino-6-morpholino-pyrimidine,
2,4,5-triamino-4-(dimethylamino)-pyrimidine
4,5,6-triamino-2-morpholino-pyrimidine,
2,4,5-triamino-6-(β -hydroxyethyl-amino)-pyrimidine,
4,5,6-triamino-2-[(β -aminoethyl)amino]-pyrimidine,
2,5,6-triamino-4-[(β -methylamino)-ethylamino]-pyrimidine,
2,5-diamino-4,6[bis-(γ -diethylamino)-propylamino]-pyrimidine,
4,5-diamino-6-[[β]- β -hydroxyethyl)-amino]-2-(methylamino)-pyrimidine,
5-amino-2,4,6-(triethylamino)-pyrimidine, and
5-amino-6-anilino-2,4-[bis-(β -hydroxyethyl)-amino]-pyrimidine.

Generally speaking, the coupling components include 1-phenyl-pyrazol-5-ones, aromatic amines, aromatic alcohols, preferably m-aminophenols, 1,3-diamino-4-nitrobenzenes, 1,4-diamino-2-nitrobenzenes and pyrazolidine-diones.

Examples of coupling components include 1-phenyl-pyrazol-5-one derivatives of the formula



wherein D represents lower alkyl such as methyl, NH_2- , $-NH-CO-D_1$ or $-NH-CO-NH-D_1$ or $NH-CS-NH-D_1$ and D_1 denotes a hydrocarbon radical with 1 to 12 carbon atoms or a heterocyclic radical, and preferably D_1 is alkyl having 1 to 6 carbon atoms such as methyl, ethyl, isopropyl, n-propyl, n-butyl; phenyl; halopenyl such as p-chlorophenyl; alkyl-phenyl having from 7 to 12 carbon atoms such as p-methylphenyl, p-ethylphenyl, o-propylphenyl; alkoxyphe-
nyl, having from 7 to 12 carbon atoms such as o-methoxyphenyl p-isopropoxyphenyl; dialkylamino-

phenyl having 1 to 4 carbon atoms in the alkyl such as p-dimethylaminophenyl; cyclohexyl; alkylcyclohexyl having from 7 to 12 carbon atoms such as methylcyclohexyl; pyridyl; and piperidyl.

Examples of readily available substituted 1-phenyl-3-aminopyrazol-5-ones are the following:

- 1-phenyl-3-amino-pyrazol-5-one,
- 1-phenyl-3-methyl-pyrazol-5-one,
- 1-phenyl-3-acetamido-pyrazol-5-one,
- 1-phenyl-3-benzamido-pyrazol-5-one,
- 1-phenyl-3-(3'-cyclohexylureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-phenylureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-p-chlorophenylureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-ethyl-2'-thioureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-n-butyl-2'-thioureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-phenyl-2'-thioureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-methylureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-ethylureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-n-propylureido)-pyrazol-5-one,
- 1-phenyl-3-(3'-isopropylureido)-pyrazol-5-one, and
- 1-phenyl-3-(3'-n-butylureido)-pyrazol-5-one.

Another coupling component for the hair dyes is based on aromatic amines and/or aromatic alcohols, preferably a compound having the formula



in which Ar represents an aromatic nucleus, preferably consisting of naphthylene, hydroxynaphthylene, aminonaphthylene, phenylene, aminophenylene, hydroxyphenylene, toluylene, alkoxyphenylene having from 7 to 10 carbon atoms and quinolyne; E represents hydrogen, hydroxyl, or amino, dialkylamino having from 1 to 4 carbon atoms in the alkyls; and F represents hydroxyl, amino or dialkylamino having from 1 to 4 carbon atoms in the alkyls.


Suitable coupling components include aromatic amines and diamines, phenols, naphthols and aminophenols. The metacomponents are preferably used in the case of the diamines, aminophenols and phenols. When diamines are used, those in which the hydrogen atoms of the amino groups are substituted by lower alkyl residues (C₁-C₄) may also be used.

The following compounds are examples of the said coupling components:


- m-cresol,
- m-phenylenediamine,
- o-cresol,
- m-aminophenol,
- anisidine,
- 2,4-diaminoanisole,
- m-toluylenediamine,
- resorcinol,
- pyrogallol,
- pyrocatechol,
- resorcinol-monoethyl ether
- resorcinol-monomethyl ether,
- m-aminoresorcinol,
- 1,5-dihydroxynaphthalene,
- 1,6-dihydroxynaphthalene,
- 1,7-dihydroxynaphthalene,
- 2,7-dihydroxynaphthalene,
- 1,5-amino- or 1,8-amino-hydroxynaphthalene,
- α -naphthol,
- 7-(dimethylamino)-4-hydroxy-1-methyl-2-quinoline,
- 8-hydroxyquinoline,
- 1,8-diaminonaphthalene,
- 2,6-dimethyl-phenol,

- 2,5-dimethyl-phenol,
- 3,4-dimethyl-phenol,
- 3,5-dimethyl-phenol,
- 5-amino-2-methyl-phenol,
- hydroquinone,
- 4-aminophenol,

Other coupling compounds include 1,3-diamino-4-nitrobenzene derivatives of the formula



and 1,4-diamino-2-nitrobenzene derivatives of the formula



in which X is an electrophilic substituent and A₂ is a member selected from the group consisting of hydrogen, alkyl of 1 to 10 carbon atoms, hydroxyalkyl of 1 to 10 carbon atoms; N,N-dialkyl-aminoalkyl of 3 to 18 carbon atoms, alkanoyl of 2 to 10 carbon atoms, substituted alkanoyl of 2 to 10 carbon atoms with a substituent selected from the group consisting of nitro, phenyl, halo, cyano, carboxy and sulfo, acyl of aromatic hydrocarbon carboxylic acids having 7 to 15 carbon atoms, substituted acyl of aromatic hydrocarbon carboxylic acids having from 7 to 15 carbon atoms with a substituent selected from the group consisting of lower alkyl, nitro, halo, cyano, carboxyl and sulfo, and the activated methylene group —CO—CH₂—X, in which X is an electrophilic substituent.

An electrophilic substituents X, the radicals containing a carbonyl, such as carboxyl, alkoxy-carbonyl, and acyl, or a nitrile, or carbonyl containing radicals further substituted with halogen, the sulfo group and the nitro group, are considered. X is preferably a nitrile, acyl, or alkoxy-carbonyl group.

Examples of X include carboxyl, cyano, alkanoyl of 2 to 10 carbon atoms, substituted alkanoyl of 2 to 10 carbon atoms with a substituent selected from the group consisting of nitro, phenyl, halo, cyano, carboxyl and sulfo, acyl of aromatic hydrocarbon carboxylic acids having 7 to 15 carbon atoms, substituted acyl of aromatic hydrocarbon carboxylic acids having from 7 to 15 carbon atoms with a substituent selected from the group consisting of lower alkyl, nitro, halo, cyano, carboxyl and sulfo, cycloalkyl-carbonyl of 6 to 10 carbon atoms, alkoxy-carbonyl of 2 to 10 carbon atoms, cycloalkoxy-carbonyl of 6 to 10 carbon atoms, phenylalkoxy-carbonyl of 8 to 16 carbon atoms, furoyl, and thenoyl.

Suitable examples for compounds of the above described diamino-nitrobenzenes are as follows:

- 1-amino-3-cyanoacetyl-amino-4-nitrobenzene,
- 1-benzoylacetyl-amino-3-amino-4-nitrobenzene,
- 1-methylamino-3-cyanoacetyl-amino-4-nitrobenzene,

1,3-di(cyanoacetyl-amino)-4-nitrobenzene,
 1,3-Di(ω -ethoxycarbonyl-acetyl-amino)-4-nitrobenzene,
 1-Amino-3(ω -ethoxycarbonyl-acetyl-amino)-4-nitrobenzene,
 1-Acetyl-amino-3-cyanoacetyl-amino-4-nitrobenzene,
 1-Amino-3-acetoacetyl-amino-4-nitrobenzene,
 1-(*p*-Nitro-benzylacetyl-amino)-3-amino-4-nitrobenzene,
 1-Trifluoro-acetoacetyl-amino-3-amino-4-nitrobenzene,
 1-(2-Hydroxy-ethyl-amino)-3-cyanoacetyl-amino-4-nitrobenzene,
 1-Amino-2-nitro-4- ω -cyanoacetyl-amino-benzene,
 1-Amino-2-nitro-4- ω -benzoyl-acetyl-amino-benzene,
 1,4-Dicyanoacetyl-amino-2-nitrobenzene,
 1-Amino-2-nitro-4- ω -ethoxycarbonyl-acetyl-amino-benzene,
 1,4-Bis-(ω -ethoxycarbonyl-acetyl-amino)-2-nitrobenzene,
 1-Amino-2-nitro-4- ω -benzoxycarbonyl-acetyl-amino-benzene,
 1-Amino-2-nitro-4-(*p*-nitrobenzoyl-acetyl-amino)-benzene,
 1-Amino-2-nitro-4-acetoacetyl-amino-benzene,
 1-Amino-2-nitro-4-nitroacetoacetyl-amino-benzene,
 1-Amino-2-nitro-4-trifluoroacetoacetyl-amino-benzene,
 1-Amino-2-nitro-4- ω -butyryl-acetyl-amino-benzene,
 1-Amino-2-nitro-4- ω -(β -Naphthoyl-acetyl-amino)-benzene,
 1-Amino-2-nitro-4-cyclohexylcarbonyl-acetyl-amino-benzene,
 1-Amino-2-nitro-4-Furoylacetyl-amino-benzene,
 1-Amino-2-nitro-4-Thenoyl-acetyl-amino-benzene,
 1,4-Bis-(ω -butoxycarbonyl-acetyl-amino)-benzene,
 1,4-Bis-(ω -cyclohexyloxycarbonyl-acetyl-amino)-benzene.

Further coupling components include pyrazolidinones such as 1-phenyl-3,5-pyrazolidine dione.

Specific examples of preferred coupling components to be used for the hair dyes according to the invention are as follows:


α -naphthol,
o-cresol,
m-cresol,
 2,6-dimethylphenol,
 2,5-dimethylphenol
 3,4-dimethylphenol,
 3,5-dimethylphenol
 pyrocatechol,
 pyrogallol
 1,5-dihydroxy-naphthalene,
 1,7-dihydroxy-naphthalene,
 5-amino-2-methylphenol,
 hydroquinone,
 2,4-diaminoanisole,
m-toluylenediamine
 4-aminophenol,
 resorcinol,
 resorcinol monomethyl ether *m*-phenylenediamine,
 3-methyl-1-phenyl-pyrazolone-5,
 3-amino-1-phenyl-pyrazolone-5,
 1-phenyl-3,5-dione-pyrazolidine,
 7-(dimethylamino)-4-hydroxy-1-methyl-quinolone-2,
 1-amino-3-(acetacetyl-amino)-4-nitrobenzene,
 1-amino-3-(cyanoacetyl-amino)-4-nitrobenzene.

In order to obtain shades which are as strong as possible and which correspond to natural hair colors to a large extent, it is very important to use a superior blue dye as shade component. In the preparation of natural

5 looking shades with the aid of coupler components for producing blue dyes, there are, however, difficulties encountered with the customary blue-couplers even when the otherwise very satisfactory tetraaminopyrimidines are used as developers.

10 It has now been found that this drawback can be corrected when, in combination with the tetraaminopyrimidines employed as developers, the following *m*-aminophenols are used as the coupling components. These *m*-aminophenols have the formula

15




20

wherein Z and Y are each selected from the group consisting of hydrogen, halogen, hydroxyl, amino, alkyl having 1 to 4 carbon atoms, alkoxy having 1 to 4 carbon atoms, alkylamino having 1 to 4 carbon atoms, dialkylamino having 1 to 4 carbon atoms in each of the alkyls; R₉ is selected from the group consisting of alkyl having 1 to 10 carbon atoms, phenyl optionally substituted, benzyl, methylenecyanamido, propionamido, ureido, thioureido, oxalyl ester having 1 to 4 carbon atoms in the alcohol moiety, M-(CH₂)_m- in which m is an integer from 1 to 4 and M is selected from the group consisting of hydroxyl, halogen, and -NR₁₁R₁₂ in which R₁₁ and R₁₂ are each hydrogen or alkyl having 1 to 4 carbon atoms, and in which R₉ and R₁₀ can together with the nitrogen atom form an optionally substituted five- or six-membered heterocyclic ring which may contain an additional nitrogen atom or an oxygen atom, and R₁₀ is selected from the group consisting of hydrogen and R₉.

More particularly the coupling component is selected from the group consisting of (a) *m*-aminophenol of the

45

50



55 wherein Z and Y are each selected from the group consisting of hydrogen, halogen, hydroxyl, amino, alkyl having 1 to 4 carbon atoms, alkoxy having 1 to 4 carbon atoms, alkylamino having 1 to 4 carbon atoms, dialkylamino having 1 to 4 carbon atoms in each of the alkyls; R₉ is selected from the group consisting of alkyl having 1 to 10 carbon atoms, hydroxyalkyl having 1 to 10 carbon atoms, phenyl, anilino, hydroxyphenyl, benzyl, methylenecyanamido, propionamido, ureido, thioureido, oxalyl ester of an alcohol having 1 to 4

65



wherein m is an integer from 1 to 4 and M is selected from the group consisting of hydroxyl, halogen, and $\text{—NR}_{11}\text{R}_{12}$, in which R_{11} and R_{12} are each hydrogen or alkyl having 1 to 4 carbon atoms and wherein R_{10} is selected from the group consisting of hydrogen and R_9 ,

wherein R_9 and R_{10} can together with the nitrogen atom form a member selected from the group consisting of a five- or six-membered heterocyclic ring optionally containing in the ring oxygen or nitrogen and

b. a water-soluble acid addition salt of (a).

A particularly preferred subgenus of said coupling component is

wherein Z and Y are each selected from the group consisting of hydrogen, methyl, methoxy, amino, chloro, and hydroxyl;

wherein R_9 is selected from the group consisting of methyl, ethyl, octyl, propyl, hydroxyethyl, phenyl, benzyl, anilino, hydroxyphenyl, ureido, thioureido, methylene-cyanamido, and diethylaminoethyl,

wherein R_{10} is selected from the group consisting of hydrogen and R_9 ,

and wherein R_9 and R_{10} together with the nitrogen atom form a member selected from the group consisting of pyrrolidino, morpholino, and piperidino.

The m -aminophenol derivatives to be employed as blue-coupling component can be used either as such or in form of their water-soluble acid addition salts with inorganic or organic acids, such as for example, hydrochloric acid, sulfuric acid, phosphoric acid, acetic acid, propionic acid, lactic acid and citric acid.

Suitable examples of blue-coupling components to be employed according to the invention are as follows:

3-(dimethylamino)-phenol,
 3-(diethylamino)-phenol,
 3-(dioctylamino)-phenol,
 3-(ethylmethylamino)-phenol,
 3-(ethylpropylamino)-phenol,
 3-(phenylethylamino)-phenol,
 3-(phenyloctylamino)-phenol,
 3-(benzylamino)-phenol,
 3-(ethylbenzylamino)-phenol,
 3-anilinophenol,
 3-(methylamino)-phenol,
 3-(ethylamino)-phenol,
 3-(octylamino)-phenol,
 3-(ethylureidoamino)-phenol,
 3-ureidophenol,
 3-(phenylureidoamino)-phenol,
 3-(β -hydroxyethyl amino)-phenol,
 3-(N - β -hydroxyethyl- N -methyl-amino)-phenol,
 3-(thioureidoamino)-phenol,
 N -ethyl- N -(3-hydroxyphenyl)-urea,
 3-(methylenecyanamido)-phenol,
 3-((2-diethylamino)-ethyl)-amino)-phenol,
 3- N -(2-diethylaminoethyl)- N -methyl-aminophenol,
 3-pyrrolidinophenol,
 3-morpholinophenol,
 3-piperidinophenol,
 5-(ethylamino)-2-methylphenol,
 5-(diethylamino)-2-methylphenol
 3-ethylamino)-4-methoxyphenol,
 3-diethylamino)-4-methoxyphenol,
 2-chloro-5-(dimethylamino)-phenol
 3,5-dihydroxy- N,N -dimethylaniline,
 2-amino-5-(diethylamino)-phenol,
 6-chloro-3-(diethylamino)-5-hydroxy-toluene,
 3-anilino-6-chloro-5-hydroxy-toluene,
 5-methyl-3-piperidino-phenol, and

4-chloro-3-pyrrolidino-phenol.

The preparation of the m -aminophenol derivatives to be used as blue-coupling components is already known from the literature. See the article of F. Effenberger et al in Chem. Ber. 103, 1456-62, (1970) and British Pat. No. 974,343. The preparation of several novel derivatives which were not found in the literature is separately described below.

Specific examples of novel m -aminophenols are as follows: 3-(diethylamino)-4-methoxy-phenol, 5-(ethylamino)-2-methyl-phenol and 5-(diethylamino)-2-methyl-phenol, as well as water-soluble acid addition salts thereof.

In the hair coloring preparations according to the invention, substantially equimolar quantities of the developer components are used based on the coupling components employed. Although an equimolar amount is preferred, it is possible to use more or less than molar amounts. Furthermore, the developer component and the coupling component may be used as pure ingredients or as mixtures. Not only can the developer component consist of mixtures of the tetraaminopyrimidines to be used according to the invention, but the coupling component can also consist of mixtures of the above-named coupling components.

In addition, the hair coloring preparations according to the invention can contain admixtures of other customary developing components and, if necessary, can also contain the customary direct dyestuffs in case the latter are needed for obtaining certain shades. From 0% to 5% direct dyestuffs may be employed. As in the case of other oxidation hair dyes, the oxidative coupling, i.e., the development of the dye can in principle be effected by atmospheric oxygen. Moreover, it is preferred to use chemical oxidizing agents. Suitable examples are especially hydrogen peroxide or its products of addition to urea, melamine and sodium borate, as well as mixtures of such hydrogen peroxide addition products with potassium peroxydisulfate.

When the tetraaminopyrimidines according to the invention are used as developer components, they have the advantage that the oxidative coupling with atmospheric oxygen readily produces highly satisfactory hair dyeing results. Thus, damage to the hair by the oxidizing agents, otherwise employed for oxidative coupling, can be prevented.

In the situation that a bleaching of the hair is simultaneously desired, then the concurrent use of chemical oxidizing agents is necessary.

For application, the hair dyes according to the invention are incorporated into suitable aqueous cosmetic preparations, such as creams, emulsions, gels or simple solutions and immediately before application to the hair, one of the above-named oxidizing agents is added. These hair dyeing preparations contain coupling and developing components in amounts of from 0.2% to 5% by weight, preferably from 1% to 3% by weight.

For the preparation of creams, emulsions or gels, the dye components are mixed with the additional ingredients customarily used in such preparations. Such additional ingredients are, for example, wetting agents or emulsifiers of the anionic or nonionic type, such as alkylbenzenesulfonates, higher fatty alcohol sulfates, higher alkylsulfonates, higher fatty acid alkanolamides, ethoxylated fatty alcohols; thickeners such as methyl cellulose, starch, higher fatty alcohols, paraffin oil and higher fatty acids. Furthermore, perfumes and hair-con-

ditioning and grooming agents, such as panthothenic acid and cholesterol may be included.

Effective amounts of the above-named additives are those customarily employed for this purpose. Effective amounts of wetting agents range from 0.5% to 30% by weight, preferably from 1% to 15% by weight; and for thickeners, an effective amount ranges from 0.1% to 25% by weight, preferably from 1% to 15% by weight, based in each case on the total weight of the preparation. As a lower limit for the above additives, a zero percent lower limit is possible, if none of the additive is utilized.

The hair coloring preparations according to the invention can be applied in a weakly acid medium, a neutral medium or especially, in an alkaline medium, preferably at a pH of 8 to 10, regardless whether a solution, an emulsion, a cream, or a gel is employed.

These preparations are applied at a temperature which usually ranges from 15° C. to 40° C. and preferably is room temperature.

After the preparation has been allowed to react for about 30 minutes, the hair coloring preparation is removed from the hair to be dyed, by rinsing. Then the hair is washed with a mild shampoo, and finally is dried.

When different developer and coupling components are used, the shades obtainable by use of the hair coloring preparations according to the invention have the advantage of providing an extraordinary variations which extend from ash blond to dark brown and green to violet. The properties of the hair colors produced are excellent as far as fastness to light and washing are concerned, as well as good resistance to abrasion is concerned; and the hair dyes once fixed can be easily removed by means of reducing agents.

The following examples are merely illustrative of the present invention without being deemed limitative in any manner thereof.

EXAMPLES

First, there are examples for the preparation of some novel tetraminopyrimidines which are utilized according to the invention, but have not been previously described in the literature.

EXAMPLE 1

Preparation of

5-Amino-2,4,6-tris-(methylamino)-pyrimidine Sulfate,
 $C_7H_{14}N_6 \cdot H_2SO_4 \cdot 2H_2O$

2,4,6-tris(methylamino)-pyrimidine was prepared according to Winkelmann, J. Pract. Chem. 115, 292 (1927) and 5.5 gm of this compound were dissolved in 50 ml water. The solution was adjusted to a pH of 4 by means of sodium acetate and was then heated to 80° C.; and a solution of 1.4 gm of $NaNO_2$ in 5 ml of H_2O was added to form a red solution. At 60° C., sodium dithionite was added until the solution became yellow. Dilute H_2SO_4 was added to the yellow solution; and the precipitate was vacuum filtered. The yield was 55%; and the melting point was 215° C.

Elemental Analysis:	% C	% H	% N
Found:	26.6	6.4	26.6
Calculated:	26.6	8.3	27.7

EXAMPLE 2

Preparation of

2,4,5-Triamino-6-(di-n-propylamino)-pyrimidine
 Dihydrochloride, $C_{10}H_{20}H_6 \cdot 2 HCl$

2,4,5-Triamino-6-(di-n-propylamino)pyrimidine dihydrochloride was prepared stepwise as follows:

1. 2,4-Diamino-6-(di-n-propylamino)-pyrimidine

2,4-Diamino-6-chloropyrimidine was prepared according to Roth et al, J. Amer. Chem. Soc. 72, 1914 (1950). To 15 gm of this compound in 130 ml of ethanol, there was added 50 gm of di-n-propylamine, and the mixture was heated in an autoclave at 200° C. for three hours (at an initial gauge pressure of 10 atm. of N_2). After the autoclave had cooled and had been opened, the reaction mixture was cooled in an ice-salt bath to precipitate dipropylamine hydrochloride. After filtration, the mother liquor was considerably concentrated to give a residue of about 30 ml, whereby 18 gm or 82.9% of theory of crude raw product precipitated. This product was used as such for subsequent synthesis.

2. 2,4-Diamino-6-(di-n-propylamino)-5-nitrosopyrimidine

18 gm of the crude product 2,4-diamino-6-(di-n-propylamino)-pyrimidine was suspended in 25 ml of water; and glacial acetic acid was added until a pH of 4 was reached. After heating to 50° C. had resulted in solution of the substance, 5.5 gm of sodium nitrite in 10 ml of water were slowly added. After a short time, a raspberry red precipitate was deposited, vacuum filtered, and dried under vacuum at room temperature. The remainder weighed 9.4 gm which was 46% theory; and had a melting point of 206°-208° C.

Elemental Analysis:	% C	% H	% N
Found:	50.40	7.61	35.27
Calculated:	49.56	7.62	35.50

3. 2,4,5-Triamino-6-(di-n-propylamino)-pyrimidine Dihydrochloride

6.5 gm of 2,4-diamino-6-(di-n-propylamino)-5-nitrosopyrimidine in 150 ml of ethanol with 0.5 gm of catalyst (10% palladium-on-charcoal) were introduced into a pressure-tested hydrogenation vessel to be hydrogenated at room temperature in a catalytic apparatus of the shaker type. After H_2 -absorption had terminated, the catalyst was removed by filtration; the solution was acidified with hydrochloric acid, and concentrated. The residue was 5.6 gm which was 78.8% of theory of brown crystals; melting point (decomp.) was 105° C. The mass spectrum showed the molecular mass to be 224 (calcd.: 224).

EXAMPLE 3

Preparation of

2,4,5-Triamino-6-morpholino-pyrimidine Sulfate,
 $C_8H_{14}N_6O \cdot H_2SO_4$

2,4,5-Triamino-6-morpholino-pyrimidine sulfate was prepared stepwise from the following compounds:

1. 2,4-Diamino-6-morpholinopyrimidine

2,4-diamino-6-chloropyrimidine was prepared according to Roth et al, J. Amer. Chem. Soc. 72, 1914 (1950). To 10 gm of this compound, (0.07 mol), 30 gm of morpholine (30 ml, 0.34 mol) were added; and while

stirring, the mixture was heated to 100° C. within one hour. The mixture was kept at this temperature for 2½ hours. Then, 10 ml of ethanol were added; and the mixture was stored in a refrigerator until morpholine hydrochloride had precipitated. After removal of this salt, the filtrate was concentrated to produce the crude product which was half oily, half crystalline residue. There was 5.9 gm which constituted 43.7% of theory; and this crude product was utilized for subsequent synthesis.

2. 2,4-Diamino-6-morpholine-5-nitroso-pyrimidine

5.9 gm of the crude 2,4-diamino-6-morpholino-pyrimidine (0.03 mol), were dissolved by heating in 25 ml of water; and glacial acetic acid was added until a pH of 4 was reached. Then the solution was heated to 80° C., and a solution of 2 gm of sodium nitrite in 5 ml of water was slowly added. After a short time, the nitroso compound was obtained as a raspberry red precipitate. Concentration and cooling produced a total yield of a 4.1 gm which was 60.3% of theory. The melting point was 231°-233° C.

Elemental Analysis:	% C	% H	% N
Found:	42.85	5.39	37.48
Calculated:	41.92	4.97	38.22

3. 2,4,5-Triamino-6-morpholino-pyrimidine Sulfate

2.5 gm of 2,4-diamino-6-morpholino-5-nitroso-pyrimidine was suspended in 15 ml of water, and 2 N HCl (5 ml) was added until the substance just dissolved. The mixture was heated to 50° C.; and sodium dithionite (Na₂S₂O₄) was added until the violet color solution took on a yellow color. Then, the solution was filtered, cooled, and adjusted to a pH 2 by means of sulfuric acid (1:1). After a short time, the pyrimidine derivative precipitated as pyrimidine sulfate; the yield was 2.6 gm which was 76.4% of theory.

Elemental Analysis (of the recrystallized compound):	% C	% H	% N
Found:	31.17	5.23	27.2
Calculated:	29.81	4.98	27.8

Melting Point: the compound sintered at 230° slow decomposition from 255° C. on.

Melting Point: the compound sintered at 230° slow decomposition from 255° C. on.

The other tetraaminopyrimidines which were used in the following Examples were reported in the literature and were prepared by the synthesis indicated in the monograph of D. J. Brown, "The Pyrimidines" in the series of Heterocyclic Compounds, Interscience Publishers, 1962, Vols. I and II.

Several m-aminophenol derivatives which were used as blue-coupling components in the hair coloring preparations according to the invention are novel and are not found in the literature. Hence, their preparation is described in the following examples.

EXAMPLE 4

3-(Diethylamino)-4-methoxyphenol Hydrochloride

8.3 gm ethyl bromide were added to a solution of 10.5 gm of 3-amino-4-methoxyphenol in 50 ml of ethanol. The reaction mixture was refluxed for 24 hours. After the solution had cooled, it was diluted with 800 ml of water; and the pH was adjusted to slight alkalinity by

the addition of sodium carbonate. The solution was repeatedly extracted with ether, and the combined extracts were dried over anhydrous sodium sulfate. After the precipitate had been filtered off, the residue was concentrated; and the remaining residue as dissolved in 50 ml of ethanol. Then, after 8.7 gm of ethyl bromide had been added, the alkylation and treatment of the product were repeated as described above with the exception that this time, the dried ether extract was only concentrated to a volume of about 300 ml and dry HCl gas was passed into the solution while it was being cooled with ice. The precipitate was vacuum filtered; the residue was dried, dissolved in ethanol, and treated with active charcoal. Addition of ether to the ethanolic solution resulted in the precipitation of the pure product, which had a melting point of 197°-198° C.

Elemental Analysis:	% C	% H	% N	Cl
Found:	57.1	7.8	6.03	15.3
Calculated:	56.3	7.8	5.4	14.6

Mass Spectrum: 195 (calcd.: 195).

EXAMPLE 5

5-(Ethylamino)-2-methylphenol

88.6 gm of ethyl bromide were added to a solution of 50 gm of 5-amino-2-methylphenol in 250 ml of ethanol; and the mixture was refluxed for four hours. After the solution had cooled, it was diluted with 1.2 liters of water and was made weakly alkaline with sodium carbonate. After repeated extraction with ether, the ether extracts were dried and concentrated. The residue was recrystallized from ethanol, and had a melting point of 128° C. to 129° C.

Elemental Analysis:	% C	% H	% N
Found:	71.5	8.6	9.3
Calculated:	70.79	8.51	9.5

Mass Spectrum: 151 (calcd.: 151)

EXAMPLE 6

5-(Diethylamino)-2-methylphenol Hydrochloride

5 gm of 5-(ethylamino)-2-methylphenol were dissolved in 50 ml of ethanol, then alkylated analogously to the procedure described in Example 4 with 5.2 gm ethyl bromide, and worked up. The hydrochloride had a melting point of 192° C.

Elemental Analysis:	% C	% H	% N	% Cl
Found:	61.3	8.4	6.5	16.5
Calculated:	62.9	8.8	6.8	15.9

Mass Spectrum: 179 (calcd.: 179)

The hair dyes according to the invention were applied in form of an aqueous preparation such as an emulsified cream. The emulsion contained

10 parts by weight of fatty alcohols having 12 to 18 carbon atoms,

10 parts by weight of fatty alcohol sulfate (sodium salt) having 12 to 18 carbon atoms and

75 parts by weight of water.

Into each emulsion, there was incorporated 0.01 mol of the tetraaminopyrimidines and couplers which are listed in the following Table I. Then, the pH-value of the emulsion was adjusted to 9.5 with ammonia, and the emulsion was made up to 100 parts by weight with water. The oxidative coupling was effected by using as an oxidizing agent either atmospheric oxygen, or a 1% hydrogen peroxide solution with the proviso that 10 parts by weight of hydrogen peroxide solution were

added to 100 parts by weight of the emulsion. The respective dyeing cream, with or without additional oxidizing agent, was applied to human hair that was 90% gray and that had not been pretreated in a special manner. After the cream had remained on the hair for 30 minutes to complete the dyeing process, the hair was washed with a customary shampoo and then dried. The shades thereby obtained are also listed in the following Table I.

TABLE 1

Example	Developer	Coupler	Shade obtained	
			With Atmospheric O ₂	with 1% H ₂ O ₂
7	2,4,5,6-Tetra-amino-pyrimidine	m-Phenylenediamine	olive	olive
8	2,4,5,6-Tetra-amino-pyrimidine	2,4-Diaminoanisole	dark green	dark green
9	2,4,5,6-Tetra-amino-pyrimidine	m-diamino toluene	yellowish	yellowish
10	2,4,5,6-Tetra-amino-pyrimidine	m-aminophenol	brown violet	brown violet
11	2,4,5,6-Tetra-amino-pyrimidine	Resorcinol	brown strawberry red	brown grayish red
12	2,4,5,6-Tetra-amino-pyrimidine	3-Amino-1-phenyl-pyrazolone-5	brownish orange	brownish orange
13	2,4,5,6-Tetra-amino-pyrimidine	3-Methyl-1-phenyl-pyrazolone-5	brownish red	brownish orange
14	2,4,5,6-Tetra-amino-pyrimidine	Resorcinol monomethyl ether	golden brown	golden brown
15	2,4,5,6-Tetra-amino-pyrimidine	Naphthol	yellowish brown	yellowish brown
16	2,4,5,6-Tetra-amino-pyrimidine	1,5-Dihydroxy-Naphthalene	havanna brown	havanna brown
17	2,4,5,6-Tetra-amino-pyrimidine	1,7-Dihydroxy-Naphthalene	olive brown	olive brown
18	4-Dimethylamino-2,5,6-Triamino-pyrimidine	m-Phenylenediamine	yellowish brown	yellowish brown
19	4-Dimethylamino-2,5,6-Triamino-pyrimidine	2,4-Diaminoanisole	olive yellow	olive yellow
20	4-Dimethylamino-2,5,6-Triamino-pyrimidine	m-Diaminotoluene	yellowish brown	yellowish brown
21	4-Dimethylamino-2,5,6-Triamino-pyrimidine	m-Aminophenol	burgundy red	burgundy red
22	4-Dimethylamino-2,5,6-Triamino-pyrimidine	Resorcinol	brownish red	brownish red
23	4-Dimethylamino-2,5,6-Triamino-pyrimidine	3-Amino-1-Phenyl-pyrazolone-5	raspberry red	raspberry red
24	4-Dimethylamino-2,5,6-Triamino-pyrimidine	m-Phenylenediamine	olive brown	olive brown
25	2-Dimethylamino-4,5,6-triamino-pyrimidine	m-Phenylenediamine	dark green	dark green
26	2-Dimethylamino-4,5,6-triamino-pyrimidine	2,4-Diaminoanisole	dark green	dark green
27	2-Dimethylamino-4,5,6-triamino-pyrimidine	m-Diaminotoluene	olive-tinged yellow	olive-tinged yellow
28	2-Dimethylamino-4,5,6-triamino-pyrimidine	m-Aminophenol	grayish violet	dark violet
29	2-Dimethylamino-4,5,6-triamino-pyrimidine	Resorcinol	reddish violet	reddish violet
30	2-Dimethylamino-4,5,6-triamino-pyrimidine	3-Amino-1-phenyl-pyrazolone-5	brick red	brick red
31	2-Dimethylamino-4,5,6-triamino-pyrimidine	3-Methyl-1-phenyl-pyrazolone-5	lake red	grayish orange
32	2-Dimethylamino-	Naphthol	hair	olive

TABLE 1-continued

Example	Developer	Coupler	Shade obtained	
			With Atmospheric O ₂	with 1% H ₂ O ₂
	4,5,6-triamino-pyrimidine		brown	brown
33	2-Dimethylamino-4,5,6-triamino-pyrimidine	3-acetoacetyl-amino-1-amino-4-nitrobenzene	yellowish brown	brass yellow
34	2-Dimethylamino-4,5,6-triamino-pyrimidine	1-Phenyl-3,5-pyrazolidine-dione	brownish orange	gray red
35	2-Dimethylamino-4,5,6-triamino-pyrimidine	o-Cresol	brownish orange	ivory
36	2-Dimethylamino-4,5,6-triamino-pyrimidine	m-Cresol	brownish orange	ivory
37	2-Dimethylamino-4,5,6-triamino-pyrimidine	2,5-Dimethylphenol	brownish red	grayish green
38	2-Dimethylamino-4,5,6-triamino-pyrimidine	3,4-Dimethylphenol	brownish orange	grayish green
39	2-Dimethylamino-4,5,6-triamino-pyrimidine	3,5-Dimethylphenol	brownish orange	grayish orange
40	2-Dimethylamino-4,5,6-triamino-pyrimidine	1,5-Dihydroxy-Naphthalene	brown	brown
41	2-Dimethylamino-4,5,6-triamino-pyrimidine	Pyrogallol	chocolate brown	zinc gray
42	2-Dimethylamino-4,5,6-triamino-pyrimidine	Pyrocatechol	reddish brown	fallow
43	2-Dimethylamino-4,5,6-triamino-pyrimidine	7-(Dimethylamino)-4-hydroxy-1-methyl-2-quinolone	grayish orange	golden blond
44	2-Dimethylamino-4,5,6-triamino-pyrimidine	5-Amino-2-methylphenol	brown	brown
45	2-Dimethylamino-4,5,6-triamino-pyrimidine	Hydroquinone	brownish red	golden blond
46	2-Piperidino-4,5,6-triamino-pyrimidine	m-Phenylenediamine	dark green	dark green
47	2-Piperidino-4,5,6-triamino-pyrimidine	m-Aminophenol	dark violet	dark violet
48	2-Piperidino-4,5,6-triamino-pyrimidine	Resorcinol	grayish ruby	grayish ruby
49	2-Piperidino-4,5,6-triamino-pyrimidine	2,4-Diaminoanisole	dark green	dark green
50	2-Piperidino-4,5,6-triamino-pyrimidine	m-Diaminotoluene	brownish orange	brownish orange
51	2-Piperidino-4,5,6-triamino-pyrimidine	3-Amino-1-phenylpyrazolone-5-	tomato red	tomato red
52	2-Morpholino-4,5,6-triamino-pyrimidine	m-Phenylenediamine	olive	olive
53	2-Morpholino-4,5,6-triamino-pyrimidine	m-Aminophenol	dark violet	dark violet
54	2-Morpholino-4,5,6-triamino-pyrimidine	Resorcinol	grayish ruby	grayish ruby
55	2-Morpholino-4,5,6-triamino-pyrimidine	2,4-Diaminoanisole	dark green	dark green
56	2-Morpholino-4,5,6-triamino-pyrimidine	m-Diaminotoluene	olive brown	olive brown
57	2-Morpholino-4,5,6-triamino-pyrimidine	3-Amino-1-phenylpyrazolone-(5)	brownish red	brownish red
58	2-Methylamino-	m-Phenylenediamine	olive	olive

TABLE 1-continued

Example	Developer	Coupler	Shade obtained	
			With Atmospheric O ₂	with 1% H ₂ O ₂
59	4,5,6-triamino-pyrimidine 2-Methylamino-4,5,6-triamino-pyrimidine	m-Aminophenol	dark purple	dark violet
60	2-Methylamino-4,5,6-triamino-pyrimidine	Resorcinol	grayish red	brownish violet
61	2-Methylamino-4,5,6-triamino-pyrimidine	2,4-Diaminoanisole	dark green	dark green
62	2-Methylamino-4,5,6-triamino-pyrimidine	m-Diaminotoluene	yellow	yellow
63	2-Methylamino-4,5,6-triamino-pyrimidine	3-Amino-1-phenyl-pyrazolone-5	brownish red	brownish red
64	6-Morpholino-2,4,5-triamino-pyrimidine	m-Phenylenediamine	olive brown	olive brown
65	6-Morpholino-2,4,5-triamino-pyrimidine	2,4-Diaminoanisole	grayish green	olive
66	6-Morpholino-2,4,5-triamino-pyrimidine	m-Diaminotoluene	golden yellow	brass yellow
67	6-Morpholino-2,4,5-triamino-pyrimidine	m-Aminophenol	grayish ruby	grayish red
68	6-Morpholino-2,4,5-triamino-pyrimidine	Resorcinol	brownish red	brownish red
69	6-Morpholino-2,4,5-triamino-pyrimidine	3-Amino-1-phenyl-pyrazolone-5	dull red	dull red
70	6-Piperidino-2,4,5-Triamino-pyrimidine	m-Phenylenediamine	bamboo yellow	grayish yellow
71	6-Piperidino-2,4,5-Triamino-pyrimidine	2,4-Diaminoanisole	olive brown	olive
72	6-Piperidino-2,4,5-Triamino-pyrimidine	m-Aminophenol	olive brown	bamboo yellow
73	6-Piperidino-2,4,5-Triamino-pyrimidine	Resorcinol	grayish red	ivory
74	6-Piperidino-2,4,5-Triamino-pyrimidine	m-Diaminotoluene	grayish yellow	olive brown
75	6-Piperidino-2,4,6-Triamino-pyrimidine	3-Amino-1-phenyl-pyrazolone-5	chamois yellow	straw yellow
76	6-(Di-n-propylamino)-2,4,5-triamino-pyrimidine	m-Phenylenediamine	honey yellow	olive
77	6-(Di-n-propylamino)-2,4,5-triamino-pyrimidine	2,4-Diaminoanisole	olive green	olive
78	6-(Di-n-propylamino)-2,4,5-triamino-pyrimidine	m-Diaminotoluene	honey yellow	honey yellow
79	6-(Di-n-propylamino)-2,4,5-triamino-pyrimidine	m-Aminophenol	grayish ruby	grayish ruby
80	6-(Di-n-propylamino)-2,4,5-triamino-pyrimidine	Resorcinol	brownish red	brownish red
81	6-(Di-n-propylamino)-2,4,5-triamino-pyrimidine	3-Amino-1-phenyl-pyrazolone-5-	reddish brown	reddish brown
82	5-Amino-2,4,6-tris-(methylamino)-pyrimidine	m-Phenylenediamine	oak brown	olive
83	5-Amino-2,4,6-tris-(methylamino)-pyrimidine	2,4-Diaminoanisole	olive	beaver brown
84	5-Amino-2,4,6-	m-diaminotoluene	brass	olive

TABLE 1-continued

Example	Developer	Coupler	Shade obtained	
			With Atmospheric O ₂	with 1% H ₂ O ₂
	tris-(methylamino)-pyrimidine		yellow	
85	5-Amino-2,4,6-tris-(methylamino)-pyrimidine	m-Aminophenol	dark ruby	brown
86	5-Amino-2,4,6-tris-(methylamino)-pyrimidine	Resorcinol	grayish red	dull red
87	5-Amino-2,4,6-tris-(methylamino)-pyrimidine	3-Amino-1-phenyl pyrazolone-5	reddish brown	light brown
88	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-(Diethylamino)-phenol	blue black	blue black
89	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-[(β -Hydroxyethyl)-amino]-phenol	blue black	blue black
90	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-[2-(Diethylamino)-ethyl-amino]-phenol	blue black	blue black
91	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-Morpholino-phenol	dark violet	dark violet
92	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-(Ethylpropylamino) phenol	blue black	blue black
93	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-(Butylamino)-phenol	blue black	blue black
94	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-(Ethylamino)-phenol	blue black	blue black
95	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-(Octylamino)-phenol	dark violet	dark violet
96	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-Pyrrolidino-phenol	blue black	blue black
97	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-Piperidino-phenol	dark blue	dark blue
98	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-(Dimethylamino)-phenol	dark blue	dark blue
99	2-Dimethylamino-4,5,6-Triamino-pyrimidine	5-(Ethylamino)-2-methyl-phenol	bluish gray	bluish gray
100	2-Dimethylamino-4,5,6-Triamino-pyrimidine	2-Chloro-5-(dimethylamino)-phenol	dark violet	dark violet
101	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-Hydroxydiphenylamine	dark blue	dark blue
102	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-(Methylenecyanamido)-phenol	dark violet	dark violet
103	2-Dimethylamino-4,5,6-Triamino-pyrimidine	2-bromo-5-(methylene-cyanamido)-phenol	blue black	blue black
104	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-Tioureido-phenol	brownish gray	dull violet
105	2-Dimethylamino-4,5,6-Triamino-pyrimidine	3-(Diethylamino)-4-methoxy-phenol	dark blue	dark blue
106	2-Dimethylamino-4,5,6-Triamino-pyrimidine	5-(Diethylamino)-2-methyl-phenol	dark blue	dark blue
107	2-Morpholino-4,5,6-triamino-pyrimidine	m-Dimethylamino-phenol	dark blue	dark blue
108	2-Morpholino-4,5,6-triamino-pyrimidine	3-Hydroxydiphenylamine	dark blue	dark blue
109	2-Morpholino-4,5,6-triamino-pyrimidine	3-(Octylamino)-phenol	dark violet	violet blue
110	2-Piperidino-	m-(Dimethylamino)-	dark	dark

TABLE 1-continued

Example	Developer	Coupler	Shade obtained	
			With Atmospheric O ₂	with 1% H ₂ O ₂
111	4,5,6-triamino-pyrimidine	phenol	blue	blue
	2-Piperidino-4,5,6-triamino-pyrimidine	m-(Octylamino)-phenol	blue violet	blue violet
112	2-Piperidino-4,5,6-triamino-pyrimidine	3-Hydroxydiphenylamine	dark blue	dark blue
113	2-Methylamino-4,5,6-triamino-pyrimidine	3-Hydroxydiphenylamine	violet blue	blue violet
114	2-Methylamino-4,5,6-triamino-pyrimidine	3-(Octylamino)-phenol	deep violet	deep violet
115	2-Methylamino-4,5,6-triamino-pyrimidine	m-(Dimethylamino)-phenol	deep violet	violet blue
116	2,4,5,6-Tetraamino-pyrimidine	m-(Dimethylamino)-phenol	dark violet	dark violet
117	2,4,5,6-Tetraamino-pyrimidine	m-(Diethylamino)-phenol	dark violet	dark violet
118	2,4,5,6-Tetraamino-pyrimidine	m-(Ethylamino)-phenol	indigo blue	indigo blue
119	2,4,5,6-Tetraamino-pyrimidine	m-(Ethylpropylamino)-phenol	dark violet	dark violet
120	2,4,5,6-Tetraamino-pyrimidine	m-[(β-Hydroxyethyl)-amino]-phenol	dark violet	dark violet
121	2,4,5,6-Tetraamino-pyrimidine	m-Piperidino-phenol	light violet	light violet
122	2,4,5,6-Tetraamino-pyrimidine	m-Pyrrolidino-phenol	violet	violet
123	2,4,5,6-Tetraamino-pyrimidine	3-Hydroxydiphenylamine	indigo blue	indigo blue

The following tests were conducted to determine the toxicological and dermatological properties of tetraaminopyrimidines. The test compound used was 2-dimethylamino-4,5,6-triamino-pyrimidine sulfate which was compared with known compounds, p-phenylenediamine and p-toluylenediamine sulfate. The following results were obtained.

EXAMPLE 124

Acute Toxicity

Male white mice of the CF/W 68-strain were used for the tests of general tolerance. The average weight of the test animals was 22 gm. The substances to be tested were administered by means of a stomach tube, whereby increasing dosages were used once. Ten mice were employed per dose. The volume of application was constant and was 0.2 cm³/10 gm body weight. The animals were observed for a period of eight days. After the test results had been calculated according to the method of Litchfield-Wilcoxon, J. Pharm. Exptl. Ther. 96, 99-108(1949), the following LD₅₀-values were obtained.

2-dimethylamino-4,5,6-triamino-pyrimidine sulfate	555 mg/kg
p-Phenylenediamine	87 mg/kg
p-Diaminotoluene Sulfate	110 mg/kg

Distilled water was used as the solvent.

EXAMPLE 125

Dermal Tolerance Tests of the Hairless Mouse

Groups of five animals were used for each preparation. Small quantities of the substances to be tested were applied in form of an aqueous 5% solution to the dorsal

skin of each animal. A single daily application was continued for 14 days. During this period, and at the end of the test, none of the animals showed any reaction.

EXAMPLE 126

Mucocutaneous Tolerance of the Rabbit Eye

Groups of albino rabbits were employed for these tests of the local tolerance. Small quantities of the substances to be tested were employed as aqueous 5% preparations; and were once dripped into the conjunctival sac of one eye of each rabbit. The reactions of the mucous membranes of the eye were evaluated according to the point system of Draize, [Appraisal of the Safety of Chemicals in Foods, Drugs and Cosmetics. Assoc. of Food and Drug Officials of the U.S., pp. 49-52 (1959)]. This evaluation which was made 2, 6, 24 and 48 hours after the application and it showed that p-phenylenediamines leads to a low-grade redness and exudation of the conjunctiva which could not be detected 24 hours after the dripping in. The other two test compounds were tolerated without reaction.

EXAMPLE 127

Skin (Tissue) Tolerance Test of White Mice after a Single Intracutaneous Application of Various Concentrations of the Preparations to be Tested


This test of local tolerance was conducted according to Barail, [J. Society Cosmetic Chemists 11, 241 (1960)]. In this test, small quantities of the compounds to be tested were transcutaneously applied to the abdominal skin of white mice, whereby increasing concentration were used. After 24 hours, the test animals were sacri-

25

fixed and the treated places of the skin were cut out and dried. The skin damages were evaluated according to a point system, in which the blood irrigation and other damages of the treated skin are taken into account. In this test, a larger animal group of ten mice was employed per preparation and per test concentration. Hence, finer differences with respect to local tolerance could be observed according to the above-described test procedure. The test results are reported in the following Table II.

5

10



It is evident from the above test results that (2-dimethylamino)-4,5,6-triamino-pyrimidine sulfate shows the best results of the tested developers with respect to general and local tolerance. Besides these good toxicological and dermatological properties, the tetraaminopyrimidines used in the hair coloring preparations according to the invention have the additional advantages that the color can readily be developed with atmospheric oxygen. These hair coloring preparations produce an extraordinary variation of shades which are distinguished by excellent fastness to light and washing, as well as by excellent resistance to abrasion, and they can easily be removed with reducing agents.

Although the present invention has been disclosed in connection with a few preferred embodiments thereof, variations and modifications may be resorted to by those skilled in the art without departing from the principles of the new invention. All of these variations and modifications are considered to be within the true spirit and scope of the present invention as disclosed in the foregoing description and defined by the appended claims.

We claim:

1. An aqueous preparation for the dyeing of hair consisting essentially of (1) from 0.2% to 5% by weight of an oxidation dyestuff combination of a developer component, and a coupling component in substantially equimolar amounts, said developer component consisting essentially of (A) a tetraaminopyrimidine of the formula

wherein R_1 , R_2 , R_3 , R_4 , R_5 and R_6 are each selected from the group consisting of hydrogen, phenyl, 6-alkyl having 1 to 4 carbon atoms, phenylalkyl having 7 to 10 carbon atoms, phenylalkenyl having 7 to 10 carbon atoms,




wherein n is an integer from 1 to 4, and X is selected from the group consisting of hydroxyl, halogen and NR_7R_8- in which R_7 and R_8 are selected from the group consisting of hydrogen and alkyl having 1 to 4 carbon atoms, and together with the nitrogen atom R_7 to R_8 form a 5 to 6 membered heterocyclic ring optionally containing an additional nitrogen atom or an oxygen atom,

and wherein R_1 and R_2 or R_3 and R_4 or R_5 and R_6 , together with the nitrogen atom form a 5 to 6 membered heterocyclic ring, optionally containing another nitrogen or oxygen atom in the ring or (B) a water-soluble acid addition salt of (A); (2) from 0% to 5% by weight of a direct dyestuff; (3) from 0% to 30% by weight of a surfactant; (4) from 0% to 25% by weight of a thickener; and (5) the balance up to 100% by weight of water.

2. The aqueous preparation of claim 1, wherein R_1 , R_2 , R_3 , R_4 , R_5 and R_6 are each selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, n-propyl, butyl, phenyl, benzyl, benzylidene; and $-(CH_2)_n-X$

wherein R_1 and R_2 or R_3 and R_4 , or R_5 and R_6 , together with the nitrogen atom form a substituent selected from the group consisting of piperidino and morpholino; and

26




wherein n is 1,2 or 3 and X is selected from the group consisting of hydroxyl, halogen and —NR₇R₈ in which R₇ and R₈ are selected from the group consisting of hydrogen and alkyl having 1 to 4 carbon atoms.

3. The aqueous preparation of claim 1, wherein said coupling component is selected from the group consisting of (a) a 1-phenyl-pyrazol-5-one, (b) an aromatic amine (c) an aromatic alcohol, and (d) a pyrazolidinedione.

4. The aqueous preparation of claim 1, wherein there is from 1% to 3% by weight of said oxidation dyestuff combination.

5. The aqueous preparation of claim 1, wherein the developer component is selected from the group consisting of (A) 2,4,5,6-tetraamino-pyrimidine, 4-dimethylamino-2,5,6-triaminopyrimidine, 2-dimethylamino-4,5,6-triamino-pyrimidine, 2-piperidino-4,5,6-triamino-pyrimidine, 2-morpholino-4,5,6-triamino-pyrimidine, 6-morpholino-2,4,5-triamino-pyrimidine, 6-piperidino-2,4,5-triamino-pyrimidine, 6-(di-n-propylamino)-2,4,5-triamino-pyrimidine, 5-amino-2,4,6-tris-(methylamino)-pyrimidine and 2-methylamino-4,5,6-triamino-pyrimidine and (B) a water soluble acid addition salt of (A).

6. The aqueous preparation of claim 1, wherein said coupling component is selected from the group consisting of (a) a m-aminophenol of the formula



wherein Z and Y are each selected from the group consisting of hydrogen, halogen, hydroxyl, amino, alkyl having 1 to 4 carbon atoms, alkoxy having 1 to 4 carbon atoms, alkylamino having 1 to 4 carbon atoms, dialkylamino having 1 to 4 carbon atoms in each of the alkyls;

wherein R₉ is selected from the group consisting of alkyl having 1 to 10 carbon atoms, hydroxyalkyl having 1 to 10 carbon atoms, phenyl, anilino, hydroxyphenyl, benzyl, methylenecyanamido, propionamido, ureido, thioureido, oxalyl ester of an alcohol having 1 to 4 carbon atoms,



wherein m is an integer from 1 to 4 and M is selected from the group consisting of hydroxyl, halogen, and —NR₁₁R₁₂, in which R₁₁ and R₁₂ are selected from the group consisting of hydrogen and alkyl having 1 to 4 carbon atoms and wherein R₁₀ is selected from the group consisting of hydrogen and R₉,

wherein R₉ and R₁₀ can together with the nitrogen atom form a five to six membered heterocyclic ring optionally containing an additional nitrogen atom or an oxygen atom and

b. a water-soluble acid addition salt of (a).

7. The aqueous preparation of claim 6, wherein Z and Y are each selected from the group consisting of hydrogen, methyl, methoxy, amino, chloro and hydroxyl;

wherein R₉ is selected from the group consisting of methyl, ethyl, octyl, propyl, hydroxyethyl, phenyl, benzyl, anilino, hydroxyphenol, ureido, thioureido, methylenecyanamido and diethylaminoethyl, wherein R₁₀ is selected from the group consisting of hydrogen and R₉,

and wherein R₉ and R₁₀ together with the nitrogen atom form a member selected from the group consisting of pyrrolidino, morpholino, and piperidino.

8. A process for the dyeing of human hair comprising applying to said hair at temperatures ranging substantially from 15° C. to 40° C. for a time sufficient to effect dyeing, an effective amount of the aqueous preparation of claim 1.

9. The process for the dyeing of human hair of claim 8, wherein said preparation also contains a chemical oxidizing agent.

* * * * *

45

50

55

60

65

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 30199 Reissue

Page 1 of 3

DATED : January 29, 1980

INVENTOR(S) : DAVID ROSE, FERDI SAYGIN, ERWIN WEINRICH


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 3, line 8: Before "hydroxyl" insert the following:
-- piperidino and morpholino; and wherein
n is 1, 2 or 3 and is selected from the
group consisting of --.

Column 14, line 5: "as" should read -- was --.

Column 21/22, Table 1, Example 100, under the column "Coupler":
"amion)-phenol" should read
-- amino)-phenol --.

Column 25: The Table II which reads



UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 30,199 Reissue

Page 2 of 3

DATED : January 29, 1980

INVENTOR(S) : DAVID ROSE, FERDI SAYGIN, ERWIN WEINRICH


It is certified that error appears in the above—identified patent and that said Letters Patent is hereby corrected as shown below:

should read:

-- TABLE II

Skin (Tissue) Test of White Mice after Intracutaneous Application (Test according to Barail) Average Values after Evaluation of 10 Animals Employed Each Time per Concentration

Points or Degree of Skin Damage	4,5,6-Triamino-2-(Dimethylamino)-pyrimidine Sulfate	p-Phenylenediamine	p-Diamino-toluene Sulfate
---------------------------------	---	--------------------	---------------------------



UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 30,199 Reissue


Page 3 of 3

DATED : January 29, 1980

INVENTOR(S) : DAVID ROSE, FERDI SAYGIN, ERWIN WEINRICH

It is certified that error appears in the above—identified patent and that said Letters Patent is hereby corrected as shown below:

TABLE II continued:



Concentration of the
Compounds to be tested in % --.

Signed and Sealed this

Seventh Day of July 1981

[SEAL]

Attest:

RENE D. TEGTMEYER

Attesting Officer

Acting Commissioner of Patents and Trademarks