

# United States Patent [19]

Bayer et al.

[11] E

Patent Number: **Re. 31,731**

[45] **Reissued** Date of Patent: **Nov. 13, 1984**

[54] **HERBICIDAL  
4-TRIFLUOROMETHYL-4-NITRODIPHENYL ETHERS**

[76] Inventors: **Horst O. Bayer**, Levittown; **Colin Swithenbank**, Perkasië; **Roy Y. Yih**, Doylestown, all of Pa.

[21] Appl. No.: **939,291**

[22] Filed: **Sep. 1, 1978**

## Related U.S. Patent Documents

Reissue of:

[64] Patent No.: **3,928,416**  
Issued: **Dec. 23, 1975**  
Appl. No.: **331,719**  
Filed: **Feb. 12, 1973**

U.S. Applications:

[63] Continuation-in-part of Ser. No. 234,651, Mar. 14, 1972, Pat. No. 3,789,276.

[51] Int. Cl.<sup>3</sup> ..... **C07C 79/46**

[52] U.S. Cl. .... **562/435; 71/94; 71/95; 71/105; 71/111; 71/116; 71/118; 71/121; 71/122; 71/124; 260/465 F; 544/163; 544/167; 548/577; 560/21; 560/133; 560/138; 564/50; 564/192; 564/212; 564/221; 568/585; 568/586**

[58] Field of Search ..... **560/21; 562/435**

[56] **References Cited**

## U.S. PATENT DOCUMENTS


3,080,225 3/1963 Wilson et al. .... 71/213  
3,154,398 10/1964 McRae ..... 71/216  
3,322,525 5/1967 Martin et al. .... 71/2.3  
3,423,470 1/1969 Rohr et al. .... 260/612  
3,652,645 3/1972 Theissen ..... 560/21  
3,784,635 1/1974 Theissen ..... 560/21

3,941,830 3/1976 Theissen ..... 560/21  
3,979,437 9/1976 Theissen ..... 560/21

Primary Examiner—Paul J. Killos

[57] **ABSTRACT**

Compounds of the formula



wherein

X is a hydrogen atom, a halogen atom, a trihalomethyl group, an alkyl group, or a cyano group, Y is a hydrogen atom, or a trihalomethyl group, and Z is an alkoxy group, an alkoxyalkoxy group, a hydroxyalkoxy group, an alkyl group, a halogen atom, an alkylamino group, a dialkylamino group, an alkylthio group, a carboxy group, a carbalkoxy group, a carboxyalkoxy group, a carbalkoxyalkoxy group, a carboxyalkyl group, a carbalkoxyalkyl group, a dialkylureido group, an alkanoylamino group, or carbalkoxyamino group,

and compositions containing these compounds exhibit herbicidal activity.

**5 Claims, No Drawings**



# HERBICIDAL 4-TRIFLUOROMETHYL-4-NITRODIPHENYL ETHERS

Matter enclosed in heavy brackets [ ] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.


## CROSS REFERENCE TO RELATED APPLICATION

This application is a continuation-in-part of our co-pending application Ser. No. 234,651, filed Mar. 14, 1972, now U.S. Pat. No. 3,789,276, granted Mar. 19, 1974.

This invention relates to novel compounds which show activity as herbicides, to novel herbicidal compositions which contain these compounds, and to new methods of controlling weeds with these herbicidal compositions.

Certain diphenyl ethers have been shown to be effective weed control agents. However, the herbicidal effectiveness of a given diphenyl ether cannot be predicted from an examination of the substituent groups attached to the phenyl rings in the ether, and often quite closely related compounds will have quite different weed control abilities. Various diphenyl ethers may have overlapping or complementary areas of activity or selectivity, and can thus be useful in combination to control a variety of weeds upon application of a single composition. Furthermore, the diphenyl ethers heretofore disclosed as herbicides are not completely effective. An ideal herbicide should give selective weed control, over the full growing season, with a single administration at low rates of application. It should be able to control all common weeds by killing them as the seed, the germinating seed, the seedling, and the growing plant. At the same time, the herbicide should not be phytotoxic to the crops to which it is applied and should decompose or otherwise be dissipated so as not to poison the soil permanently. The known diphenyl ether herbicides fall short of these ideals, and it would thus be desirable to have new herbicides which show even more selective control of undesirable plants among desirable crop plants or which complement the known diphenyl ethers in activity.

In accordance with the present invention, there is provided a new class of novel diphenyl ethers having the formula



wherein X is a hydrogen atom, a halogen atom, preferably a fluorine atom or a chlorine atom, a trihalomethyl

group, preferably a trifluoromethyl group, a (C<sub>1</sub>-C<sub>4</sub>)alkyl group, preferably a methyl group, or a cyano group,

Y is a hydrogen atom, a halogen atom, preferably a fluorine atom or a chlorine atom, or a trihalomethyl group, preferably a trifluoromethyl group, and

Z is a hydroxy group, an alkoxy group, preferably having 1 to 6 carbon atoms, and most preferably 1 to 4 carbon atoms, an alkyl group, preferably having 1 to 4 carbon atoms, a halogen atom, preferably a chlorine atom or a fluorine atom, an amino group, preferably having up to 6 carbon atoms, an alkylthio group, preferably having 1 to 4 carbon atoms, a cyano group, a carboxy group, a carbalkoxy group, —CO<sub>2</sub>R, preferably having 1 to 4 carbon atoms in the alkoxy moiety, a carboxyalkyl group, —R'CO<sub>2</sub>H, preferably having up to 4 carbon atoms, a carbalkoxyalkyl group, —R'CO<sub>2</sub>R, preferably having up to 6 carbon atoms, an alkanoyloxy group, —OCOR, preferably having up to 4 carbon atoms, optionally substituted with a halogen atom, or a carbamoyloxy group, —OCONH<sub>2</sub>, —OCONHR, or —OCONR<sub>2</sub>, preferably having up to 6 carbon atoms.

In the above definitions of the Z substituent, R represents an alkyl group, and R' represents a divalent alkylene group. The alkyl or alkylene portion of the alkyl-containing X and Z substituents can have either a straight- or branched-chain or a cyclic spatial configuration.

As used in the present specification and claims, the term "alkoxy group" is intended to include both unsubstituted alkoxy groups as well as substituted alkoxy groups which have one or more of the hydrogen atoms replaced by a substituent group. Among the substituted alkoxy groups which Z can represent are alkoxy groups of preferably up to 4 carbon atoms substituted with a halogen atom, a hydroxy group, a (C<sub>1</sub>-C<sub>4</sub>)alkoxy group, a carboxy group, a carbalkoxy group, preferably having up to 4 carbon atoms in the ester alkoxy group, a trihaloalkyl group, preferably a trifluoromethyl group, an alkenyl group, an alkynyl group, preferably an ethynyl group, an amino group, an alkyl- or dialkyl-amino group, including heterocyclic substituents such as morpholino, piperazino, piperidino, and the like, and preferably having a total of up to 4 carbon atoms, an alkylthio group, preferably having up to 4 carbon atoms, a alkylsulfonyl group, preferably having up to 4 carbon atoms, an epoxy group, an alkylcarbonyl group, including halo-substituted alkylcarbonyl, and preferably having up to 4 carbon atoms in the alkyl group, most preferably methylcarbonyl, a carbamoyl group, including alkyl- or dialkylcarbamoyl, preferably having a total of up to 4 carbon atoms in the alkyl substituents.

The term "amino group" as used in the present specification and claims is intended to include an unsubstituted amino group, —NH<sub>2</sub>, as well as amino groups having one or both hydrogen atoms replaced by substituent groups. Among the substituted amino groups which C can represent are amino groups substituted with one or two alkyl groups, preferably having a total of up to 6 carbon atoms, halo-, hydroxy-, or alkoxy-substituted alkyl groups, preferably having a total of up to 6 carbon atoms, one or two alkylthio carbonyl groups, preferably having a total of up to 4 carbon atoms in the alkyl moiety, carboxy groups, carbalkoxy groups, preferably having up to 4 carbon atoms in the alkoxy group, carbamoyl groups, including alkyl or dialkylcarbamoyl



groups, preferably having up to 4 carbon atoms in the alkyl moiety, alkylcarbonyl groups, preferably having up to 4 carbon atoms, or halo-substituted alkylcarbonyl groups, preferably having up to 4 carbon atoms.

The substituted amino groups can also be heterocyclic amino groups, such as piperidino, piperazino, morpholino, pyrrolidinyl, and the like. When the Z substituent is or contains a carboxy group, either the free acid or the salt form can be used.

When Z is an alkyl group, it may be optionally substituted with a hydroxy group, a (C<sub>1</sub>-C<sub>4</sub>)alkoxy group, or a halogen atom, preferably a chlorine atom.

These novel compounds have been found to show unexpected activity as weed control agents. In a preferred embodiment of the invention, X is a halogen atom or a cyano group, Y is a hydrogen atom or a halogen atom, and Z is an alkoxy group.

Examples of the compounds of the invention embraced by Formula I include:

2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-4-nitro-m-tolyl ether,  
 2-bromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-butyl-4-nitrophenyl ether,  
 2, $\alpha,\alpha,\alpha$ -tetrafluoro-p-tolyl-4-nitro-3-n-propoxyphenyl ether,  
 2-chloro-6, $\alpha,\alpha,\alpha$ -tetrafluoro-p-tolyl-3-methylthio-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-propyl-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether,  
 $\alpha,\alpha,\alpha$ -trifluoro-2-iodo-p-tolyl-3-ethoxy-4-nitrophenyl ether,  
 2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-4-nitro-3-n-propoxyphenyl ether,  
 $\alpha,\alpha,\alpha\alpha',\alpha',\alpha'$ -hexafluoro-2,4-xylyl-3-n-butoxy-4-nitrophenyl ether,  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether,  
 2-chloro-6-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-4-nitro-3-n-propoxyphenyl ether,  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carboxy-4-nitrophenyl ether,  
 2,6-dibromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-methoxymethoxy-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2-hydroxyethoxy)-4-nitrophenyl ether,  
 2, $\alpha,\alpha,\alpha$ -tetrafluoro-p-tolyl-4-nitro-3-n-propylaminophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-dimethylamino-4-nitrophenyl ether,  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carbethoxy-4-nitrophenyl ether,  
 2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carbethoxy-4-nitrophenyl ether,  
 2-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2-carboxy ethoxy)-4-nitrophenyl ether,  
 $\alpha,\alpha,\alpha\alpha',\alpha',\alpha'$ -hexafluoro-2,4-xylyl-3-carbethoxymethyl-4-nitrophenyl ether,  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2-carboxy propyl)-4-nitrophenyl ether,  
 2, $\alpha,\alpha,\alpha$ -tetrafluoro-p-tolyl-3-carbethoxymethoxy-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(3,3-diethylureido)-4-nitrophenyl ether,  
 2-chloro-6-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetamido-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carbethoxyamino-4-nitrophenyl ether,

2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-chloro-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-hydroxy-4-nitrophenyl ether,  
 2-bromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-butynyloxy-4-nitrophenyl ether  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2-methyl)propynyloxy-4-nitrophenyl ether,  
 2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2,2,2-trifluoro)ethoxy-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2-dimethylaminoethoxy)-4-nitrophenyl ether,  
 2-bromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetoxy-4-nitrophenyl ether,  
 $\alpha,\alpha,\alpha\alpha',\alpha',\alpha'$ -hexafluoro-2,4-xylyl-3-(2-hydroxyethylamino)-4-nitrophenyl ether  
 $\alpha^4,\alpha^4,\alpha^4$ -trifluoro-2,4-xylyl-3-amino-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-morpholino-4-nitrophenyl ether,  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(N-methylcarbamoyloxy)-4-nitrophenyl ether,  
 2-chloro-6, $\alpha,\alpha,\alpha$ -tetrafluoro-p-tolyl-3-propionamido-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-chloroacetamido-4-nitrophenyl ether,  
 $\alpha^4,\alpha^4,\alpha^4$ -trifluoro-2,4-xylyl-3-(2,3-epoxypropoxy)-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2,3-dihydroxypropoxy)-4-nitrophenyl ether,  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2-methylthioethoxy)-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-ethyl-3-methylureido)-4-nitrophenyl ether,  
 2-bromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(2-methylsulfonylthoxy)-4-nitrophenyl ether,  
 $\alpha,\alpha,\alpha\alpha',\alpha',\alpha'$ -hexafluoro-2,4-xylyl-3-(3-methylureido)-4-nitrophenyl ether,  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethylthiocarbonylamido-4-nitrophenyl ether,  
 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-cyano-4-nitrophenyl ether,  
 2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carboxy-4-nitrophenyl ether,  
 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carbamoylethoxy)-4-nitrophenyl ether,  
 2-bromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(3-oxobutoxy)-4-nitrophenyl ether,  
 and the like.

The novel diphenyl ethers of the invention are useful both as preemergence and as postemergence herbicides. Preemergence herbicides are ordinarily used to treat the soil in which the desired crop is to be planted by application either before seeding, during seeding, or, as in most applications, after seeding and before the crop emerges. Postemergence herbicides are those which are applied after the plants have emerged and during their growth period.

Among the crops on which the diphenyl ethers of the invention can be advantageously employed are, for example, cotton, soybeans, peanuts, safflower, beans, peas, carrots, corn, wheat, and other cereal crops.

Diphenyl ethers of the invention are useful for controlling weeds in rice crops. When used in transplanted rice crops, the ethers can be applied either preemergence or postemergence to the weeds—that is, they can be applied to the growth medium of the transplanted plants either before the weed plants have emerged or



while they are in their early stages of growth. The ethers can be applied to the growth medium either before or after the rice has been transplanted to that medium.

The diphenyl ethers of the invention can be applied in any amount which will give the required control of weeds. A preferred rate of application of the herbicides of the invention is from about 0.1 to about 12, and most preferably about 0.125 to 4, pounds of the diphenyl ether per acre.

Under some conditions, the diphenyl ethers of the invention may be advantageously incorporated into the soil or other growth medium prior to planting a crop. This incorporation can be carried out by any convenient means, including by simple mixing with the soil, by applying the diphenyl ether to the surface of the soil and then disking or dragging into the soil to the desired depth, or by employing a liquid carrier to accomplish the necessary penetration and impregnation.

A diphenyl ether of the invention can be applied to the growth medium or to plants to be treated either by itself or, as is generally done, as a component in a herbicidal composition or formulation which also comprises an agronomically acceptable carrier. By agronomically acceptable carrier is meant any substance which can be used to dissolve, disperse, or diffuse a herbicidal compound in the composition without impairing the effectiveness of the herbicidal compound and which by itself has no detrimental effect on the soil, equipment, crops, or agronomic environment. Mixtures of the diphenyl ethers of the invention may also be used in any of these herbicidal formulations. The herbicidal compositions of the invention can be either solid or liquid formulations or solutions. For example, the diphenyl ethers can be formulated as wettable powders, emulsifiable concentrates, dusts, granular formulations, aerosols, or flowable emulsion concentrates. In such formulations, the compounds are extended with a liquid or solid carrier and, when desired, suitable surfactants are incorporated.

It is usually desirable, particularly in postemergence applications, to include adjuvants, such as wetting agents, spreading agents, dispersing agents, stickers, adhesives, and the like, in accordance with agricultural practices. Examples of adjuvants which are commonly used in the art can be found in the John W. McCutcheon, Inc. publication "Detergents and Emulsifiers Annual".

The diphenyl ether compounds of this invention can be dissolved in any appropriate solvent. Examples of solvents which are useful in the practice of this invention include alcohols, ketones, aromatic hydrocarbons, halogenated hydrocarbons, dimethylformamide, dioxane, dimethyl sulfoxide, and the like. Mixtures of these solvents can also be used. The concentration of the solution can vary from about 2% to about 98% with a preferred range being about 25% to about 75%.

For the preparation of emulsifiable concentrates, the diphenyl ether can be dissolved in organic solvents, such as benzene, toluene, xylene, methylated naphthalene, corn oil, pine oil, o-dichlorobenzene, isophorone, cyclohexanone, methyl oleate, and the like, or in mixtures of these solvents, together with an emulsifying agent which permits dispersion in water. Suitable emulsifiers include, for example, the ethylene oxide derivatives of alkylphenols or long-chain alcohols, mercaptans, carboxylic acids, and reactive amines and partially esterified polyhydric alcohols. Solvent-soluble sulfates

or sulfonates, such as the alkaline earth salts or amine salts of alkylbenzenesulfonates and the fatty alcohol sodium sulfates, having surface-active properties can be used as emulsifiers either alone or in conjunction with an ethylene oxide reaction product. Flowable emulsion concentrates are formulated similarly to the emulsifiable concentrates and include, in addition to the above components, water and a stabilizing agent such as a water-soluble cellulose derivative or a water-soluble salt of a polyacrylic acid. The concentration of the active ingredient in emulsifiable concentrates is usually about 10% to 60% and in flowable emulsion concentrates, this can be as high as about 75%.

Wettable powders suitable for spraying, can be prepared by admixing the compound with a finely divided solid, such as clays, inorganic silicates and carbonates, and silicas and incorporating wetting agents, sticking agents, and/or dispersing agents in such mixtures. The concentration of active ingredients in such formulations is usually in the range of about 20% to 98%, preferably about 40% to 75%. A dispersing agent can constitute about 0.5% to about 3% of the composition, and a wetting agent can constitute from about 0.1% to about 5% of the composition.

Dusts can be prepared by mixing the compounds of the invention with finely divided inert solids which may be organic or inorganic in nature. Materials useful for this purpose include, for example, botanical flours, silicas, silicates, carbonates and clays. One convenient method of preparing a dust is to dilute a wettable powder with a finely divided carrier. Dust concentrates containing about 20% to 80% of the active ingredient are commonly made and are subsequently diluted to about 1% to 10% use concentration.

Granular formulations can be prepared by impregnating a solid such as granular fuller's earth, vermiculite, ground corn cobs, seed hulls, including bran or other grain-hulls, or similar material. A solution of one or more of the diphenyl ethers in a volatile organic solvent can be sprayed or mixed with the granular solid and the solvent then removed by evaporation. The granular material can have any suitable size, with a preferable size range of 16 to 60 mesh. The diphenyl ether will usually comprise about 2 to 15% of the granular formulation.

The diphenyl ethers of the invention can also be mixed with fertilizers or fertilizing materials before their application. In one type of solid fertilizing composition in which the diphenyl ethers can be used, particles of a fertilizer or fertilizing ingredients, such as ammonium sulfate, ammonium nitrate, or ammonium phosphate, can be coated with one or more of the ethers. The solid diphenyl ethers and solid fertilizing material can also be admixed in mixing or blending equipment, or they can be incorporated with fertilizers in granular formulations. Any relative proportion of diphenyl ether and fertilizer can be used which is suitable for the crops and weeds to be treated. The diphenyl ether will commonly be from about 5% to about 25% of the fertilizing composition. These compositions provide fertilizing materials which promote the rapid growth of desired plants, and at the same time control the growth of undesired plants.

The diphenyl ethers of the invention can be applied as herbicidal sprays by methods commonly employed, such as conventional high-gallonage hydraulic sprays, low gallonage sprays, airblast spray, aerial sprays and dusts. For low volume applications a solution of the



compound is usually used. The dilution and rate of application will usually depend upon such factors as the type of equipment employed, the method of application, the area to be treated and the type and stage of development of the weeds.

For some applications, it may be desirable to add one or more other herbicides along with diphenyl ethers of the invention. Examples of other herbicides which can be incorporated to provide additional advantages and effectiveness include:

#### CARBOXYLIC ACIDS AND DERIVATIVES

2,3,6-trichlorobenzoic acid and its salts  
2,3,5,6-tetrachlorobenzoic acid and its salts  
2-methoxy-3,5,6-trichlorobenzoic acid and its salts  
2-methoxy-3,6-dichlorobenzoic acid and its salts  
2-methyl-3,6-dichlorobenzoic acid and its salts  
2,3-dichloro-6-methylbenzoic acid and its salts  
2,4-dichlorophenoxyacetic acid and its salts and esters  
2,4,5-trichlorophenoxyacetic acid and its salts and esters  
2-methyl-4-chlorophenoxyacetic acid and its salts and esters  
2-(2,4,5-trichlorophenoxy)propionic acid and its salts and esters  
4-(2,4-dichlorophenoxy)butyric acid and its salts and esters  
4-(2-methyl-4-chlorophenoxy)butyric acid and its salts and esters  
2,3,6-trichlorophenylacetic acid and its salts  
3,6-endoxohexahydrophthalic acid  
dimethyl 2,3,5,6-tetrachloroterephthalate  
trichloroacetic acid and its salts  
2,2-dichloropropionic acid and its salts  
2,3-dichloroisobutyric acid and its salts

#### CARBAMIC ACID DERIVATIVES

ethyl N,N-di(n-propyl)thiolcarbamate  
propyl N,N-di(n-propyl)thiolcarbamate  
ethyl N-ethyl-N-(n-butyl)thiolcarbamate  
propyl N-ethyl-N-(n-butyl)thiolcarbamate  
2-chloroallyl N,N-diethyldithiocarbamate  
N-methyldithiocarbamic acid salts  
ethyl 1-hexamethyleneiminecarbothiolate  
isopropyl N-phenylcarbamate  
isopropyl N-(m-chlorophenyl)carbamate  
4-chloro-2-butyryl N-(m-chlorophenyl)carbamate  
methyl N-(3,4-dichlorophenyl)carbamate

#### Phenols

dinitro-o-(sec-butyl)phenol and its salts  
pentachlorophenol and its salts

#### SUBSTITUTED UREAS

3-(3,4-dichlorophenyl)-1,1-dimethylurea  
3-phenyl-1,1-dimethylurea  
3-(3,4-dichlorophenyl)-3-methoxy-1,1-dimethylurea  
3-(4-chlorophenyl)-3-methoxy-1,1-dimethylurea  
3-(3,4-dichlorophenyl)-1-n-butyl-1-methylurea  
3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea  
3-(4-chlorophenyl)-1-methoxy-1-phenylurea  
3-(3,4-dichlorophenyl)-1,1,3-trimethylurea  
3-(3,4-dichlorophenyl)-1,1-diethylurea  
dichloral urea

#### SUBSTITUTED TRIAZINES

2-chloro-4,6-bis(ethylamino)-s-triazine  
2-chloro-4-ethylamino-6-isopropylamino-s-triazine  
2-chloro-4,6-bis(methoxypropylamino)-s-triazine

2-methoxy-4,6-bis(isopropylamino)-s-triazine  
2-chloro-4-ethylamino-6-(3-methoxypropylamino)-s-triazine  
2-methylmercapto-4,6-bis(isopropylamino)-s-triazine  
5 2-methylmercapto-4,6-bis(ethylamino)-s-triazine  
2-methylmercapto-4-ethylamino-6-isopropylamino-s-triazine  
2-chloro-4,6-bis(isopropylamino)-s-triazine  
2-methoxy-4,6-bis(ethylamino)-s-triazine  
10 2-methoxy-4-ethylamino-6-isopropylamino-s-triazine  
2-methylmercapto-4-(2-methoxyethylamino)-6-isopropylamino-s-triazine

#### DIPHENYL ETHER DERIVATIVES

15 2,4-dichloro-4'-nitrodiphenyl ether  
2,4,6-trichloro-4'-nitrodiphenyl ether  
2,4-dichloro-6-fluoro-4'-nitrodiphenyl ether  
3-methyl-4'-nitrodiphenyl ether  
3,5-dimethyl-4'-nitrodiphenyl ether  
20 2,4'-dinitro-4-trifluoromethyldiphenyl ether  
2,4-dichloro-3'-methoxy-4'-nitrodiphenyl ether

#### ANILIDES

N-(3,4-dichlorophenyl)propionamide  
25 N-(3,4-dichlorophenyl)methacrylamide  
N-(3-chloro-4-methylphenyl)-2-methylpentanamide  
N-(3,4-dichlorophenyl)trimethylacetamide  
N-(3,4-dichlorophenyl)- $\alpha,\alpha$ -dimethylvaleramide  
30 N-isopropyl-N-phenylchloroacetamide  
N-n-butoxymethyl-N-(2,6-diethylphenyl)chloroacetamide  
N-n-methoxymethyl-N-(2,6-diethylphenyl)chloroacetamide

#### URACILS

35 5-bromo-3-s-butyl-6-methyluracil  
5-bromo-3-cyclohexyl-1,6-dimethyluracil  
3-cyclohexyl-5,6-trimethylenuracil  
40 5-bromo-3-isopropyl-6-methyluracil  
3-tert-butyl-5-chloro-6-methyluracil

#### NITRILES

2,6-dichlorobenzonitrile  
45 diphenylacetoneitrile  
3,5-dibromo-4-hydroxybenzonitrile  
3,5-diiodo-4-hydroxybenzonitrile

#### OTHER ORGANIC HERBICIDES

50 2-chloro-N,N-diallylacetamide  
N-(1,1-dimethyl-2-propynyl)-3,5-dichlorobenzamide  
maleic hydrazide  
3-amino-1,2,4-triazole  
monosodium methanearsonate  
55 disodium methanearsonate  
N,N-dimethyl- $\alpha\alpha$ -diphenylacetamide  
N,N-di(n-propyl)-2,6-dinitro-4-trifluoromethylaniline  
N,N-di(n-propyl)-2,6-dinitro-4-methylaniline  
N,N-di(n-propyl)-2,6-dinitro-4-methylsulfonylaniline  
60 O-(2,4-dichlorophenyl)-O-methyl-isopropylphosphoramidothioate  
4-amino-3,5,6-trichloropiclinic acid  
2,3-dichloro-1,4-naphthoquinone  
di(methoxythiocarbonyl)disulfide  
65 3-isopropyl-1H-2,1,3-benzothiadiazin-(4)3H-one-2,2-dioxide  
6,7-dihydrodipyridol[1,2-a:2',1'-c]pyrazidinium salts  
1,1'-dimethyl-4,4'-bipyridinium salts



3,4,5,6-tetrahydro-3,5-dimethyl-2-thio-2H-1,3,5-thiadiazine.

When mixtures of herbicides are employed, the relative proportions which are used will depend upon the crop to be treated and the degree of selectivity in weed control which is desired.


The diphenyl ethers of the invention or their precursors can be prepared by reacting a suitably substituted phenol, or the potassium or sodium salt of the phenol, with a suitably substituted halobenzene, such as a chloro-

ro- or fluorobenzene, in the presence of an alkaline agent.

The following examples will further illustrate this invention but are not intended to limit it in any way. In Table I, typical diphenyl ethers of the invention are listed, with their melting points and elemental analyses. Specific, illustrative preparations of the compounds of Examples 3, 7, 10, 19, 21, 25, 16, 37, 47, 48, 49, 51, 55, and 56 are described after Table I.

TABLE I


Diphenyl Ethers - Physical Data



Example No.	X	Y	Z	m.p. (°C.)		Analysis				
						% C	% H	% N	% Cl	% F
1	H	H	OC <sub>2</sub> H <sub>5</sub>	72-75	found	54.67	3.75	4.19		15.25
					reqs.	55.05	3.70	4.28		17.42
2	Cl	H	OCH <sub>3</sub>	95-100	found	48.78	2.89	4.01	9.65	14.91
					reqs.	48.33	2.61	4.03	10.20	16.40
3	Cl	H	OC <sub>2</sub> H <sub>5</sub>	83-84	found	49.85	3.33	3.68	9.90	15.51
					reqs.	49.80	3.07	3.87	9.80	15.75
4	Cl	H	OC <sub>2</sub> H <sub>7</sub> -i	49.5-51	found	51.02	3.58	3.52	9.62	15.58
					reqs.	51.18	3.49	3.73	9.42	15.16
5	Cl	H	OC <sub>3</sub> H <sub>7</sub> -n	75-76	found	51.36	3.60	3.62	9.43	15.00
					reqs.	51.18	3.49	3.73	9.42	15.16
6	Cl	H	OC <sub>4</sub> H <sub>9</sub> -n	51-52	found	52.07	4.03	3.38	9.28	14.65
					reqs.	52.42	3.88	3.59	9.10	14.63
7	CN	H	OC <sub>2</sub> H <sub>5</sub>	143-145	found	54.84	3.36	8.01		15.85
					reqs.	52.18	3.01	7.61		15.48
8	CN	H	OC <sub>3</sub> H <sub>7</sub> -n	96.5-98	found	55.70	3.65	7.56		15.56
					reqs.	55.74	3.58	7.65		15.56
9	CN	H	CH <sub>3</sub>	86-88.5	found	55.95	2.80	8.62		17.68
					reqs.	55.90	2.81	8.72		17.69
10	Cl	H	OH	68-70	found	47.07	2.11	4.00	10.76	17.00
					reqs.	46.79	2.12	4.20	10.63	17.08
11	Cl	H	OCH <sub>2</sub> CF <sub>3</sub>	78-80	found	43.30	1.77	3.22	8.60	27.60
					reqs.	43.35	1.94	3.37	8.54	27.40
12	Cl	H	OCH <sub>2</sub> CH=CH <sub>2</sub>	76.78.5	found	51.76	2.77	3.75	9.51	15.32
					reqs.	5.42	2.97	3.75	9.48	15.25
13	Cl	H	OCH <sub>2</sub> C≡CH	89-93	found	51.83	2.22	3.52	9.61	15.31
					reqs.	51.70	2.44	3.77	9.54	15.33
14	Cl	H	OCH <sub>2</sub> CH <sub>2</sub> C≡CCH <sub>3</sub>	93-94	found	53.99	2.95	3.35	8.93	14.12
					reqs.	54.10	3.28	3.50	8.87	14.25
15	Cl	H	OCH <sub>2</sub> CH <sub>2</sub> OH	76-77	found	47.53	2.92	3.55	9.88	14.98
					reqs.	47.70	2.94	3.71	9.38	15.09
16	Cl	H	OCH <sub>2</sub> CH <sub>2</sub> OC <sub>2</sub> H <sub>5</sub>	62-64	found	50.75	3.82	3.37	8.48	14.00
					reqs.	50.32	3.73	3.45	8.74	14.05
17	Cl	H	OCH <sub>2</sub> CH <sub>2</sub> N(CH <sub>3</sub> ) <sub>2</sub>	oil	found	50.30	3.76	6.42	9.19	13.80
					reqs.	50.44	3.98	6.92	8.76	14.03
18	Cl	H	OCOCH <sub>3</sub>	85-89	found	47.82	2.50	3.65	9.58	14.79
					reqs.	47.90	2.42	3.73	9.44	15.20
19	Cl	H	NHC <sub>2</sub> H <sub>5</sub>	82-83	found	50.01	3.23	7.82	9.95	15.48
					reqs.	49.94	3.35	7.77	9.83	15.80
20	Cl	H	N(CH <sub>3</sub> ) <sub>2</sub>	82-83	found	50.27	3.46	7.95	9.62	15.50
					reqs.	49.97	3.35	7.77	9.83	15.80
21	Cl	H	N(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	*170° C./0.01 mm	found	52.80	3.89	7.10	9.25	14.58
					reqs.	52.50	4.15	7.20	9.12	14.68
22	Cl	H	NHCH <sub>2</sub> CH <sub>2</sub> OH	85-87	found	46.22	3.33	7.06	8.50	15.30
					reqs.	47.82	3.21	7.44	9.42	15.13
23	Cl	H	OCH <sub>2</sub> CO <sub>2</sub> H	94-96	found	45.96	2.21	3.71	9.17	13.99
					reqs.	46.00	2.32	3.57	9.05	14.56
24	Cl	H	OCH <sub>2</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	76-77	found	48.49	2.93	3.25	8.58	13.65
					reqs.	48.60	3.13	3.34	8.46	13.62
25	Cl	H	OCH(CH <sub>3</sub> )CO <sub>2</sub> H	103-109	found	47.34	2.63	3.52	8.78	13.70

TABLE I-continued

Diphenyl Ethers - Physical Data




Example No.	X	Y	Z	m.p. (°C.)		Analysis				
						% C	% H	% N	% Cl	% F
26	Cl	H	OCH(CH <sub>3</sub> )CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	71-72.5	reqs.	47.35	2.73	3.45	8.73	14.05
					found	49.78	3.40	3.07	8.31	12.90
27	Cl	H	NH <sub>2</sub>	85.5-89.5	reqs.	49.85	3.49	3.23	8.18	13.15
					found	48.6	2.87	8.01	10.37	17.09
28	Cl	H		*180-185/0.04 mm	reqs.	46.93	2.42	3.42	10.66	17.14
					found	50.61	3.39	6.80	8.95	13.71
29	Cl	H	OCH <sub>2</sub> CH <sub>2</sub> C≡CH	104-105.5	reqs.	50.70	3.50	6.96	8.80	14.15
					found	52.67	3.05	3.44	9.41	14.33
30	Cl	H	OCH(CH <sub>3</sub> )C≡CH	40-42	reqs.	53.00	2.87	3.63	9.20	14.80
					found	52.59	2.77	3.61	9.66	15.27
31	Cl	H	OCH <sub>2</sub> CH <sub>2</sub> COCH <sub>3</sub>	74-76	reqs.	53.00	2.87	3.63	9.20	14.80
					found	50.44	3.27	3.36	8.81	14.49
32	Cl	H	OCH(CH <sub>3</sub> )COCH <sub>3</sub>	oil	reqs.	50.60	3.25	3.47	8.75	14.12
					found	48.07	3.26	3.17	8.18	14.26
33	Cl	H	OCONHCH <sub>3</sub>	85-88	reqs.	50.60	3.25	3.47	8.75	14.12
					found	45.57	3.21	8.02	9.48	14.76
34	Cl	H	NHCOC <sub>2</sub> H <sub>5</sub>	oil	reqs.	46.10	2.58	7.18	9.08	14.60
					found	49.50	3.14	7.01	9.12	14.71
35	Cl	H	NHCOCH <sub>2</sub> Cl	oil	reqs.	49.40	3.12	7.20	9.12	14.68
					found	44.51	1.85	6.60	17.34	14.90
36	Cl	H	CH <sub>3</sub>	*135° C./0.8 mm	reqs.	44.03	2.22	6.85	17.33	13.93
					found	50.91	2.81	4.31	10.63	16.95
37	Cl	H	Cl	*153° C./0.24 mm	reqs.	50.70	2.73	4.22	10.69	17.19
					found	44.15	1.58	4.09	18.13	18.48
38	Cl	H		49-53	reqs.	44.34	1.72	3.98	20.14	16.19
					found	47.88	2.50	3.32	10.58	14.82
39	Cl	H	OCH <sub>2</sub> CHOHCH <sub>2</sub> OH	59-64	reqs.	49.30	2.84	3.60	9.11	14.73
					found	47.49	3.32	3.35	8.88	14.73
40	Cl	H	OCH <sub>2</sub> CH <sub>2</sub> SCH <sub>3</sub>	42-45	reqs.	49.00	3.35	3.58	9.05	14.58
					found	47.09	2.95	3.34	8.80	14.05
41	Cl	H	N(C <sub>3</sub> H <sub>7</sub> -n) <sub>2</sub>	oil	reqs.	47.12	3.21	3.44	8.69	13.98
					found	54.04	4.62	6.34	8.71	13.55
42	Cl	H	OCH <sub>2</sub> CH <sub>2</sub> SO <sub>2</sub> CH <sub>3</sub>	127.5-129.5	reqs.	54.75	4.84	6.72	8.50	13.67
					found	43.53	2.71	2.99	8.27	13.19
43	Cl	H	N(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>2</sub> OH	oil	reqs.	43.69	2.98	3.19	8.06	12.96
					found	48.95	3.65	7.18	9.29	14.87
44	Cl	H	NHCONHCH <sub>3</sub>	204-208	reqs.	49.18	3.61	7.17	9.07	14.59
					found	46.52	2.56	10.76	9.40	14.47
45	Cl	H	NHCOSC <sub>2</sub> H <sub>5</sub>	111-112	reqs.	46.22	2.85	10.78	9.10	14.63
					found	45.76	2.88	6.77	8.58	13.24
46	Cl	H	N(COSC <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>	99-100	reqs.	45.66	2.88	6.66	8.42	13.55
					found	45.08	3.14	5.53	7.20	11.03
47	Cl	H	CN	95-103	reqs.	44.84	3.17	5.50	6.97	11.20
					found	49.92	1.82	7.68	11.57	15.69
48	Cl	H	CO <sub>2</sub> H	140-150	reqs.	49.05	1.77	8.18	10.35	16.65
					found	46.26	1.86	3.45	11.03	14.48
49	Cl	H	CO <sub>2</sub> CH <sub>3</sub>	oil	reqs.	46.50	1.95	3.87	9.82	15.78
					found	47.77	2.64	3.48	10.49	12.93
50	Cl	H	OCH(CH <sub>3</sub> )CO <sub>2</sub> CH <sub>3</sub>	68-70	reqs.	47.90	2.42	3.73	9.45	15.20
					found	48.58	2.98	3.15	8.58	13.73
51	Cl	H	OCH(CH <sub>3</sub> )CONH <sub>2</sub>	108-111	reqs.	48.60	2.98	3.35	8.43	13.58
					found	47.61	3.21	6.70	8.98	14.37
					reqs.	47.50	2.99	6.93	8.78	14.10



TABLE I-continued

Diphenyl Ethers - Physical Data



Example No.	X	Y	Z	m.p. (°C.)		Analysis				
						% C	% H	% N	% Cl	% F
52	Cl	H	OCH(CH <sub>3</sub> )CONHCH <sub>3</sub>	121-126	found	48.68	3.53	6.54	8.78	13.72
					reqs.	48.80	3.37	6.70	8.48	13.62
53	Cl	H	OCH(CH <sub>3</sub> )CON(CH <sub>3</sub> ) <sub>2</sub>	83-85	found	50.17	3.78	6.25	8.37	13.22
					reqs.	50.00	3.73	6.48	8.20	13.18
54	Cl	H	N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> CH <sub>2</sub> OH		found	50.44	3.99	6.92	8.76	14.08
					reqs.	50.27	4.10	6.89	8.86	14.18
55	Cl	H	N(C <sub>2</sub> H <sub>5</sub> )CONHCH <sub>3</sub>	127.5-128.5	found	49.51	3.91	9.95	8.64	12.60
					reqs.	48.87	3.62	10.06	8.49	13.64
56	Cl	Cl	OC <sub>2</sub> H <sub>5</sub>	100.5-102	found	45.26	2.43	3.36	18.00	12.33
					reqs.	45.47	2.54	3.54	17.90	14.39

\*boiling point

## EXAMPLE 3

Preparation of  
2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether

## Method A

a.

1,3-Bis(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)benzene

A solution of potassium hydroxide (3.26 g. 0.05 mole, 85%) in water (~3 g.) is added slowly dropwise to a solution of resorcinol (2.75 g. 0.025 mole) and 3,4-dichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene (10.75 g. 0.05 mole) in sulfolane (125 ml) at 150°-160° C., with stirring. When the addition is complete, the strongly colored reaction mixture is stirred at 150°-160° C. overnight, then cooled, diluted with benzene (200 ml), and washed cautiously with water (700 ml). Hexane (200 ml) is added and the mixture washed with water (600 ml), dilute sulfuric acid (600 ml), dilute sodium hydroxide solution (600 ml), and water (600 ml), dried, and the solvent removed to give 1,3-bis(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)benzene (8.6 g. 65%) b.p. 160°-70° C./0.1 mm.

b.

1,3-Bis(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-4-nitrobenzene

1,3-Bis(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-benzene (12 g. 0.0255 mole) is added to a mixture of concentrated nitric acid (12 g.) and sulfuric acid (15 g.) at 5° C. The temperature is then allowed to rise to 25°-30° C. with manual stirring and mild ice bath cooling and after 10-20 minutes, the oil solidifies. The mixture is taken up in water/benzene (400 ml)/hexane (400 ml) and the organic phase is washed with water, dried, filtered through activated silica gel (~20 g.), and the solvents removed. The residue is recrystallized from isopropanol

to give 1,3-bis(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-4-nitrobenzene, (7.4 g. 56%) m.p. 110°-111.5° C.

c.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether

A 10% solution of potassium hydroxide in ethanol (10 ml) is added to a solution of 1,3-bis(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-4-nitrobenzene (2 g. 0.0039 mole) in dioxane (20 ml). After forty minutes at room temperature, the solution is heated to 45° C. for eight minutes, then cooled, diluted with benzene (50 ml) and hexane (50 ml) and washed with water (3×100 ml), dried, and the solvents removed. The residue is recrystallized from isopropanol to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenylether (1.21 g. 86%) m.p. 83°-84° C.

## Method B

a.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-hydroxyphenyl ether

A mixture of the di-potassium salt of resorcinol (186.3 g., 1 mol), 3,4-dichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene (53.7 g., 0.25 mol), and sulfolane (100 ml.) is stirred for 30 minutes. Benzene (500 ml) and water (200 ml) are added and the organic phase is washed with water (3×200 ml), diluted with hexane (500 ml) and washed again with water, dried, filtered through activated silica gel (15g.), the solvents removed, and the residue distilled to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-hydroxyphenyl ether (45.1 g., 62%) b.p. 112°-124° C./0.3 mm.

b.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetoxyphenyl ether

A mixture of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-hydroxyphenylether (184 g.) and acetic anhydride (334



## 15

g.) is heated on a steam bath for 1 hour and cooled. The mixture is washed with 5% sodium carbonate solution (2×500 ml.) and distilled to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetoxyphenyl ether (84 g., 40%) b.p. 107°–117° C./0.09 mm.

c.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetoxy-4-nitrophenyl ether

A solution of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetoxyphenyl ether (249 g., 0.75 mol) in 1,2-dichloroethane (1200 ml.) is stirred 2.2 hours at 20°–30° C. with a cooled mixture of concentrated sulfuric acid (276 g.) and nitric acid, 70% (227 g.). Hexane (700 ml.) is added and the oil layer washed once with water, 3 times with dilute sodium bicarbonate, and once more with water, dried, filtered through activated silica gel (~40 g.), the solvents removed. The product is crystallized from hexane-benzene to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetoxy-4-nitrophenyl ether (207.5 g., 73%) m.p. 83°–89° C.

d.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-hydroxy-4-nitrophenyl ether

A solution of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetoxy-4-nitrophenyl ether (204.9 g., 0.545 mol) in methanol (2900 ml.) is stirred 1 hour at 20° C. with potassium carbonate (103 g., 0.745 mol). Ninety percent of the methanol is removed; and benzene (1 liter), 7–8% sulfuric acid solution (1600 ml.) are added and stirred 1.5 hours at 25° C. The oil layer is washed twice more with water (200 ml. each), dried, filtered through activated silica gel (40 g.), the solvents removed, and the residue crystallized in hexane to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-hydroxy-4-nitrophenyl ether (165.3 g. 90%) m.p. 68.5°–73° C.

e.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether

A solution of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-hydroxy-4-nitrophenyl ether (60 g., 0.018 mol, 73% pure) and dimethylformamide (100 g.) is converted to the potassium phenoxide and stirred with ethyl bromide (35 g., 0.32 mol) 3 hours at 60° C. and 5 hours at 80° C. Perchloroethylene (150 g.) is added and the solution washed twice with water (~250 ml. each) at 50° C. The solvents are removed to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether (56 g., 83%, 71% pure).

## EXAMPLE 7

## Preparation of

2-Cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether

A solution of potassium hydroxide (2.6 g., 0.04 mole) 87.3% pure of 3-ethoxy-4-nitrophenol (7.3 g., 0.04 mole) in methanol (30 ml) is stripped to dryness under reduce pressure. A residue of potassium 3-ethoxy-4-nitrophenoxide is dissolved in sulfolane (200 g.) and 4-chloro-3-cyano-benzotrifluoride (8.2 g., 0.04 mole) is added. Gas-liquid chromatography shows the reaction to be complete after stirring at 110° C. for 4½ hours and 135° C. for 2½ hours. The reaction mixture is cooled and poured into deionized water and the precipitate which forms is filtered off and air dried. Recrystallization from

## 16

isopropanol yields 2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether (7.4 g. 53%) m.p. 143°–145° C.

## EXAMPLE 19

## Preparation of

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethylamino-4-nitrophenyl ether

A solution of 1,3-bis(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-4-nitrobenzene (12.8 g. 0.025 mol), and ethylamine (6.7 g. 0.15 mol) in dioxane (120 ml.) is heated in a pressure bottle 4.5 hours at 50°–55° C. and 4.3 hours at 90°–95° C. Benzene (200 ml.), hexane (70 ml.) and water (500 ml.) are added and the organic phase is washed with water (500 ml.), 10% sodium bicarbonate solution (200 ml.) and water (200 ml), dried, filtered through activated silica gel (25 g.), the solvents removed, and the residue is crystallized from hexane to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethylamino-4-nitrophenyl ether (7.9 g. 88%) m.p. 82°–83° C.

## EXAMPLE 21

## Preparation of

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-diethylamino-4-nitrophenyl ether

A solution of 1,3-bis(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-4-nitrobenzene (12.8 g. 0.025 mol) in p-dioxane (130 ml.) is heated under reflux for 26 hours at 65°–95° C. with diethylamine (50 g. 0.66 mol). Benzene (~200 ml.) and water (~500 ml.) are added followed by hexane (~70 ml.) and the oil layer is separated, washed with water (500 ml), 10% sodium bicarbonate solution (200 ml), and water (200 ml.), dried, filtered through activated silica gel (~25 g.), the solvents removed. The residual oil is distilled in vacuo to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-diethylamine-4-nitrophenyl ether (8.15 g 84%) b.p. 180°–190° C./0.01 mm.

## EXAMPLE 25

## Preparation of

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carboxyethoxy)-4-nitrophenyl ether

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carbethoxyethoxy)-4-nitrophenyl ether (8.6 g. of 0.02 mol), potassium hydroxide 86% (2.6 g. 0.04 mol), ethanol (8 ml.), dioxane (8 ml.) and water (100 ml.) are heated at 90°–95° C. for 30 minutes. Ether (200 ml.) and water (200 ml.) are added and the mixture acidified with dilute sulfuric acid, the water layer extracted 3 times with ether (200 ml. each), dried, and the ether removed to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carboxyethoxy)-4-nitrophenyl ether (6.4 g. 79%) m.p. 108°–109° C.

## EXAMPLE 26

## Preparation of

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carbethoxyethoxy)-4-nitrophenyl ether

Potassium 2-nitro-5-(2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)phenoxide (7.4 g. 0.02 mol), ethyl 2-bromopropionate (3.6 g. 0.02 mol) and sulfolane (50 ml.) are heated 1.5 hours at 90°–95° C. Benzene (100 ml.) and hexane (100 ml.) are added and the solution is washed with dilute sodium carbonate solution then with water, dried, and the solvents removed. The residue is crystal-



lized from pentane to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carbethoxyethoxy)-4-nitrophenyl ether (6.2 g., 71%) m.p. 71°–74° C.

## EXAMPLE 37

Preparation of  
2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-chloro-4-nitrophenyl ether

a.

## 3-Chloro-4-nitrofluorobenzene

m-Chlorofluorobenzene (240 g. 1.85 moles) is added to a mixture of sulfuric acid (185 g. 1.85 moles) and nitric acid (166 g., 1.85 moles) at –5° C. in 3.5 hours, stirred 13 hours, then benzene (200 ml.) and hexane (200 ml.) are added. The extract is washed with water (1×300 ml.), sodium carbonate solution (1×300 ml.), and water (1×300 ml.), dried and the solvents removed. The residue is distilled to give 138 g. of mixed isomers. The 4-nitro isomer crystallizes and is filtered off to give 3-chloro-4-nitrofluorobenzene (51 g. 16.7%) m.p. 36°–38° C.

## EXAMPLES 47–49

Preparation of  
2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carbomethoxy-4-nitrophenyl ether

a.

## 3-Cyano-4-nitrofluorobenzene

m-Fluorobenzonitrile (96.8 g., 0.8 mole) is added in 2½ hours to a mixture of concentrated sulfuric acid (600 ml.) and potassium nitrate (80.9 g., 0.8 mole) at 3°–6° C., then allowed to warm to 25° C. The mixture is poured over cracked ice (3000 ml.), extracted with chloroform (5×250 ml.), dried and the solvent removed. The residue is extracted with pentane and dried to give 3-cyano-4-nitrofluorobenzene (115 g., 86.5%) m.p. 102°–104° C.

b.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-cyano-4-nitrophenyl ether

The potassium phenoxide of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-cresol (13.5 g 0.0688 mole) prepared in sulfolane at 5° C. is added to a solution of 3-cyano-4-nitrofluorobenzene (11.4 g., 0.0688 mole) in sulfolane at 120° C. in 4 hours, stirred 18 hours and cooled. Benzene (200 ml.) and hexane (100 ml) are added and the solution is water washed (5×250 ml.), dried, filtered and the solvents removed. The residue is crystallized to give 2-chloro- $\alpha,\alpha,\alpha$ -p-tolyl-3-cyano-4-nitrophenyl ether (16.3 g., 69%) m.p. 95°–103° C. 85% pure.

c.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carboxy-4-nitrophenyl ether

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-cyano-4-nitrophenyl ether (11.2 g. 0.0327 mole), acetic acid (25 ml.), and hydrobromic acid (12 ml. of 47.8% purity) are heated at 120° C., in a pressure bottle, for 2 days, poured over cracked ice and extracted with benzene (2×150 ml). The benzene solution is dried, filtered, the solvent removed, and the residue crystallized from pentane to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carboxy-4-nitrophenyl ester (7.5 g. 63.5%) m.p. 140°–150° C., 85% pure.

d.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carbomethoxy-4-nitrophenylether

Hydrogen chloride is bubbled thru a solution of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carboxy-4-nitrophenyl ether (2.3 g. 0.0064 mole) in methanol (50 ml.) for 10 hours at 32° C., stirred overnight and the solvent removed to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carbomethoxy-4-nitrophenyl ether (1.5 g. 40%)

## EXAMPLE 51

Preparation of  
2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carbamoylethoxy)-4-nitrophenyl ether

a.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-chloroformylethoxy)-4-nitrophenyl ether

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carboxyethoxy)-4-nitrophenyl ether (34.8 g. 0.086 mol), thionyl chloride (20.4 g. 0.172 mol), and benzene (150 ml.) are stirred 5 hours at 95° C. and 16 hours at 25° C. The benzene is removed to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-chloroformylethoxy)-4-nitrophenyl ether (33.4 g. 92%).

b.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carbamoylethoxy)-4-nitrophenyl ether

A solution of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-chloroformylethoxy)-4-nitrophenyl ether (4.2 g. 0.01 mol) in ether (50 ml.) is added to an ether solution (200 ml.) saturated with gaseous ammonia at zero temperatures. After 30 minutes, water (100 ml.) is added and the ether layer separated. The aqueous phase is extracted with ether (3×200 ml.) and the combined extracts dried, filtered through activated silica gel (~20 g.), and the solvent removed to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carbamoylethoxy)-4-nitrophenyl ether (2.4 g. 60%) m.p. 108°–111° C.

## EXAMPLE 55

Preparation of  
2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(3-methyl-1-ethylureido)-4-nitrophenyl ether

a.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(N-ethylchloroformamido)-4-nitrophenyl ether

A mixture of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethylamino-4-nitrophenyl ether (3.6 g. 0.01 mol), phosgene (18.9 g. 0.19 mol), 2,6-lutidine (2.2 g. 0.02 mol), and benzene (~130 ml.) are heated in a pressure bottle 64 hours at 90°–95° C. The mixture is cooled, filtered and the solvent removed to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(N-ethylchloroformamido)-4-nitrophenyl ether.

b.

2-Chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(3-methyl-1-ethylureido)-4-nitrophenyl ether

A solution of 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(N-ethylchloroformamide)-4-nitrophenyl ether (4.4 g., 0.01 mol), methylamine (3.3 g 0.11 mol) and benzene (~60 ml.) is allowed to stand 25 minutes at zero °C., filtered,



and the solvent removed. Benzene (~100 ml.) and hexane (50 ml.) are added and the solution is washed with water (100 ml.) and aqueous 10% sodium carbonate solution (2×100 ml.), dried, and the product absorbed on activated silica gel (~25 g.). The product is eluted with a mixture of benzene (400 ml.) and methanol (40 ml) the solvents removed and the product recrystallized to give 2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(3-methyl-1-ethylureido)-4-nitrophenyl ether (2.6 g. 62%) m.p. 127.5°–128.5° C.

## EXAMPLE 56

Preparation of  
2,6-Dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether

a.

3,4-Dichloro-5-nitro- $\alpha,\alpha,\alpha$ -trifluorotoluene

3,4-Dichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene (862 g. 4.0 mols) is added to a stirred mixture of concentrated sulfuric acid (4400 g.) and nitric acid (3400 g.) at 35° C. The mixture is stirred 70 minutes at 95° C. and allowed to separate. The oil layer is washed once with water and twice distilled to give 3,4-dichloro-5-nitro- $\alpha,\alpha,\alpha$ -trifluorotoluene (188 g. 18%) b.p. 115°–118° C./15 mm, 88% pure.

b.

5-Amino-3,4-dichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene

500 ml. of an ethanolic solution containing 3,4-dichloro-5-nitro- $\alpha,\alpha,\alpha$ -trifluorotoluene (188 g. 0.72 mol), and platinum oxide (Adam's catalyst) (0.2 g.) is reduced at room temperature in a low pressure hydrogenation apparatus to give 5-Amino-3,4-dichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene (129.9 g. 78%) b.p. 65°–70° C./1–2 mm.

c.

3,4,5-Trichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene

A solution of sodium nitrite (39 g.) in water (85 ml.) is added over 1 hour to a solution of 5-amino-3,4-dichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene (117.5 g., 0.51 mol) in 1700 ml. concentrated hydrochloric acid at –6° C. and the solution stirred for 1 hour then filtered. The filtrate is added to a solution of cuprous chloride (76.5 g.) in concentrated hydrochloric acid (500 ml.) over 5 minutes at 0° to 8° C. and gradually heated to 80° C. over 80 minutes. The reaction mixture is cooled to 35° C. and extracted with hexane (2×300 ml.). The extract is washed with water, 2% sodium hydroxide solution, dried and distilled to give 3,4,5-trichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene (70 g., 55%) b.p. 82°–86° C./10 mm, 95% pure.

d.

1,3-Bis(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)benzene

A mixture of 3,4,5-trichloro- $\alpha,\alpha,\alpha$ -trifluorotoluene (10 g. 0.04 mol), and the dipotassium salt of 1,3-dihydroxybenzene (4 g. 0.021 mol) in 150 ml. sulfolane is stirred and heated 70 minutes at ~120° C. The cooled reaction mixture is diluted with benzene (350 ml.) and washed once with water (1 l). Hexane (200 ml.) is added, and the solution washed with water (3×500 ml.) dried, filtered through activated silica gel (~25 g.), and the solvents removed. The residual oil is crystallized from a mixture of pentane and benzene to give 1,3-

bis(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)benzene (5.3 g. 49%) m.p. 121°–122° C.

e.

1,3-Bis(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-4-nitrobenzene

A cooled mixture of concentrated sulfuric acid (6.5 ml.) and nitric acid (4.4 ml.) is added with stirring to an ice cold solution of 1,3-bis(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-benzene (11.1 g. 0.021 mol) in 1,2-dichloroethane (30 ml.). After 30 minutes at room temperature, the phases are allowed to separate and the organic phase washed twice with water. Benzene (200 ml.) is added and the solution washed twice with dilute sodium carbonate solution, dried, filtered through activated silica gel (~25 g.), and the solvents removed. The residual crystals are triturated with pentane, filtered, and dried to obtain 1,3-bis(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-4-nitrobenzene (9.9 g., 82%) m.p. 137.5°–140.5° C., 90% pure.

f.

2,6-Dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether

A solution of potassium hydroxide, 86% (1.9 g., 0.029 mol) in ethanol (20 ml.) is added to a solution of 1,3-bis(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-4-nitrobenzene (8.0 g., 0.014 mol) in p-dioxane (70 ml.) and warmed 1 hour at 50° C. The solution is cooled and benzene (~250 ml.) is added and crystals of potassium 2-nitro-5-(2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyloxy)-phenoxide (2.9 g. 52%) are collected. Treatment with acid yields the free phenol, 2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-hydroxy-4-nitrophenyl ether (2.0 g. 40%) m.p. 84.5°–86.5° C. This phenol (1.7 g. 0.0046 mol) is reconverted to the potassium salt, dissolved in dimethylformamide (20 ml.) and treated with ethyl iodide (1.2 g. 0.0077 mol) 2.5 hours at 50°–70° C. The reaction mixture is diluted with benzene (~100 ml.) and hexane (~50 ml.), washed with water (3×100 ml.), dried, filtered through activated silica gel (~15 g.), and the solvents removed to give 2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether (0.8 g. 44%) m.p. 100.5°–102° C.

From the filtrate of the 2.9 g. of phenoxide there is recovered ethyl 2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl ether (3.0 g. 82%) b.p. 78° C./5 mm, and an additional amount -trifluoro-p-tolyl- of the product 2,6-dichloro- $\alpha,\alpha,\alpha$ -3-ethoxy-4-nitrophenyl ether (0.35 g. 6%) m.p. 88°–93° C.

## EXAMPLES 57 to 72

Following the procedures of Examples 1 to 56, other diphenyl ethers of Formula I are prepared. Among the compounds which are prepared by these procedures are:

- $\alpha,\alpha,\alpha,\alpha',\alpha',\alpha'$ -hexafluoro-2,4-xylyl-3-ethoxy-4-nitrophenyl ether
- $\alpha^4,\alpha^4,\alpha^4$ -trifluoro-2,4-xylyl-3-n-propoxy-4-nitrophenyl ether,
- 2-chloro-6, $\alpha,\alpha,\alpha$ -tetrafluoro-p-tolyl-3-ethyl-4-nitrophenyl ether,
- 2-iodo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-methoxy-4-nitrophenyl ether,
- 2-chloro-6-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-methylthio-4-nitrophenyl ether,



2-bromo- $\alpha,\alpha,\alpha,\alpha'$ -trifluoro-4,6-xylyl-3-ethoxy-4-nitrophenyl ether,  
2-chloro- $\alpha,\alpha,\alpha,\alpha',\alpha'$ -hexafluoro-4,6-xylyl-3-methoxy-4-nitrophenyl ether,  
2-bromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethoxy-4-nitrophenyl ether,  
2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-n-butyl-4-nitrophenyl ether,  
2-chloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-bromo-4-nitrophenyl ether,  
2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-acetoxy-4-nitrophenyl ether,  
2-bromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-diethylamino-4-nitrophenyl ether,  
2,6-dichloro- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-ethylamino-4-nitrophenyl ether,  
2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-(1-carbethoxyethoxy)-4-nitrophenyl ether,  
2-cyano- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carbomethoxy-4-nitrophenyl ether, and  
2-bromo- $\alpha,\alpha,\alpha$ -trifluoro-p-tolyl-3-carboxy-4-nitrophenyl ether.

These diphenyl ethers have herbicidal properties.  
It should be noted that the diphenyl ethers of the invention can also be named correctly using different systems of nomenclature. For example, the diphenyl ether of Example 3 can also be named as 2-cyano-4-trifluoromethyl-3'-ethoxy-4'-nitrodiphenyl ether. However, within the specification and claims of this invention the system of nomenclature exemplified in Examples 1-56 has been followed.  
The following examples show the herbicidal properties of the diphenyl ethers of the invention.

EXAMPLE 73

This example shows the herbicidal activity of diphenyl ethers of the invention towards a number of common weeds. Using the procedure described below, diphenyl ethers were evaluated for control of the following weeds:  
At 10 pounds per acre:

Monocots

barnyardgrass (*Echinochloa crusgalli*)  
crabgrass (*Digitaria* spp.)  
nutsedge (*Cyperus esculentus*)  
wild oats (*Avena fatua*)

Dicots

bindweed (*Convolvulus arvensis*)  
curly dock (*Remex crispus*)  
velvetleaf (*Abutilon theophrasti*)  
wild mustard (*Brassica haber*)  
At 2 and 4 pounds per acre:

Monocots

barnyardgrass (*Echinochloa crusgalli*)  
\*\*Bermudagrass (*Cynodon dactylon*)  
crabgrass (*Digitaria* spp.)  
\*downy brome (*Bromus tectorum*)  
foxtail (*Setaria faberii*)  
Johnsongrass (*Sorghum halepense*)  
nutsedge (*Cyperus esculentus*)  
quackgrass (*Agropyron repens*)  
\*ryegrass (*Lolium perenne*)  
\*wild oats (*Avena fatua*)  
\*yellow millet (*Panicum miliaceum*)

Dicots

bindweed (*Convolvulus arvensis*)  
cocklebur (*Xanthium pensylvanicum*)  
\*\*coffeeweed (*Sesbania macrocarpa*)  
\*curly dock (*Rumex crispus*)  
\*lambsquarters (*Chenopodium album*)  
morningglory (*Ipomoea purpurea*)  
\*pigweed (*Amaranthus retroflexus*)  
10 \*\*ragweed (*Ambrosia artemisiifolia*)  
\*smartweed (*Polygonum pensylvanicum*)  
\*\*tomato (*Lycopersicon esculentum*)  
velvetleaf (*Abutilon theophrasti*)  
\*wild carrot (*Daucus carota*)  
15 \*wild mustard (*Brassica haber*)  
\*Examples 1 to 9 only  
\*\*Examples 10 to 56 only

The following test procedure is employed. Seeds of selected crops and weeds are planted in soil in flats. For preemergence tests, the flats are treated with the test compound immediately after the planting. For post-emergence tests, the seeds are allowed to germinate, and after 2 weeks the flats are treated with the test compound. The compound to be evaluated is dissolved in acetone, diluted with water, and sprayed over the flats using a carrier volume equivalent to 50 gallons per acre at the rate of application (pounds per acre, lb/A.) specified in the tables. About 2 weeks after the application of the test compound, the state of growth of the plants is observed and the phytotoxic effect of the compound is evaluated. Table II gives the average percent control achieved by the test compounds in terms of the percent of the plants which are killed by the compounds.

TABLE II

		HERBICIDAL ACTIVITY (% control)						
Compound of			Preemergence			Postemergence		
Example No.	lb./A.		10	4	2	10	4	2
40	1	M*	42	76		82	57	
		D*	35	45		100	58	
	2	M	97	97		100	99	
		D	100	80		100	100	
	3	M	99	89	**84	100	99	+ 96
		D	100	99	**66	100	100	+ 97
45	4	M		98			80	
		D		100			100	
	5	M	97	84		100	99	
		D	100	78		100	100	
	6	M	65	77		97	91	
		D	70	55		100	94	
50	7	M	61	65	65	100	75	82
		D	60	57	54	100	77	80
	8	M	81	61	52	85	48	35
		D	82	57	46	100	80	66
	9	M	99	90	81	100	97	82
		D	92	66	67	100	81	75
55	10	M		70	66		67	17
		D		100	70		98	85
	11	M		81	88		82	100
		D		98	99		99	100
	12	M		86	72		77	99
		D		96	93		100	100
60	13	M		76	78		77	100
		D		90	72		98	100
	14	M		64	30		61	91
		D		78	22		94	100
	15	M		99	68		79	76
		D		100	90		99	94
65	16	M		86	67		77	82
		D		94	75		96	94
	17	M		51	20		45	39
		D		84	31		89	98
	18	M		67	99		28	28



TABLE II-continued

Compound of Example No.	lb./A.	HERBICIDAL ACTIVITY (% control)					
		Preemergence			Postemergence		
		10	4	2	10	4	2
19	D	100	100		86	98	
	M	91	77		84	94	
	D	96	91		98	97	
20	M	88	80		78	80	
	D	99	93		88	97	
21	M	91	79		74	90	
	D	99	90		96	94	
22	M	72	63		60	70	
	D	98	77		99	100	
23	M	90	55		82	71	
	D	64	87		100	100	
24	M	91	62		86	84	
	D	68	97		99	100	
25	M	87	40		97	88	
	D	88	93		100	100	
26	M	98	75		99	93	
	D	100	99		100	100	
27	M	76	74		82	99	
	D	98	90		100	100	
28	M	74	79		73	73	
	D	95	79		99	100	
29	M	74	66		79	97	
	D	73	84		100	100	
30	M	81	78		75	91	
	D	99	94		100	100	
31	M	74	68		61	68	
	D	99	69		100	100	
32	M	86	81		86	100	
	D	100	81		100	100	
33	M	53	79		32	65	
	D	99	74		98	100	
34	M	35	83		32	47	
	D	52	85		89	90	
35	M	40	60		45	63	
	D	40	72		98	80	
36	M	93	100		87	94	
	D	100	100		100	98	
37	M	99	100		92	100	
	D	100	100		100	100	
38	M	47	62		43	51	
	D	86	68		98	58	
39	M	65	77		64	61	
	D	99	80		98	100	
40	M	86	88		64	81	
	D	92	98		100	100	
41	M	60	84		71	56	
	D	96	62		95	94	
42	M	62	83		38	63	
	D	80	60		98	88	
43	M	0	99		2	61	
	D	48	100		47	80	
44	M	66	0		60	7	
	D	96	17		98	10	
45	M	13	68		17	53	
	D	58	77		88	78	
46	M	0	34		0	0	
	D	20	87		30	4	
47	M	61	80		11	24	
	D	40	80		72	84	
48	M	99	100		75	84	
	D	97	100		100	97	
49	M	92	100		86	77	
	D	95	100		100	100	
50	M	90	98		97	100	
	D	83	100		100	100	
51	M	79	96		75	86	
	D	100	100		100	98	
52	M	79	98		87	93	
	D	67	75		100	100	
53	M	98	98		79	81	
	D	83	100		100	100	
54	M	83	98		79	71	
	D	75	98		100	100	
55	M	91			69	31	
	D	73			95	96	
56	M	91	100		99	100	

TABLE II-continued

Compound of Example No.	lb./A.	HERBICIDAL ACTIVITY (% control)					
		Preemergence			Postemergence		
		10	4	2	10	4	2
5	D	83	100		100	99	

\*\* $\frac{1}{2}$  lb./A.+ $\frac{1}{2}$  lb./A.

10 \*M = Monocots; D = Dicots

## EXAMPLE 74

15 This example shows the selective herbicidal activity of diphenyl ethers of the invention in a number of agronomic crops. Following the general test procedure of Example 73, diphenyl ethers are evaluated for significant tolerance (as shown by 50% or less kill of the test crop at levels of application which give more than 50% kill of many or all of the weeds of Example 73) towards some or all of the following common agronomic crops (not all compounds tested against all crops): alfalfa, snapbeans, corn, cotton, cucumbers, peanuts, rape, rice, 20 safflower, soybeans, tomatoes, and wheat.

25 Tolerance to snapbeans in preemergence applications is shown by the compounds of Examples 4 and 26. Tolerance to corn in preemergence applications is shown by the compounds of Examples 4, 11, 19, 21, 34, 40, 52, 30 and 54 and in postemergence applications by the compounds of Examples 4, 18, 19, 21, 26, 34, 36, 40, 49, 52, 53, 54, and 56. Tolerance to cotton in preemergence applications is shown by the compounds of Examples 3, 35 4, and 30 and in postemergence or layby applications by the compound of Example 3. Tolerance to peanuts in preemergence applications is shown by the compounds of Examples 3, 19, 21, 30, 34, 36, 40, 48, 49, 50, 52, 53, 40 and 56 and in postemergence applications by the compounds of Examples 4, 18, 34, 36, 37, 40, 48, 53, and 54. Tolerance to rice in preemergence applications is shown by the compounds of Examples 4, 19, 21, 30, 34, 40, 52, and 54 and in postemergence applications or in 45 applications on transplanted rice by the compounds of Examples 3, 11, 18, 20, 30, 34, 40, 48, 53, and 54. Tolerance to safflower in preemergence applications is shown by the compound of Example 3. Tolerance to soybeans in preemergence applications is shown by the compounds of Examples 3, 4, 11, 18, 19, 21, 26, 30, 34, 40, 48, 49, 50, 52, 53, and 54, in postemergence or layby applications by the compounds of Examples 3 and 34. Tolerance to wheat in preemergence applications is shown by the compounds of Example 4, 18, 19, 26, 34, 40, 50, 54, and 56 and in postemergence applications by the compounds of Examples 4, 18, 21, 30, 34, 36, 48, and 54. 55


60 It is to be understood that changes and variations may be made without departing from the spirit and scope of the invention as defined by the appended claims.

We claim:

65 [1. A compound of the formula



25



wherein

X is a hydrogen atom, a halogen atom, a trifluoromethyl group, or a (C<sub>1</sub>-C<sub>4</sub>)alkyl group,

Y is a hydrogen atom, a halogen atom, or a trifluoromethyl group, and

Z is a carboxy group, a carbalkoxy group having up to 4 carbon atoms in the alkoxy moiety, a carboxyalkyl group having up to 4 carbon atoms, a carbalkoxyalkyl group having up to 6 carbon atoms.】

【2. A compound according to claim 1 wherein Y is a hydrogen atom.】

【3. A compound according to claim 2 wherein X is a halogen atom.】


4. A compound according to claim 【3】 9 wherein Z is a carbalkoxy group.

5. A compound according to claim 4 wherein X is a chlorine atom and Z is a carbomethoxy group.

【6. A compound according to claim 1 wherein X is a halogen atom, Y is a halogen atom, and Z is a carbalkoxy group.】


7. A compound of the formula:

26




wherein Z is a carbalkoxy group having 1 to 4 carbon atoms in the alkoxy moiety.

8. A compound of the formula:



9. A compound of the formula



wherein X is a halogen atom and Z is a carboxy group or a carbalkoxy group having up to 4 carbon atoms in the alkoxy moiety.

\* \* \* \* \*