



US00RE37004E

**United States Patent** [19]

[11] E

**Patent Number: Re. 37,004****Takahashi et al.**[45] **Reissued Date of Patent: Dec. 26, 2000**

[54] **WATER-SOLUBLE MONOAZODYE CONTAINING A TRIAZINYL OR PYRIMIDINYL GROUP AND AT LEAST ONE VINYL SULFONYL TYPE GROUP IN ITS STRUCTURE**

[75] Inventors: **Yousuke Takahashi**, Kitakyushu; **Yukihara Shimizu**, Sagamihara; **Toshio Hihara**, Kitakyushu; **Kiyoshi Himeno**, Munakata, all of Japan

[73] Assignee: **Dystar Japan Ltd.**, Osaka, Japan

[21] Appl. No.: **08/916,366**

[22] Filed: **Aug. 22, 1997**

#### Related U.S. Patent Documents

Reissue of:

[64] Patent No.: **5,380,827**  
 Issued: **Jan. 10, 1995**  
 Appl. No.: **08/049,410**  
 Filed: **Apr. 20, 1993**

#### [30] Foreign Application Priority Data

Apr. 20, 1992 [JP] Japan ..... 4-125380  
 Apr. 28, 1992 [JP] Japan ..... 4-134426  
 Oct. 23, 1992 [JP] Japan ..... 4-307867

[51] **Int. Cl.<sup>7</sup>** ..... **C09B 62/028**; C09B 62/085;  
 C09B 62/245; C09B 62/51

[52] **U.S. Cl.** ..... **534/638**; 534/632; 534/642

[58] **Field of Search** ..... 534/632, 638

#### [56] References Cited

##### U.S. PATENT DOCUMENTS

5,380,827 1/1995 Takahashi et al. .... 534/638  
 5,391,718 2/1995 Tzikas et al. .... 534/637  
 5,527,886 6/1996 Auerbach et al. .... 534/638  
 5,530,104 6/1996 Tzikas et al. .... 534/642  
 5,731,421 3/1998 Tzikas et al. .... 534/638  
 5,747,657 5/1998 Lehr ..... 534/638 X

##### FOREIGN PATENT DOCUMENTS

0 133 843 3/1985 European Pat. Off. .  
 2 110 325 6/1972 France .  
 55-163276 12/1980 Japan .  
 56-91083 7/1981 Japan .  
 2 239 024 6/1991 United Kingdom .


*Primary Examiner*—Fiona T. Powers

*Attorney, Agent, or Firm*—Oblon, Spivak, McClelland, Maier & Neustadt, P.C.


#### [57] ABSTRACT

A water-soluble monoazodye having the following formula (I) in its free acid form:


(I)



wherein X is  $-\text{CH}=\text{CH}_2$  or  $-\text{CH}_2\text{CH}_2\text{W}$  wherein W is a leaving group removable by alkali;  $\text{R}^1$  is methyl,  $-\text{NHCONH}_2$  or  $-\text{NHCOT}^1$  wherein  $\text{T}^1$  is methyl, ethyl,  $-\text{CH}_2\text{CH}_2\text{COOH}$  or  $-\text{CH}=\text{CHCOOH}$ ;  $\text{R}^2$  is hydrogen, lower alkyl or lower alkoxy; and Z is



wherein P is cyano or halogen, each of Q and U which are independent of each other, is halogen, V is halogen or



and Y is halogen or a group of the following formula (1) or (2):



(1)



(2)

wherein  $\text{R}^3$ , A,  $\text{R}^4$  and E are as defined in the disclosure and G is as defined above, has a moderate yellow to orange color and is particularly suitable for dyeing cellulose fibers or nitrogen-containing fibers.

**3 Claims, No Drawings**

1


**WATER-SOLUBLE MONOAZODYE  
CONTAINING A TRIAZINYL OR  
PYRIMIDINYL GROUP AND AT LEAST ONE  
VINYL SULFONYL TYPE GROUP IN ITS  
STRUCTURE**

Matter enclosed in heavy brackets [ ] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.


This present invention relates to a water-soluble monoazodye of a moderate yellow color having a novel chemical structure. Particularly, it relates to a water-soluble monoazodye having from 2 to 4 reactive groups in the chemical structure and being particularly suitable for dyeing cellulose fibers or nitrogen-containing fibers.

It is common to use water-soluble reactive dyes to dye fibers containing cellulose fibers or nitrogen-containing fibers. Such water-soluble reactive dyes are required to be excellent in level dyeing and capable of dyeing the objects in deep color, and yet the dyed products are required to be excellent in various fastnesses. These properties of reactive dyes delicately differ depending upon the basic structures, substituents or reactive groups of the dyes or depending upon the combination thereof. Therefore, dyes having various chemical structures have been proposed.

Japanese Examined Patent Publications No. 18474/1983 and No. 18471/1983 disclose reactive dyes having the following structural formulas (A) and (B), as examples of dyes of moderate yellow type:



(A)



(B)

In each of these reactive dyes, the diazo component is a component of a naphthalene type, but these reactive dyes are still slightly inadequate in the level dyeing property, and they are not fully satisfactory also in the build-up property.

The present invention has been made in view of the above problems, and it is an object of the present invention to provide a water-soluble monoazodye useful as a reactive dye of moderate yellow to orange type which is excellent not only in the chlorine fastness and light fastness but also in the level dyeing property and which is also excellent in the build-up property.


As a result of extensive studies, the present inventors have found it possible to accomplish the above object by using a novel monoazo compound having a specific chemical structure. The present invention has been accomplished on the basis of this discovery.

2

Thus, the present invention provides a water-soluble monoazodye having the following formula (I) in its free acid form:

5

(I)




10

wherein X is  $\text{CH}=\text{CH}_2$  or  $-\text{CH}_2\text{CH}_2\text{W}$  wherein W is a leaving group removable by alkali;  $\text{R}^1$  is methyl,  $-\text{NHCONH}_2$  or  $-\text{NHCOT}^1$  wherein  $\text{T}^1$  is methyl, ethyl,  $-\text{CH}_2\text{CH}_2\text{COOH}$  or  $-\text{CH}=\text{CHCOOH}$ ;  $\text{R}^2$  is hydrogen, lower alkyl or lower alkoxy; and Z is

15

20

25



wherein P is cyano or halogen, each of Q and U which are independent of each other, is halogen, V is halogen or

30

35

40

45

50

55

60

65

70

75

80

85

90

95

100

105

110

115

120

125

130

135

140

145

150

155

160

165

170

175

180

185

190

195

200

205

210

215

220

225

230

235


240

245

250

255

260



55

60

65

70

75

80

85

90

95

100

105

110


115

120

125

130

135



(1)

60

65

70

75

80

85

90

95

100

105

110

115

120

125

130

135

140

145

150

155

160

165

170

175

180

185

190

195

200

205

210

215

220

225

230

235

240

245

250

255

260

265

270

275

280

285

290

295

300

305

310

315

320

325

330

335

340

345

350

355

360

365

370

375

380

385

390

395

400

405

410

415

420

425

430

435

440

445

450

455

460

465

470

475

480

485

490

495

500

505

510

515

520

525

530

535

540

545

550

555

560

565

570

575

580

585

590

595

600

605

610

615

620

625

630

635

640

645

650

655

660

665

670

675

680

685

690

695

700

705

710

715

720

725

730

735

740

745

750

755

760

765

770

775

780

785

790

795

800

805

810

815

820

825

830

835

840

845

850

855

860

865

870

875

880

885

890

895

900

905

910

915

920

925

930

935

940

945

950

955

960

965

970

975

980

985


990

995

1000

3

-continued



(2)


wherein R<sup>3</sup> is hydrogen or lower alkyl, A is a C<sub>1</sub>-C<sub>4</sub> alkyl or naphthalene group substituted by at least one selected from the group consisting of a sulfonic acid group and a carboxylic acid group, or phenyl substituted by at least one selected from the group consisting of a sulfonic acid group and a carboxylic acid group, and which may be substituted by at least one selected from the group consisting of hydroxyl, halogen, C<sub>1</sub>-C<sub>2</sub> alkyl and C<sub>1</sub>-C<sub>2</sub> alkoxy, R<sub>4</sub> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl, E is phenylene unsubstituted or substituted by lower alkyl, lower alkoxy, halogen or a sulfonic acid group, a naphthalene group unsubstituted or substituted by a sulfonic acid group, C<sub>1</sub>-C<sub>4</sub> alkylene or C<sub>1</sub>-C<sub>4</sub> alkyleneoxy C<sub>1</sub>-C<sub>4</sub> alkylene, and X is as defined above.

In the above water-soluble monoazodye of the present invention, X in the formula (I) represents —CH=CH<sub>2</sub> or —CH<sub>2</sub>CH<sub>2</sub>W. However, when X is —CH<sub>2</sub>CH<sub>2</sub>W, it will be converted to —CH=CH<sub>2</sub> by the removal of W-H by alkali. Thus, these two groups may be regarded as equivalent.

The group represented by W which is removable by the action of alkali, may, for example, be a sulfuric acid ester group, a thiosulfuric acid ester group, a phosphoric acid ester group, an acetic acid ester group or a halogen atom. Particularly preferred is a sulfuric acid ester group.

Y in the formula (I) is halogen or a group of the above formula (1) or (2).

When Y is halogen, the halogen includes fluorine, chlorine and bromine. However, when V is halogen, Y is preferably chlorine, and when V is




Y is preferably fluorine.

When Y is the group of the above formula (1), R<sub>4</sub> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl, preferably hydrogen, methyl or ethyl.

W in the group Y has the same meaning as the leaving group removable by alkali described above. W in the group X and W in the group Y may be the same or different.


When P, Q or u is halogen, such halogen includes fluorine, chlorine and bromine, preferably chlorine and fluorine.

Specific examples of




4

include:




Particularly preferred are:




The halogen for V in the formula (I) includes fluorine, chlorine and bromine. However, when Y is halogen, V is preferably chlorine, and when Y is other than halogen, V is preferably chlorine or fluorine.

Specific examples of the group




wherein A is a C<sub>1</sub>-C<sub>4</sub> alkyl or naphthalene group substituted by at least one selected from the group consisting of a sulfonic acid group and a carboxylic acid group, include the following groups:











5

-continued






Specific examples of the same group wherein A is phenyl substituted by at least one selected from the group consisting of a sulfonic acid group and a carboxylic acid group, and which may be substituted by at least one selected from the group consisting of hydroxyl, halogen, C<sub>1</sub>-C<sub>2</sub> alkyl and C<sub>1</sub>-C<sub>2</sub> alkoxy, include the following groups:



6

-continued



5

10

15

20

25

30

35

40

45


50

55




60

65




Among them, the following groups are particularly preferred for



when Y is halogen:




For E, the phenylene unsubstituted or substituted by lower alkyl, lower alkoxy, halogen or a sulfonic acid group and the naphthalene group unsubstituted or substituted by a sulfonic acid group include the following groups:




7

-continued




8

-continued




In the above formulas, the bond indicated by \* is the bond which is connected to the




group. Further, the C<sub>1</sub>-C<sub>4</sub> alkylene or the C<sub>1</sub>-C<sub>4</sub> alkyleneoxy C<sub>1</sub>-C<sub>4</sub> alkylene for E may, for example, be the following groups:

- β-(β-chloroethylsulfonyl)ethyl,
- γ-(β-chloroethylsulfonyl)propyl,
- δ-(β-chloroethylsulfonyl)butyl,
- γ-(β-sulfatoethylsulfonyl)propyl,
- β-(vinylsulfonyl)ethyl,
- β-[β-(β-chloroethylsulfonyl)ethoxy]ethyl,
- γ-[γ-(β-sulfatoethylsulfonyl)propoxy]propyl, and
- β-[β-(vinylsulfonyl)ethoxy]ethyl.

When V is halogen,




is preferably a group wherein R<sub>4</sub> is hydrogen and E is phenylene, C<sub>1</sub>-C<sub>4</sub> alkylene or C<sub>1</sub>-C<sub>4</sub> alkyleneoxy C<sub>1</sub>-C<sub>4</sub> alkylene, such as




In the formula (I), R<sup>1</sup> is methyl, —NHCONH<sub>2</sub> or —NHCOT<sup>1</sup> wherein T<sup>1</sup> is methyl, ethyl, —CH<sub>2</sub>CH<sub>2</sub>COOH or —CH=CHCOOH. The monoazo dye wherein R<sup>1</sup> is —NHCONH<sub>2</sub> is particularly excellent in that an object dyed therewith undergoes no color change even against a strong alkaline aqueous solution at the time of the post mercerizing treatment, and that it has no thermotropy.

Further, when Z is



wherein P, Q and U are as defined above



wherein each of V' and Y' is halogen, R<sup>1</sup> is preferably —NHCOCH<sub>2</sub>CH<sub>2</sub>COOH or —NHCOCH=CH<sub>2</sub>COOH with a view to improvement of the solubility of the dye.

5 For R<sup>2</sup>, the lower alkyl is preferably methyl or ethyl, and the lower alkoxy is preferably methoxy or ethoxy.


For R<sup>3</sup>, the lower alkyl is usually methyl or ethyl.

10 In the water-soluble monoazodye of the present invention, —SO<sub>3</sub>H and —SO<sub>2</sub>X in the diazo component in the above formula (I) may be substituted at any positions of the o-, m- and p-positions. It is preferred that one of them is located at the p-position and the other is located at the o-position.


15 The water-soluble monoazodye of the present invention may be used in the free acid form or in the form of its salt. The salt may, for example, be an alkali metal salt or an alkaline earth metal salt. Particularly preferred is a lithium salt, a sodium salt or a potassium salt.

20 The most preferred dyes of the present invention are represented by the following formulas (I-a), (I-b) and (z-c).


(I-a)



wherein X' and R<sup>2'</sup> have the same meanings as the above X and R<sup>2</sup>, respectively, A' is




and Y' is fluorine or chlorine.



11


wherein X<sup>n</sup> and R<sup>2n</sup> have the same meanings as the above X and R<sup>2</sup> respectively, R<sup>1n</sup> is [—NHCH<sub>2</sub>CH<sub>2</sub>COOH or —NHCH=CHCOOH] NHC<sub>2</sub>H<sub>4</sub>COOH or —NHC<sub>2</sub>H<sub>4</sub>COOH, and V<sup>n</sup> is fluorine or chlorine.




wherein X<sup>m</sup> and R<sup>2m</sup> have the same meanings as the above X and R<sup>2</sup>, respectively, R<sup>1m</sup> is [—NHCH<sub>2</sub>CH<sub>2</sub>COOH or —NHCH=CHCOOH] NHC<sub>2</sub>H<sub>4</sub>COOH or —NHC<sub>2</sub>H<sub>4</sub>COOH, and when P<sup>m</sup> is chlorine, each of Q<sup>m</sup> and U<sup>m</sup> is fluorine, and when P<sup>m</sup> is cyano, each of Q<sup>m</sup> and U<sup>m</sup> is chlorine.

The water-soluble monoazodye of the present invention can be produced as follows.


Namely, an azo compound of the following formula (II):




wherein R<sup>1</sup>, R<sup>2</sup> and X are as defined above, is condensed with a reactive compound of the following formula (III):



wherein Z<sup>1</sup> is




wherein P, Q and U are as defined above or




wherein Hal is halogen, and Hal is halogen in an aqueous medium at a temperature of from 0° to 15° C., whereby a water-soluble monoazodye of the following formula (IV):

12

(IV)




wherein R<sup>1</sup>, R<sup>2</sup>, X and Z<sup>1</sup> are as defined above, is produced. In a case where Z in the formula (I) is




wherein one of V or Y is halogen, an azo compound of the following formula (V):

(V)




wherein R<sup>1</sup>, R<sup>2</sup>, X and Hal are as defined above, is condensed with an amine of the following formula (VI), (VII) or (VIII):


(VI)



(VII)




(VIII)




wherein R<sup>3</sup>, R<sup>4</sup>, A, E and X are as defined above, in an aqueous medium at a temperature of from 30 to 40° C., whereby a water-soluble monoazodye of the following formula (IX) is obtained:

(IX)




65


wherein  $R^1, R^2$  are as defined above, and when  $y^2$  is halogen,  $V^2$  is



when  $y^2$  is the group of the formula (1) or (2),  $V^2$  is halogen. Further, to produce a water-soluble monoazodye of the formula (I) wherein  $V$  is



and  $Y$  is the group of the formula (1) or (2), a monoazodye of the following formula (X):



wherein  $R^1, R^2, R^3, A$  and  $X$  are as defined above, and  $y^3$  is halogen, is condensed with an amino compound of the above formula (VII) or (VIII) in an aqueous medium at a temperature of from  $85^\circ$  to  $95^\circ$  C. at a pH of from 4 to 6.

The feature in the chemical structure of the water-soluble monoazodye of the present invention resides in that the diazo component has a phenyl group substituted by  $-\text{SO}_3\text{H}$  and  $-\text{SO}_2\text{X}$  and the coupling component has a halotriazinyl group, a pyrimidinyl group or a triazinyl group substituted by the group of the formula (1) or (2). By virtue of this characteristic chemical structure, the build-up property and the level dyeing property at the time of dyeing are remarkably improved.

The water-soluble monoazodye of the present invention can be widely used as a dyestuff for dyeing fibers and cloths, a colorant for tinting paper and synthetic resins and as a colorant for e.g. an ink for an ink jet printer. However, it is particularly suitable for use as a dyestuff.

When the water-soluble monoazodye of the present invention is to be used as a dyestuff, the fibers to be dyed may, for example, be cellulose fibers such as cotton, viscose rayon, cupra ammonium rayon or kemp, or nitrogen-containing fibers such as polyamide, wool or silk. Particularly preferred are cellulose fibers. Such fibers may be used in the form of mixed fibers with e.g. polyester, triacetate or polyacrylonitrile.

To dye cellulose fibers or nitrogen-containing fibers by means of the water-soluble monoazodye of the present invention, it is common to employ an acid-binding agent, for example, an inorganic alkali such as sodium hydrogencarbonate or sodium carbonate, or an organic base such as triethylamine. The dyeing can be conducted in the presence of such an acid-binding agent in accordance with a conventional method.

The water-soluble monoazodye of the present invention may be used in combination with the other water-soluble dye.

As the dyeing method, a dipping method is particularly preferred, and the dyeing temperature is usually from  $40^\circ$  to  $80^\circ$  C.



Further, the water-soluble monoazodye of the present invention can be applied also to other dyeing method such as a cold pad batch method, a pad steam method, or a textile printing method.

The water-soluble monoazodye of the formula (I) of the present invention is particularly suitable for dyeing cellulose fibers or nitrogen-containing fibers to a yellow to orange color, and it is excellent in the buildup property. Further, the dyed products are excellent not only in the chlorine-fastness and the light-fastness but also in the level dyeing property.

Now, the present invention will be described in further detail with reference to Examples. However, it should be understood that the present invention is by no means restricted to such specific Examples.

EXAMPLE 1 and COMPARATIVE EXAMPLE 1

0.3 g or 1.2 g of each of monoazodyes of the following formulas (I-1) and (C):






## 15

was dissolved in 300 ml of water, and 20 g of Glauber's salt was added thereto to obtain a dye bath. In this dye bath, 15 g of a non-mercerized cotton cloth was dipped, and the temperature was raised to 60° C. over a period of 30 minutes. Then, 4.5 g of sodium carbonate was added thereto, and dyeing was conducted at 60° C. for one hour, followed by washing with water, soaping, washing with water and drying to obtain a yellow dyed cloth.

Here, the monoazodye of the above formula (I-1) was prepared by reacting 1 mol of a monoazo compound of the following formula (II-1):



(II-1)


with 1 mol of 3-sulfoaniline in an aqueous medium at 35° C. for 4 hours while maintaining the pH at a level of from 5.0 to 5.5, followed by salting out with potassium chloride. [max: 404 nm (water)]

Measurement of the build-up property

With respect to each dyed cloth, its surface reflectance was measured by a color difference meter (Macbeth® spec

## EXAMPLE 2

0.3 g or 0.6 g of a monoazodye [max: 412 nm (water)] of the following formula (I-2):



(I-2)

trometer MS-2020, manufactured by a division of Kolmorgen Corporation) to obtain a percentage exhaustion. The build-up property was represented by a relative value of the dyed density obtained by using 1.2 g of the dye to the dyed density obtained by using 0.3 g of the dye.

Level dyeing property

With respect to the dyed cloth obtained by using 0.3 g of each monoazodye, the uniformity in dyeing was visually evaluated.

The results are shown in Table 1.

TABLE 1

	Monoazodye	Build-up property (1.2 g/0.3 g)	Level dyeing property
Example 1	(I-1)	250	Remarkably uniformly dyed
Comparative Example 1	(C)	150	Non-uniformity in the dyed density observed

5

10

30

55

60


65

was dissolved in 300 ml of water, and 20 g of Glauber's salt was added thereto to obtain a dye bath. In this dye bath, 15 g of a cotton cloth was dipped, and the temperature was raised to 60° C. over a period of 30 minutes. Then, 4.5 g of sodium carbonate was added thereto, and dyeing was conducted at 60° C. for one hour, followed by washing with water, soaping, washing with water and drying to obtain a yellow dyed product having a good level dyeing property. Further, the build-up property was also good.

## 17

Here, the monoazodye of the above formula (I-2) was prepared as follows.

Namely, 1 mol of a monoazodye of the above formula (II-1) was reacted with 1 mol of 2-aminoethane sulfonic acid in an aqueous medium at a temperature of from 30 to 40° C. under stirring for 10 hours while maintaining the pH at a level of 8 with sodium carbonate to obtain a compound of the following formula:



Then, this compound was reacted with 1.2 mol of 3-(βhydroxyethyl)sulfonylaniline sulfuric acid ester in an aqueous medium at a temperature of from 90 to 95° C. under

stirring for 10 hours while maintaining the pH at a level of from 5 to 6 with sodium acetate, followed by salting out with potassium chloride to obtain the monoazodye of the above formula (I-2).

## EXAMPLE 3

The monoazodyes shown in the following Tables 2 and 3 (in their free acid forms) were prepared in accordance with the method of Example 1 or 2, and cotton cloths were dyed

with them in the same manner as in Example 1, whereby their level dyeing properties were excellent, and their build-up properties were also excellent.

TABLE 2

(f)













No.	XO <sub>2</sub> S	R <sup>1</sup>	R <sup>2</sup>	Y	V	λ <sub>max</sub> nm (Water)	Color of dyed cloth
3-1			H	F		405	Yellow
3-2			H	Cl		405	Yellow
3-3			H	Cl		405	Yellow
3-4			H	F		408	Yellow

TABLE 2-continued

(1)









No.	R <sup>1</sup>	R <sup>2</sup>	Y	V	λ <sub>max</sub> nm (Water)	Color of dyed cloth
3-5	 $\text{HO}_3\text{SOH}_4\text{C}_2\text{O}_2\text{S}$				413	Yellow
3-6		—H		$\text{—NHC}_3\text{H}_6\text{SO}_3\text{H}$	415	Yellow
3-7		—H	—Cl	$\text{—NHCHCH}_2\text{COOH}$   COOH	408	Yellow
3-8		—CH <sub>3</sub>	—F	$\text{—NC}_2\text{H}_4\text{COOH}$   CH <sub>3</sub>	410	Yellow

TABLE 2-continued

(1)

No.				$\lambda_{max}$ nm (Water)	Color of dyed cloth
3-9				432	Orange
3-10				428	Yellow
3-11				403	Yellow

TABLE 2-continued

(1)

No.	$-R^1$	$-R^2$	$-Y$	$-V$	$\lambda_{max}$ nm (Water)	Color of dyed cloth
3-12		$-H$	$-F$	$-V$	403	Yellow
3-13		$-H$	$-F$	$-V$	430	Orange
3-14		$-H$	$-F$	$-V$	407	Yellow
3-15		$-H$	$-Cl$	$-V$	405	Yellow








TABLE 2-continued

(1)










No.		$-R^1$	$-R^2$	$-Y$	$-V$	$\lambda_{max}$ nm (Water)	Color of dyed cloth
3-16		$-NHCONH_2$	$-H$	$-Cl$		403	Yellow
3-17		$-NHCONH_2$	$-H$	$-Cl$		402	Yellow
3-18		$-NHCONH_2$	$-H$	$-F$		402	Yellow
3-19		$-CH_3$	$-H$	$-Cl$		391	Yellow

TABLE 2-continued

(1)









No.			$\lambda_{max}$ nm (Water)	Color of dyed cloth
3-20	<p><math>XO_2S</math></p>  <p><math>SO_3H</math></p>	<p>—R<sup>1</sup></p> <p>—CH<sub>3</sub></p>  <p><math>SO_3H</math></p> <p><math>HO_3SOH_4C_2O_2S</math></p>	<p>—R<sup>2</sup> —Y</p> <p>—OCH<sub>3</sub> —F</p>	<p>—V</p> <p>418</p> <p>Yellow</p>
3-21	<p><math>HO_3SOH_4C_2O_2S</math></p>  <p><math>SO_3H</math></p> <p>—NHCONH<sub>2</sub></p>	<p>—H</p> <p>—Cl</p>  <p><math>C_2H_5</math></p> <p><math>SO_3H</math></p>	<p>—H</p> <p>—Cl</p>	<p>403</p> <p>Yellow</p>
3-22	<p><math>HO_3SOH_4C_2O_2S</math></p>  <p><math>SO_3H</math></p> <p>—NHCONH<sub>2</sub></p>	<p>—H</p> <p>—Cl</p>  <p><math>SO_3H</math></p> <p><math>SO_3H</math></p>	<p>—H</p> <p>—Cl</p>	<p>403</p> <p>Yellow</p>



TABLE 2-continued

(1)

No.						$\lambda_{max}$ nm (Water)	Color of dyed cloth
3-23	$XO_2S$	$SO_3H$	$-NHCONH_2$	$-V$		403	Yellow
3-24	$HO_3SOH_4C_2O_2S$	$SO_3H$	$-NHCONH_2$	$-H$		417	Yellow
3-25	$HO_3SOH_4C_2O_2S$	$SO_3H$	$-NHCOCH_2CH_2COOH$	$-OCH_3$	$-Cl$	420	Yellow
3-26	$HO_3SOH_4C_2O_2S$	$SO_3H$	$-NHCOCH=CHCOOH$	$H$	$-NHC_2H_4OC_2H_4SO_2C_2H_4OSO_3H$	409	Yellow

TABLE 2-continued

(1)











No.	$R^1$	$R^2$	$Y$	$V$	$\lambda_{max}$ nm (Water)	Color of dyed cloth	
3-27			H			419	Yellow
3-28			$-NHCOCH_3$	$-Cl$	429	Orange	
3-29		$-H$		$-Cl$	407	Yellow	
3-30		$-H$		$-F$	413	Yellow	

TABLE 2-continued

(1)











No.		$-R^1$	$-R^2$	$-Y$	$-V$	$\lambda_{max}$ nm (Water)	Color of dyed cloth
3-31		$-NHCONH_2$	$-H$	$C_3H_7$ $-NC_2H_4SO_2C_2H_4Cl$	$-NHC_2H_4SO_3H$	414	Yellow
3-32		$-NHCONH_2$	$-H$			412	Yellow
3-33		$-NHCONH_2$	$-H$			414	Yellow
3-34		$-NHCOCH_2-CH_2COOH$	$-CH_3$		$-Cl$	411	Yellow

TABLE 2-continued

(1)











No.	$R^1$	$R^2$	$Y$	$V$	$\lambda_{max}$ nm (Water)	Color of dyed cloth
3-35	 $XO_2S$	 $SO_2X$	 $SO_3H$	 $-V$	433	Orange
3-36	 $HO_3SOH_4C_2O_2S$	H	 $C_2H_4SO_2C_2H_4Br$	 $C_2H_4SO_2C_2H_4Br$	432	Orange
3-37	 $HO_3SOH_4C_2O_2S$	H	 $SO_3H$	 $SO_2C_2H_4OSO_3H$	402	Yellow

TABLE 2-continued

(1)







No.	R <sup>1</sup>	R <sup>2</sup>	—Y	Cl	λ <sub>max</sub> nm (Water)	Color of dyed cloth
3-38			—R <sup>2</sup>	—V	413	Yellow
3-39		—NHCOCH <sub>2</sub> —CH <sub>2</sub> COOH	—OCH <sub>3</sub>	—F	426	Orange
3-40		—NHCOCH=CHCOOH	—OCH <sub>3</sub>		434	Orange
						

TABLE 2-continued

(1)















No.	$R^1$	$R^2$	$Y$	$V$	$\lambda_{max}$ nm (Water)	Color of dyed cloth
3-41	 $XO_2S-$	 $-R^2$	 $-Y$	 $-V$	425	Orange
3-42	 $HO_3SOH_4C_2O_2S-$	 $-OCH_3$	 $-OCH_3$	 $-F$	423	Orange

TABLE 3


No.		$-R^1$	$-R^2$	$-P$	$-Q$	$-U$	$\lambda_{max}$ nm (water)	Color of dyed cloth
3-43		$-NHCOCH=CHCOOH$	$-OC_2H_5$	$-Cl$	$-F$	$-F$	422	Yellow
3-44		$-NHCOCH_2-CH_2COOH$	$-OC_2H_5$	$-CN$	$Cl$	$-Cl$	423	Yellow
3-45		$-NHCOCH_2-CH_2COOH$	$-OCH_3$	$-Cl$	$-F$	$-F$	420	Yellow
3-46		$-NHCOCH=CHCOOH$	$-OCH_3$	$-Cl$	$-Cl$	$-Cl$	421	Yellow

What is claimed is:

[1. A water-soluble monoazo dye having the following formula (I) in its free acid form:



45




(I)

50

wherein V is halogen or

55




Y is halogen or a group of the following formula (1) or (2):

60

wherein X is  $-CH=CH_2$  or  $-CH_2CH_2W$  wherein W is a leaving group removable by alkali,  $R^1$  is methyl,  $-NHCONH_2$ , or  $-NHCOT^1$  wherein  $T^1$  is methyl, ethyl,  $-CH_2CH_2COOH$  or  $-CH=CHCOOH$ ;  $R^2$  is hydrogen, lower alkyl or lower alkoxy; and

(1)



65


45

-continued




wherein R<sup>3</sup> is hydrogen or lower alkyl, A is a C<sub>1</sub>-C<sub>4</sub> alkyl or naphthalene group substituted by at least one selected from the group consisting of a sulfonic acid group and a carboxylic acid group, or phenyl substituted by at least one selected from the group consisting of a sulfonic acid group and a carboxylic acid group, and which may be substituted by at least one selected from the group consisting of hydroxyl, halogen, C<sub>1</sub>-C<sub>2</sub> alkyl and C<sub>1</sub>-C<sub>2</sub> alkoxy, R<sub>4</sub> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl, E is phenylene unsubstituted or substituted by lower alkyl, lower alkoxy, halogen or a sulfonic acid group, a naphthalene group unsubstituted or substituted by a sulfonic acid group, C<sub>1</sub>-C<sub>4</sub> alkylene or C<sub>1</sub>-C<sub>4</sub> alkyleneoxy C<sub>1</sub>-C<sub>4</sub> alkylene, and X is as defined above.]

2. A water-soluble monoazo dye having the following formula (I) in its free acid form:




wherein X is -CH=CH<sub>2</sub> or -CH<sub>2</sub>CH<sub>2</sub>W wherein W is a leaving group removable by alkali; R<sup>1</sup> is -NHCOT<sup>1</sup> wherein T<sup>1</sup> is -CH<sub>2</sub>CH<sub>2</sub>COOH or -CH=CHCOOH; R<sup>2</sup> is hydrogen, lower alkyl or lower alkoxy; and Z is




wherein P is cyano or halogen, each of Q and U which are independent of each other, is halogen.

[3. The monoazo dye according to claim 1, wherein V is




[4. The monoazo dye according to claim 1, wherein V is




[5. The monoazo dye according to claim 1, wherein Y is fluorine or chlorine, and V is

46




[6. The monoazo dye according to claim 1, wherein R<sup>1</sup> is -NHCONH<sub>2</sub>.]


[7. The monoazo dye according to claim 1, wherein Y is




[8. The monoazo dye according to claim 7, wherein E is



[9. The monoazo dye according to claim 1, wherein Y is




[10. The monoazo dye according to claim 1, wherein Y is



and V is fluorine or chlorine.]

[11. The monoazo dye according to claim 1, wherein Y is




V is fluorine or chlorine, and R<sup>1</sup> is -NHCOCH<sub>3</sub>, -NHCOCH<sub>2</sub>CH<sub>2</sub>COOH or -NHCOCH=CH-COOH.]

[12. The monoazo dye according to claim 11, wherein R<sup>1</sup> is -NHCOCH<sub>2</sub>CH<sub>2</sub>COOH or -NHCOCH=CH-COOH.]

13. A water-soluble monoazo dye, wherein said dye is a compound having the following formula:



47



wherein  $X'''$  is  $-CH=CH_2$  or  $-CH_2CH_2W$  wherein  $W$  is a leaving group removable by alkali;

48

$R^{2'''}$  is hydrogen, lower alkyl or lower alkoxy;

$R^{1'''}$  is  $-NHCOOH_2CH_2COOH$  or  $-NHCOCH=CHCOOH$ ;


5  $P'''$  is chlorine or cyano;

$Q'''$  is fluorine or chlorine;

$U'''$  is fluorine or chlorine;

10 wherein when  $P'''$  is chlorine, each of  $Q'''$  and  $U'''$  is fluorine, and when  $P'''$  is cyano, each of  $Q'''$  and  $U'''$  is chlorine.

14. A monoazo dye according to claim 13, having the formula in its free acid form:



\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : Re. 37,004

DATED : December 26, 2000

INVENTOR(S) : Yousuke TAKAHASHI, et al.

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title page Item [75], the 1<sup>st</sup> inventor's residence; and  
and the 2<sup>nd</sup> inventor's name and residence are listed incorrectly.

Item [75] should read as follows:

[75] Inventors: **Yousuke Takahashi**, Nishinomiya;  
**Yukiharu Shimizu**, Hasaki-machi;  
**Toshio Hihara**, Kitakyushu; **Kiyoshi  
Himeno**, Munakata, all of Japan

Signed and Sealed this

Twenty-ninth Day of May, 2001

Attest:



NICHOLAS P. GODICI

Attesting Officer

Acting Director of the United States Patent and Trademark Office