



US00RE37936E

(19) **United States**
 (12) **Reissued Patent**
Huang et al.

(10) **Patent Number: US RE37,936 E**
 (45) **Date of Reissued Patent: Dec. 10, 2002**

(54) **PESTICIDAL 1-POLYARYLPYRAZOLES**

(75) Inventors: **Jamin Huang**, Chapel Hill, NC (US);
Scot Kevin Huber, Raleigh, NC (US)

(73) Assignee: **Aventis CropScience USA Inc.**,
 Research Triangle Park, NC (US)

(21) Appl. No.: **09/903,990**

(22) Filed: **Jul. 13, 2001**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **5,922,884**
 Issued: **Jul. 13, 1999**
 Appl. No.: **08/963,631**
 Filed: **Nov. 4, 1997**

U.S. Applications:

(60) Provisional application No. 60/030,128, filed on Nov. 4, 1996.

(51) **Int. Cl.⁷** **A01N 43/56; C07D 231/44**

(52) **U.S. Cl.** **548/367.4; 514/404; 514/407**

(58) **Field of Search** **548/367.4; 514/404, 514/407**

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,614,533	A	9/1986	Schallner et al.	71/92
4,771,066	A	9/1988	Gehring et al.	514/404
4,804,675	A	2/1989	Jensen-Korte et al.	514/407
4,891,057	A	1/1990	Sohn et al.	71/72
4,945,165	A	7/1990	Jensen-Korte et al.	548/362
5,082,949	A	1/1992	Sohn et al.	548/378
5,187,185	A	2/1993	Outcalt et al.	514/408
5,223,525	A	6/1993	Wu et al.	514/398
5,232,940	A	8/1993	Hatton et al.	514/407
5,236,938	A	8/1993	Huang et al.	514/341

FOREIGN PATENT DOCUMENTS

DE 3633840 4/1988


DE	19511269	10/1995
EP	0154115	9/1985
EP	0201852	11/1986
EP	0235628	9/1987
EP	0285893	10/1988
EP	0295117	12/1988
EP	0385809	9/1990
EP	0403300	12/1990
EP	0500209	8/1992
EP	0679650	11/1995
EP	0780381	6/1997
WO	87/03781	7/1987
WO	93/06089	4/1993
WO	94/21606	9/1994
WO	96/25401	8/1996

Primary Examiner—Robert W. Ramsuer

(74) *Attorney, Agent, or Firm*—Burns, Doane, Swecker & Mathis, L.L.P.

(57) **ABSTRACT**

A compound of formula (I):



compositions containing them and methods of use to control pests.

21 Claims, No Drawings

PESTICIDAL 1-POLYARYLPYRAZOLES

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.

CROSS-REFERENCE TO RELATED APPLICATION

This application claims the priority of U.S. Provisional Patent Application No. 60/030,128, filed Nov. 4, 1996, incorporated by reference herein in its entirety and relied upon.

The invention relates to new 1-arylpyrazoles and derivatives thereof which have some valuable properties either as pesticides or as intermediates to make other pesticides. The invention further pertains to compositions of said compounds and methods, using said compounds either as intermediates to make other pesticides, or for the control of pests particularly insects, in particular to the application of said compounds or compositions in agricultural methods of use or for animal protection, particularly as pesticides, for controlling arthropods.

International Patent Publication No. WO 87/03781 and European Patent Publication No. 295117, 154115, 201852 describe insecticidal 1-(substituted phenyl)pyrazoles. Other prior art is also found in the text of these patent applications or the patents issued therefrom.

International Patent Publications No. WO 93/06089 and WO 94/21606 also describe insecticidal 1-(4-SF₅ substituted phenyl) heterocycles which may be pyrroles as well as imidazoles or pyrazoles. The teaching of these patents is not substantially different from International Patent Publication No. WO 87/03781 or from European Patent Publication No. 0295117 as far as pyrazoles are concerned.

It is an object of the present invention to provide new pesticidal compounds of the 1-arylpyrazole family together with processes for their preparation.

Another object of the present invention is to provide pesticidal compositions and pesticidal methods of use of the pesticidal pyrazole compounds against arthropods, especially insects, particularly in agricultural or horticultural crops, forestry, veterinary medicine or livestock husbandry, or in public health.


A third object of the present invention is to provide very active compounds with broad spectrum pesticidal activity, as well as compounds with selective special activity, e.g., aphicidal, miticidal, foliar insecticidal, soil insecticidal, systemic, antifeeding or pesticidal activity via seed treatment.

A fourth object of the present invention is to provide compounds with substantially enhanced and more rapid activity, especially against insects and more particularly insects in their larval stages.

A fifth objective of the present invention is to provide compounds with greatly improved (faster and greater) penetration into pest species when topically applied and to thus provide enhanced movement of the compounds to the pesticidal site(s) of action within the pest.

These and other objectives of the invention are met in whole or part and shall become readily apparent from the description of the present invention which follows.

The invention thus relates to compounds having the general formula (I):



wherein

X is N or C—R₂;

Y is N or C—R₃;

W is N or C—R₄;

R₂ and R₃ are independently selected from H, halogen, hydroxy, alkyl, cycloalkyl, haloalkyl, alkoxy, haloalkoxy, CN, NO₂, —S(O)_nR₈, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, hydroxycarbonyl, aminosulfonyl, alkylaminosulfonyl, and dialkylaminosulfonyl;

R₄ is H, halogen, alkyl, alkoxy, CN, NO₂, haloalkyl, haloalkoxy, thiocyanato, formyl, alkylcarbonyl, —CH=N—OH, [—CH=N—O-alkyl] —CH=N—O-alkyl, —S(NH₂)(=NH), —S(O)_nR₈, mercapto, haloalkylcarbonyl, or a —S—radical so that two molecules are bound together to form a disulfide compound;

R₅ is hydrogen, halogen, —NR₉R₁₀, —N=CR₁₁R₁₉, —S(O)_nR₈, formyl, alkylcarbonyl, haloalkylcarbonyl, cyano, alkyl, haloalkyl, hydrazino, alkoxy carbonyl, alkylthiocarbonyl, 1H-pyrrol-1-yl or 1H-pyrazol-1-yl;

R₈ is alkyl or haloalkyl, alkenyl or alkynyl, or a cycloalkyl ring containing 3 to 5 carbon atoms;

R₁₁ is H, or alkyl;

R₁₉ may also be hydrogen, alkyl, haloalkyl, alkoxy, haloalkoxy, alkylthio, amino, monoalkylamino, dialkylamino, or R₁₉ is phenyl, thienyl, pyridyl or furyl, each of which is unsubstituted or substituted with alkyl, haloalkyl, halogen, NO₂, CN, alkoxy, haloalkoxy, OH, alkylcarbonyl, alkylcarbonyloxy;

R₉ and R₁₀ independent of one another, are H, alkyl, haloalkyl, alkylcarbonyl, haloalkylcarbonyl, R₈S(O)_n, formyl, alkenyl, alkynyl, alkoxy carbonyl, alkylthiocarbonyl, aroyl; or are joined so as together form a divalent radical having 4 to 6 atoms in the chain, this divalent radical being alkylene, alkyleneoxyalkylene or alkyleneaminoalkylene, preferably to form a morpholine, pyrrolidine, piperidine or piperazine ring; the alkyl portion of R₉ and R₁₀ may be substituted by R₇;

R₇ is cyano, nitro, alkoxy, haloalkoxy, R₈S(O)_n, —C(O) alkyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, —CO₂H, halogen, hydroxy, aminosulfonyl, alkylaminosulfonyl or dialkylaminosulfonyl;

Z is N or C—R₁₆;

n is zero, one or two;

R_{12} , R_{13} , R_{15} , R_{16} , are independently selected from hydrogen, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, alkylsulfenyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfenyl, haloalkylsulfinyl, haloalkylsulfonyl, formyl, alkylcarbonyl, alkoxycarbonyl;

R_{22} , R_{23} , R_{24} , R_{25} , R_{26} are *hydrogen*, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, haloalkylcarbonyl, formyl, alkylcarbonyl, thioamide, amide, and alkoxycarbonyl, SF_5 , $R_8S(O)_n$; preferably R_{24} is halogen, haloalkyl or haloalkoxy; or R_{22} and R_{23} or R_{23} and R_{24} or R_{25} and R_{26} may also be together a divinylidene group ($-CH=CH-CH=CH-$) or a methylene diether ($-O-CH_2-O-$) or halomethylene diether ($-O-CF_2-O-$), so as to form a cyclic ring vicinal to the phenyl ring; or pesticidally acceptable salts thereof.

By the term "pesticidally acceptable salts" is meant salts the anions and cations of which are known and accepted in the art for the formation of pesticidally acceptable salts. Preferably such salts are water soluble. Suitable acid addition salts formed from compounds of formula (I) containing an amine group, include salts with inorganic acids for example hydrochlorides, phosphates, sulfates and nitrates, and salts with organic acids for example acetates. Suitable salts with bases formed from compounds of formula (I) containing a suitably acidic group include alkali metal (for example sodium or potassium) salts, ammonium salts and organic amine (for example diethanolamine or morpholine) salts.

In the present invention, some words are used in a specific sense:

The term "lower alkyl- $S(O)_n$ " means a radical of the formula $-S(O)_n-$ lower alkyl. The term " $R_{10}S(O)_n$ " means a radical of the formula $-S(O)_nR_{10}$. The term "aminocarbonyl" means a carbamoyl radical, that is, a radical of the formula $-C(O)NH_2$. Similarly, the term "alkylaminocarbonyl" means an [alkylcarbomoyl] *alkylcarbomoyl* radical, that is, a radical of the formula $-C(O)-NH-alkyl$; and the term "dialkylaminocarbonyl" means a dialkylcarbomoyl radical, that is, a radical of the formula $-C(O)-N(alkyl)_2$ in which the alkyl moieties can be the same or different. The term "aminosulfonyl" means a sulfamoyl radical, that is, $-SO_2NH_2$. Similarly, the term "alkylaminosulfonyl" means an alkylsulfamoyl radical, that is, a radical of the formula $-SO_2NH-alkyl$; while the term "dialkylaminosulfonyl" means a dialkylsulfamoyl radical, which has the formula $-SO_2N(alkyl)_2$ wherein the alkyl moieties can be the same or different.

The term "halo" before the name of a radical means that this radical is partially or completely halogenated, that is to say, substituted by F, Cl, Br, or I, in any combination, preferably by F or Cl. The term "halogen" means F, Cl, Br or I. The term "lower" before the name of a radical having a carbon skeleton means that this carbon skeleton has less than 6 carbon atoms. When the name of any substituent is repeated, it keeps the same meaning unless otherwise specified. The term "aryl" designates a carbon and/or heteroatom-containing aromatic radical which is preferably phenyl optionally substituted with one or more substituents selected from halogen, methyl and methoxy, especially phenyl, halophenyl, tolyl or xylyl. The term "aroyl" designates a carbonyl aromatic radical, that is, aryl-C(O)—, which is preferably a benzoyl, methylbenzoyl, halobenzoyl or xylyl-carbonyl radical. The term "acyl" designates an alkylcarbonyl radical. The various individual radicals (such as alkyl,

alkenyl, alkynyl, alkoxy and alkylene or the like) generally contain up to six carbon atoms.

A preferred class of compounds of formula (I) is that wherein

X is N;

Y is C— R_3 ;

W is C— R_4 ;

R_3 is H, halogen, hydroxy, alkyl, cycloalkyl, haloalkyl, alkoxy, haloalkoxy, CN, NO_2 , $-S(O)_nR_8$, alkylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, hydroxycarbonyl, aminosulfonyl, alkylaminosulfonyl, dialkylaminosulfonyl;

R_4 is H, halogen, alkyl, alkoxy, CN, NO_2 , haloalkyl, haloalkoxy, thiocyanato, formyl, alkylcarbonyl, $-CH=N-OH$, $-CH=N-O-alkyl$, $-S(NH_2)(=NH)$, $-S(O)_nR_8$, mercapto, haloalkylcarbonyl, or a $-S-$ radical so that two molecules are bound together to form a disulfide compound;

R_5 is hydrogen, halogen, $-NR_9R_{10}$, $-N=CR_{11}R_{19}$, $-S(O)_nR_8$, formyl, alkylcarbonyl, haloalkylcarbonyl, cyano, lower alkyl, hydrazino, alkoxycarbonyl, alkylthiocarbonyl, 1H-pyrrol-1-yl or 1H-pyrazol-1-yl, preferably, amino $-NR_9R_{10}$;

R_8 is alkyl or haloalkyl, alkenyl or alkynyl; or a cycloalkyl ring containing 3 to 5 carbon atoms; preferably R_8 is lower alkyl;

R_{11} is H or alkyl;

R_{19} may also be hydrogen, alkyl, haloalkyl, alkoxy, haloalkoxy, alkylthio, amino, monoalkylamino, dialkylamino; or R_{19} is phenyl, thienyl, pyridyl or furyl, all of them being optionally substituted with alkyl, haloalkyl, halogen, NO_2 , CN, alkoxy, haloalkoxy, OH, alkylcarbonyl, alkylcarbonyloxy;

R_9 and R_{10} are H, alkyl, haloalkyl, alkylcarbonyl, haloalkylcarbonyl, $R_8-S(O)_n$, formyl, alkenyl, alkynyl, alkoxycarbonyl, alkylthiocarbonyl, aroyl; or are joined so as together form a divalent radical having 4 to 6 atoms in the chain, this divalent radical being alkylene, alkyleneoxyalkylene or alkyleneaminoalkylene, preferably to form a morpholine, pyrrolidine, piperidine or piperazine ring; the alkyl portion of R_9 and R_{10} may be substituted by R_7 ;

R_7 is cyano, nitro, alkoxy, haloalkoxy, $R_8S(O)_n$, $-C(O)alkyl$, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, $-CO_2H$, halogen, hydroxy, aminosulfonyl, alkylaminosulfonyl or dialkylaminosulfonyl;

Z is N or C— R_{16} ;

n is zero, one or two;

R_{12} , R_{13} , R_{15} , R_{16} , are separately hydrogen, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, alkylsulfenyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfenyl, haloalkylsulfinyl, haloalkylsulfonyl, formyl, alkylcarbonyl, alkoxycarbonyl;

R_{22} , R_{23} , R_{24} , R_{25} , R_{26} are *hydrogen*, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, haloalkylcarbonyl, formyl, alkylcarbonyl, thioamide, amide, and alkoxycarbonyl, SF_5 , $R_8S(O)_n$, preferably R_{24} is halogen, haloalkyl or haloalkoxy; or R_{22} and R_{23} or R_{23} and R_{24} or R_{25} and R_{26} may also be together a divinylidene group

5

(—CH=CH—CH=CH—) or a methylene diether (—O—CH₂—O—) or halomethylene diether (—O—CF₂—O—) so as to form a cyclic ring vicinal to the phenyl ring; or a pesticidally acceptable salt thereof.

Another preferred class of the compounds of formula (I) are those with one or more of the following features wherein:

X is N or C—R₂;

Y is N or C—R₃;

W is N or C—R₄;

R₃ is CN or halogen;

R₄ is H, halogen, formyl, or —S(O)_nR₈;

R₅ is hydrogen, halogen, C₁–C₃ alkyl, C₁–C₃ haloalkyl, or —NR₉R₁₀;

R₈ is methyl, ethyl, —CF₃, —CFCl₂, —CF₂Cl;

R₁₂ and R₁₆ are independently selected from F, Cl, Br and H;

R₁₃ and R₁₅ are H;

R₂₄ is —CF₃, —OCF₃, —CHF₂, —S(O)_nCF₃, —CFCl₂, —CF₂Cl, —OCF₂Cl, —OCFCl₂, Cl, Br or F; or

Z is CCl, CF, CBr or N.

A further especially preferred class of compounds are those wherein:

X is N;

Y is C—R₃;

W is C—R₄;

R₁₂ and R₁₆ are Cl or Br;

R₁₃ and R₁₅ are H;

R₂₄ is —CF₃, —OCF₃ or Br;

R₅ is amino;

R₉ and R₁₀ are H, alkyl or alkylcarbonyl;

R₈ is methyl or ethyl or CF₃, CCl₂F, CClF₂; and

R₃ is CN or halogen.


For the above preferred compounds, there are optimum combinations of substituent groups.

Further, preferred S(O)_nR₈ substituents in formula (I) are: methylthio, methylsulfinyl, methylsulfonyl, ethylsulfinyl, ethylsulfonyl, ethylthio, cyclopropylsulfinyl, cyclopropylthio, cyclopropylsulfonyl, isopropylsulfinyl, isopropylsulfonyl, isopropylthio, trifluoromethylthio, trifluoromethylsulfinyl, trifluoromethylsulfonyl, dichlorofluoromethylthio, dichloromethylsulfinyl, dichloromethylsulfonyl, chlorodifluoromethylthio, chlorodifluoromethylsulfonyl or chlorodifluoromethylsulfinyl.

Some compounds are useful as intermediates to make other pesticides, others are useful directly as pesticides. Compounds wherein R₂₃ or R₂₄ or R₂₅ is formyl are preferred as intermediates as well as compounds wherein R₅ is H or compounds wherein simultaneously R₄ is not halogenated and R₂₂ and R₂₃ and R₂₄ and R₂₅ and R₂₆ are H.

The compounds of general formula (I) can be prepared by the application or adaptation of known methods (i.e., methods heretofore used or described in the chemical literature including the Chemical Abstracts) employing as starting material the compounds of formula (II)

6




(II)

wherein the various substituents have the same meaning as in formula (I) and halo is a halogen atom, preferably bromine or iodine.

The compounds of formula (II) can be prepared by methods or processes similar to those described in International patent applications WO 87/03781, 93/06089, 94/21606, in European patent applications 295117, 403300, 385809, 0500209, 679650, 285,893, and 780,381, U.S. Pat. Nos. 5,232,940, 5,236,938, 5,187,185, 5,223,525 and German Patent application 19511269 or by other methods known to the skilled addressee. The skilled addressee understands and is generally knowledgeable of Chemical Abstracts.


According to a first method of preparation of compounds of formula (I) from compounds of formula (II), a compound of formula (II) is caused to react with a boric acid or ester, preferably in presence of a coupling catalyst, so as to form a compound of formula (III)



(III)

wherein the substituents have the same meaning as previously indicated, and B(OR₃₀)₂ represents a boric acid or ester group (R₃₀ is preferably hydrogen, alkyl, or a divalent lower alkylene radical such that two R₃₀O radicals may form a cyclic borate ester),

said compound (III) being, in a second step, caused to react with a compound of formula (VI)



(VI)

wherein R₂₀ is bromine or iodine or O—SO₂CF₃.


The first step of this first process is generally and preferably conducted in an organic solvent, for example an hydrocarbon such as toluene or xylene; an amide such as dimethylformamide or N-methylpyrrolidone; an ether such as tetrahydrofuran, dimethoxyethane or

7

2-methoxyethylether; the temperature is generally between 50° C. and 150° C.; utilizing as catalysts organic derivatives of palladium including palladium acetate, tetrakis(triphenylphosphine)palladium(0) or Pd₂(dibenzylidene acetone)₃, generally in the presence of a base such as an alkaline hydroxide or carbonate acetate ion or an amine. The boron derivative used as a reactant with the compound of formula (II) is preferably a cyclic diboron ester or acid of formula (R₃₀O)₂B—B(OR₃₀)₂.

The second step of this first process is advantageously conducted in an organic solvent, for example an hydrocarbon such as toluene or xylene; an amide such as dimethylformamide or N-methylpyrrolidone; an ether such as tetrahydrofuran, dimethoxyethane or 2-methoxyethylether; the temperature is generally between 50° C. and 150° C.; utilizing as catalysts organic derivatives of palladium including palladium acetate, tetrakis(triphenylphosphine)palladium(0) or Pd₂(dibenzylidene acetone)₃, generally in the presence of a base such as an alkaline hydroxide or carbonate acetate ion or an amine.

According to another method of preparation of compound of formula (I), a compound of formula (II) is caused to react with a compound of formula (IV):




(IV)

wherein the substituents have the same meaning as previously indicated.

The process is generally and preferably conducted in an organic solvent, for example an hydrocarbon such as toluene or xylene; an amide such as dimethylformamide or N-methylpyrrolidone; an ether such as tetrahydrofuran, dimethoxyethane or 2-methoxyethylether; the temperature is generally between 50° C. and 150° C.; utilizing as catalysts organic derivatives of palladium including palladium acetate, tetrakis(triphenylphosphine)palladium(0) or Pd₂(dibenzylidene acetone)₃, generally in the presence of a base such as an alkaline hydroxide or carbonate acetate ion or an amine.

According to still another method of preparation of compound of formula (I), a compound of formula (II) is caused to react in the presence of a catalyst as described above, with a hexaalkylstannane[(alkyl or cycloalkyl)₃Sn]₂ so as to form a compound of formula (V):



(V)

which in a second step is caused to react with a compound of formula (IV) in the presence of a coupling catalyst.


The first step of the process is generally conducted in an organic solvent, for example an hydrocarbon such as toluene

8

or xylene; an amide such as dimethylformamide or N-methylpyrrolidone; an ether such as tetrahydrofuran, dimethoxyethane or 2-methoxyethylether; at a temperature generally between 50° C. and 150° C.; utilizing as catalysts organic derivatives of palladium including palladium acetate, tetrakis(triphenylphosphine)palladium(0) or Pd₂(dibenzylidene acetone)₃.

The second step of this second process is generally conducted in an organic solvent, for example an hydrocarbon such as toluene or xylene; an amide such as dimethylformamide or N-methylpyrrolidone; an ether such as tetrahydrofuran, dimethoxyethane or 2-methoxyethylether; the temperature is generally between 50° C. and 150° C.; as catalysts, organic derivatives of palladium such as Pd(P-phenyl)₃, Pd₂(dibenzylidene acetone)₃, Pd(O—CO—CH₃)₂ may be cited.


According to still another method of preparation of compound of formula (I), a compound of formula (II) is caused to react with a compound of formula (VII)



(VII)

preferably in the presence of a coupling catalyst as described above, and preferably in a solvent as described above, and preferably at a temperature from 50° C. to 150° C.

According to another method of preparation of a compound of formula (I), a compound of formula (VIII):



(VIII)

35

40

45

50

55

60

is reacted in a step to form a pyrrole, imidazole, triazole, or pyrazole of formula (I). Such reactions are known and can be found in Katritzky, Rees, and Scriven, Comprehensive Heterocyclic Chemistry II, volumes 2,3,4 1996, Pergamon Press, London.

The invention is illustrated by the following examples, which are not considered as limiting the invention but are given to better enable use of it.

EXAMPLE 1

Preparation of 1-[2,6-dichloro-4-(4-trifluoromethylphenyl)phenyl]-3-cyano-4-trifluoromethylthio-5-aminopyrazole

A mixture of 6 g of [5-amino-3-cyano-1-(2,6-dichloro-4-bromophenyl)4-trifluoromethylthiopyrazole] 5-amino-3-cyano-1-(2,6-dichloro-4-bromophenyl)-4-

trifluoromethylthiopyrazole (prepared according to procedures reported in U.S. Pat. No. 5,232,940), 4-trifluoromethylphenylboronic acid (5.3 g), K_2CO_3 (5.8 g), tris(dibenzylidene acetone)dipalladium (0.6 g) and diglyme was heated at 130° C. for 25 hours. After cooling to 20° C., the mixture was poured into water and extracted with diethyl ether. The ether solutions were combined, dried, filtered and the filtrate concentrated and purified by chromatography. The desired product as a solid (3.9 g, mp 181–184° C.) was obtained.

EXAMPLE 2

Preparation of 1-[2,6-dichloro-4-(4-trifluoromethylphenyl)phenyl]-3-cyano-4-trifluoromethylsulfonyl-5-aminopyrazole

A solution of 1-[2,6-dichloro-4-(4-trifluoromethylphenyl)phenyl]-3-cyano-4-trifluoromethylthio-5-aminopyrazole (0.5 g), *m*-chloroperbenzoic acid (2.2 g) and 1,2-dichloroethane was heated at reflux for 40 hours. After cooling to 20° C., the mixture was dissolved in diethyl ether and washed with sat. aqueous $NaHCO_3$ solution twice, then saturated aqueous $NaHSO_3$ solution twice, dried, filtered, and the filtrate concentrated and purified by chromatography. The desired product as a solid (0.13 g, mp 197–202° C.) was obtained.

EXAMPLE 3

Preparation of 1-[2,6-dichloro-4-(tributyltin)phenyl]-3-cyano-4-trifluoromethylthio-5-aminopyrazole

A mixture of 1-(2,6-dichloro-4-bromophenyl)-3-cyano-4-trifluoromethylthio-5-aminopyrazole (5 g), bis-(tributyltin) (6.7 g) tetrakis(tri-phenylphosphine)palladium(0) (1.3 g) and 1-methyl-2-pyrrolidinone was heated to reflux for 75 minutes, cooled to 20° C., poured to water and extracted with methyl *t*-butyl ether. After concentration to dryness, methylene chloride was added and stirred with saturated aqueous KF solution for 16 hours. More water was added and extracted with methylene chloride. The methylene chloride solutions were combined, dried, filtered and the filtrate concentrated and purified via chromatography. The desired product as a solid (1.9 g, mp 116–117° C.) was obtained.

EXAMPLE 4

Preparation of 1-[2,6-dichloro-4-(2,3,5,6-tetrafluoropyrid-4-yl)phenyl]-3-cyano-4-trifluoromethylthio-5-aminopyrazole

A mixture of 1-[2,6-dichloro-4-(tributyltin)phenyl]-3-cyano-4-trifluoromethylthio-5-aminopyrazole (0.5 g), 4-bromo-2,3,5,6-tetrafluoropyridine (0.1 mL), tetrakis(triphenylphosphine)palladium(0) (0.1 g) and THF was refluxed for 26 hours. After cooling to 20° C., methylene chloride was added and stirred with saturated aqueous KF solution for 16 hours. The mixture was filtered with diethyl ether and dried, filtered again and the filtrate concentrated and purified via chromatography. The desired product as a solid (87 mg, mp 100–104° C.) was obtained.

EXAMPLE 5

Preparation of 1-[2,6-dichloro-4-(4-trifluoromethylphenyl)phenyl]-3-cyano-5-aminopyrazole

A mixture of 1-(2,6-dichloro-4-bromophenyl)-3-cyano-5-aminopyrazole (45 g, prepared according to procedures

reported in U.S. Pat. No. 5,232,940), 4-trifluoromethylphenylboronic acid (45 g), 2M of Na_2CO_3 (75 mL), tris(dibenzylidene acetone)dipalladium (6.4 g), toluene and ethanol was heated at 130° C. for 25 hours. After cooling to 20° C., the mixture was poured into water and extracted with diethyl ether. The ether solutions were combined, dried, filtered and the filtrate concentrated and purified via chromatography. The desired product as a solid (30.6 g, mp 195–199° C.) was obtained.

EXAMPLE 6

Preparation of 1-[2,6-dichloro-4-(4-trifluoromethylphenyl)phenyl]-3-cyano-4-bromo-5-aminopyrazole

A mixture of 1-[2,6-dichloro-4-(4-trifluoromethylphenyl)phenyl]-3-cyano-5-aminopyrazole (1 g), *N*-bromosuccinimide (0.5 g) and acetonitrile was stirred at 20° C. for 90 minutes, then concentrated and the residue was mixed with diethyl ether, washed with saturated aqueous $NaHCO_3$ solution, dried, filtered and the filtrate concentrated and purified via chromatography. The desired product as a solid (1.04 g, mp 178–180° C.) was obtained.

EXAMPLE 7

Preparation of 4-dichlorofluoromethylsulfonyl-1-[2-fluoro-4-(4-trifluoromethylphenyl)phenyl]imidazole

Step A

A solution of 4-bromo-2-fluoroaniline (3.8 g) in 20 ml of triethyl orthoformate was stirred at room temperature. After stirring for 1 h, tetrahydrofuran (40 ml), aminoacetonitrile hydrochloride (2.8 g), and diisopropylamine (5.3 ml) were added and stirring continued for an additional hour. After 1 h, the mixture was poured into 150 ml of ice water and extracted with 150 ml of dichloromethane. The organic solution was washed twice with water and dried over magnesium sulfate. Evaporation of solvents afforded a light purple solid (3.1 g), which was washed with hexane and filtered.

The solid was dissolved in 40 ml of dichloromethane at reflux, cooled to room temperature, and treated with tetramethylguanidine (1.6 ml), added dropwise over twenty minutes. After stirring at room temperature overnight, the mixture was cooled in an ice/salt bath and dichlorofluoromethylsulfonyl chloride (1.5 ml) was added. After 1 h, the mixture was diluted with 100 ml of methylene chloride and washed with saturated sodium bicarbonate solution, then with water, then dried over magnesium sulfate. After filtration and evaporation, chromatography on silica gel afforded 5-amino-4-dichlorofluoromethylsulfonyl-1-(2-fluoro-4-bromophenyl)imidazole (1.285 g).

Step B

1 g of 5-amino-4-dichlorofluoromethylsulfonyl-1-(2-fluoro-4-bromophenyl)imidazole was dissolved in 15 ml of tetrahydrofuran, cooled to 0° C., and treated with *t*-butylnitrite. After 1 h, the mixture is evaporated. Silica gel chromatography afforded 4-dichlorofluoromethylsulfonyl-1-(2-fluoro-4-bromophenyl)imidazole (0.555 g).

Step C

50 mg of 4-dichlorofluoromethylsulfonyl-1-(2-fluoro-4-bromophenyl)imidazole and 37 mg of trifluoromethylphenylboronic acid in 3 ml of toluene were treated with 0.13 ml of 2M aqueous potassium carbonate and a catalytic amount (about 5 mg) of palladium tetrakis(triphenylphosphine). The mixture was heated at 90° C. for 12 h, then cooled to room temperature and evaporated. Silica gel chromatography

afforded 4-dichlorofluoromethylsulfenyl-1-[2-fluoro-4-(4-trifluoromethylphenyl)phenyl]imidazole (27.3 mg, mass spec m/e 438).


In a manner similar to that employed in examples 1 to 7, the following compounds in Tables 1-4 were also prepared. The last column of the table indicates the physical characteristic of the compound obtained from the mass spectrum

analysis. It is the m/e value from the mass spectrum of the molecular ion. The number of the compound is for identification only.

N/A=Not Applicable


AroCH is [4-OH 3-methoxy phenyl]-CH, so that AroCH=N— is the group [[4-OH3-methoxy phenyl] CH=N—] 4-OH-3-methoxyphenyl]-CH=N—

TABLE 1




CMP. NO	R12	R13	R15	R16	R22	R23	R24	R25	R26	R3	R4	R5	m/e
1	Cl	H	H	Cl	(CH=CH)2	H	H	H	H	CN	SCF3	NH2	478
2	Cl	H	H	Cl	H	H	H	H	H	CN	SCF3	NH2	428
3	Cl	H	H	Cl	Me	H	H	H	H	CN	SCF3	NH2	442
4	Cl	H	H	Cl	H	NH2	H	H	H	CN	SCF3	NH2	443
5	Cl	H	H	Cl	H	NO2	H	H	H	CN	SCF3	NH2	473
6	Cl	H	H	Cl	H	NHAc	H	H	H	CN	SCF3	NH2	485
7	Cl	H	H	Cl	H	CHO	H	H	H	CN	SCF3	NH2	456
8	Cl	H	H	Cl	H	H	OMe	H	H	CN	SCF3	NH2	458
9	Cl	H	H	Cl	H	H	SMe	H	H	CN	SCF3	NH2	474
10	Cl	H	H	Cl	H	H	F	H	H	CN	SCF3	NH2	446
11	Cl	H	H	Cl	H	F	H	H	H	CN	SCF3	NH2	446
12	Cl	H	H	Cl	H	Cl	F	H	H	CN	SCF3	NH2	480
13	Cl	H	H	Cl	H	H	Cl	H	H	CN	SCF3	NH2	462
14	Cl	H	H	Cl	H	H	Br	H	H	CN	SCF3	NH2	506
15	Cl	H	H	Cl	H	Cl	H	Cl	H	CN	SCF3	NH2	496
16	Cl	H	H	Cl	H	H	CF3	H	H	CN	SCF3	NH2	496
17	Cl	H	H	Cl	H	CF3	H	CF3	H	CN	SCF3	NH2	564
18	Cl	H	H	Cl	(CH=CH)2	H	H	H	H	CN	SMe	NH2	424
19	Cl	H	H	Cl	H	H	H	H	H	CN	SMe	NH2	374
20	Cl	H	H	Cl	Me	H	H	H	H	CN	SMe	NH2	388
21	Cl	H	H	Cl	H	NH2	H	H	H	CN	SMe	NH2	389
22	Cl	H	H	Cl	H	NO2	H	H	H	CN	SMe	NH2	419
23	Cl	H	H	Cl	H	NHAc	H	H	H	CN	SMe	NH2	431
24	Cl	H	H	Cl	H	H	CHO	H	H	CN	SMe	NH2	402
25	Cl	H	H	Cl	H	H	OMe	H	H	CN	SMe	NH2	404
26	Cl	H	H	Cl	H	F	H	H	H	CN	SMe	NH2	392
27	Cl	H	H	Cl	H	Cl	F	H	H	CN	SMe	NH2	426
28	Cl	H	H	Cl	Cl	H	Cl	H	H	CN	SMe	NH2	442
29	Cl	H	H	Cl	H	H	CF3	H	H	CN	SMe	NH2	442
30	Cl	H	H	Cl	H	CF3	H	CF3	H	CN	SMe	NH2	510
31	Cl	H	H	Cl	H	Br	H	H	H	CN	SMe	NH2	452
32	Cl	H	H	Cl	Me	H	H	H	H	CN	SCF3	Van-CH=N—	576
33	Cl	H	H	Cl	H	NO2	H	H	H	CN	SCF3	Van-CH=N—	607
34	Cl	H	H	Cl	H	NHAc	H	H	H	CN	SCF3	Van-CH=N—	619
35	Cl	H	H	Cl	H	H	CHO	H	H	CN	SCF3	Van-CH=N—	590
36	Cl	H	H	Cl	H	H	OMe	H	H	CN	SCF3	Van-CH=N—	592
37	Cl	H	H	Cl	H	Cl	F	H	H	CN	SCF3	Van-CH=N—	614
38	Cl	H	H	Cl	Cl	H	Cl	H	H	CN	SCF3	Van-CH=N—	630
39	Cl	H	H	Cl	H	H	CF3	H	H	CN	SCF3	Van-CH=N—	630
40	Cl	H	H	Cl	H	H	CF3	H	H	CN	S(O)2CCl2F	NH2	560
41	Cl	H	H	Cl	H	H	CF3	H	H	CN	S(O)2CCl2F	NH2	544
42	Cl	H	H	Cl	H	H	CF3	H	H	CN	SCCl2F	NH2	528
43	Cl	H	H	Cl	H	H	CF3	H	H	CN	SOCl2F	NH2	512
44	Cl	H	H	Cl	H	H	CF3	H	H	CN	S(O)2CF3	NH2	528
45	Cl	H	H	Cl	H	H	CF3	H	H	PhC(O)	SCF3	NH2	575
46	Cl	H	H	Cl	(CH=CH)2	H	H	H	H	CN	SCF3	H	463
[47]	[Cl]	[H]	[H]	[Cl]	[H]	[(CH=)2]	[H]	[H]	[H]	[CN]	[SCF3]	[H]	[463]
47	Cl	H	H	Cl	H	(CH=CH)2	H	H	H	CN	SCF3	H	463
48	Cl	H	H	Cl	H	H	H	H	H	CN	SCF3	H	413
49	Cl	H	H	Cl	H	NO2	H	H	H	CN	SCF3	H	458
50	Cl	H	H	Cl	H	H	OMe	H	H	CN	SCF3	H	443
51	Cl	H	H	Cl	H	H	SMe	H	H	CN	SCF3	H	459
52	Cl	H	H	Cl	H	H	F	H	H	CN	SCF3	H	431
53	Cl	H	H	Cl	H	F	H	H	H	CN	SCF3	H	431
54	Cl	H	H	Cl	H	Cl	F	H	H	CN	SCF3	H	465
55	Cl	H	H	Cl	H	H	Cl	H	H	CN	SCF3	H	447
56	Cl	H	H	Cl	H	H	Br	H	H	CN	SCF3	H	491
57	Cl	H	H	Cl	H	Cl	H	Cl	H	CN	SCF3	H	481

TABLE 1-continued




CMP. NO	R12	R13	R15	R16	R22	R23	R24	R25	R26	R3	R4	R5	m/e
58	Cl	H	H	Cl	Cl	H	C	H	H	CN	SCF3	H	481
59	Cl	H	H	Cl	H	H	CF3	H	H	CN	SCF3	H	481
60	Cl	H	H	Cl	H	CF3	H	H	H	CN	SCF3	H	481
61	Cl	H	H	Cl	H	CF3	H	CF3	H	CN	SCF3	H	549
62	Cl	H	H	Cl	H	Cl	H	H	H	CN	SCF3	H	447
63	Cl	H	H	Cl	H	Br	H	H	H	CN	SCF3	NH2	491
64	Cl	H	H	Cl	H	H	CF3	H	H	PhC(O)	S(O)2CCl2F	NH2	639
65	Cl	H	H	Cl	H	H	Br	H	H	CN	SCCl2F	NH2	538
66	Cl	H	H	Cl	Cl	H	CF3	H	Cl	CN	SCF3	NH2	564
67	Cl	H	H	Cl	H	H	OCF3	H	H	CN	SCF3	NH2	512
68	Cl	H	H	Cl	H	H	CF3	H	H	PhC(O)	S(O)2CClF2	NH2	623
69	Cl	H	H	Cl	H	H	CF3	H	H	CN	H	NH2	396
70	Cl	H	H	Cl	H	H	CF3	H	H	CN	Br	NH2	474
71	Cl	H	H	Cl	H	H	CF3	H	H	CN	SEt	NH2	456
72	Cl	H	H	Cl	H	H	CF3	H	H	CN	SMe	NH2	442
73	Cl	H	H	Cl	H	H	CF3	H	H	CN	SCN	NH2	453
74	Cl	H	H	Cl	H	H	CF3	H	H	CN	Cl	NH2	430
75	Cl	H	H	Cl	H	H	CF3	H	H	CN	SOMe	NH2	458
76	Cl	H	H	Cl	H	H	CF3	H	H	CN	S(O)Et	NH2	472
77	Cl	H	H	Cl	H	H	Br	H	H	CN	S(O)2CCl2F	NH2	570
78	Cl	H	H	Cl	H	H	CF3	H	H	CN	S(O)2CH3	NH2	474
79	Cl	H	H	Cl	H	H	CF3	H	H	CN	S(O)2Et	NH2	488
80	Cl	H	H	Cl	H	H	CF3	H	H	CONH2	C(OH)(CF3)2	NH2	580
81	Cl	H	H	Cl	H	H	CF3	H	H	CN	C(OH)(CF3)2	NH2	562
82	Cl	H	H	Cl	H	H	CF3	H	H	CN	H	NHAc	438
83	Cl	H	H	Cl	H	H	CF3	H	H	Ac	SMe	NH2	459
84	Cl	H	H	Cl	H	H	CF3	H	H	CN	CF(CF3)2	NH2	564
85	Cl	H	H	Cl	H	H	CF3	H	H	CN	SCCl2F	H	513
86	Cl	H	H	Cl	H	H	CF3	H	H	CN	S(O)2CCl2F	H	545
87	Cl	H	H	Cl	H	Cl	Cl	H	H	CN	H	NH2	396
88	Cl	H	H	Cl	H	H	OCF3	H	H	CN	H	NH2	412
[89]	[Cl]	[H]	[H]	[Cl]	[H]	[H]	[H]	[H]	[H]	[CN]	[H]	[NH2]	[346]
89	Cl	H	H	Cl	H	H	H	H	H	CN	H	NH2	346
90	Cl	H	H	Cl	H	H	CO2Me	H	H	CN	H	NH2	386
91	Cl	H	H	Cl	H	Cl	Cl	H	H	CN	SCF3	NH2	496
92	Cl	H	H	Cl	F	H	H	H	H	CN	SCF3	NH2	446
93	Cl	H	H	Cl	H	H	CO2Me	H	H	CN	SCF3	NH2	486
94	Cl	H	H	Cl	H	H	CF3	H	H	CN	SCCl2F	Br	591
95	Cl	H	H	Cl	H	H	CF3	H	H	Ac	S(O)2CH3	NH2	491
96	Cl	H	H	Cl	Cl	H	H	H	Cl	CN	SCF3	NH2	496
97	Cl	H	H	Cl	Cl	H	Cl	H	H	CN	SCF3	NH2	496
98	Cl	H	H	Cl	Cl	H	CF3	H	Cl	CN	SCF3	NH2	564
99	Cl	H	H	Cl	CF3	H	H	H	H	CN	SCF3	NH2	496
100	Cl	H	H	Cl	NO2	H	CF3	H	H	CN	SCF3	NH2	541
101	Cl	H	H	Cl	Me	H	F	H	H	CN	SCF3	NH2	460
102	Cl	H	H	Cl	H	F	H	F	H	CN	SCF3	NH2	464
103	Cl	H	H	Cl	F	H	H	H	F	CN	SCF3	NH2	464
104	Cl	H	H	Cl	H	OCF3	H	H	H	CN	SCF3	NH2	512
[105]	[Cl]	[H]	[H]	[Cl]	[H]	[H]	[H]	[Cl]	[H]	[CN]	[SCF3]	[NH2]	[480]
105	Cl	H	H	Cl	H	F	H	Cl	H	CN	SCF3	NH2	480
106	Cl	H	H	Cl	H	CN	H	H	H	CN	SCF3	NH2	453
107	Cl	H	H	Cl	H	OMe	H	H	H	CN	SCF3	NH2	458
108	Cl	H	H	Cl	H	H	CH2CN	H	H	CN	SCF3	NH2	467
109	Cl	H	H	Cl	H	H	Ms	H	H	CN	SCF3	NH2	506
110	Cl	H	H	Cl	H	H	SO2NH2	H	H	CN	SCF3	NH2	507
111	Cl	H	H	Cl	H	H	NMe2	H	H	CN	SCF3	NH2	471
112	Cl	H	H	Cl	OCH2O	H	H	H	H	CN	SCF3	NH2	472
113	Cl	H	H	Cl	H	H	CN	H	H	CN	SCF3	NH2	453
114	Cl	H	H	Cl	H	H	CN	H	H	CN	SCF3	NH2	453
115	Cl	H	H	Cl	F	H	F	H	H	CN	SCF3	NH2	464
116	Cl	H	H	Cl	H	H	CONH2	H	H	CN	SCF3	NH2	471
117	Cl	H	H	Cl	H	Cl	CN	H	H	CN	SCF3	NH2	487
118	Cl	H	H	Cl	H	CF3	H	H	H	CN	SCF3	NH2	496
119	Cl	H	H	Cl	F	F	F	F	F	CN	SCF3	NH2	518
120	Cl	H	H	Cl	H	H	OPh	H	H	CN	SCF3	NH2	520
121	Cl	H	H	Cl	Cl	H	CF3	H	H	CN	SCF3	NH2	530

TABLE 1-continued




CMP. NO	R12	R13	R15	R16	R22	R23	R24	R25	R26	R3	R4	R5	m/e
[122]	[Cl]	[H]	[H]	[Cl]	[H]	[OCF2O]	[H]	[H]	[CN]	[SCF3]	[NH2]	[508]	
122	Cl	H	H	Cl	H	OCF2O	H	H	CN	SCF3	NH2	508	
123	Cl	H	H	Cl	H	F	NH2	H	H	CN	SCF3	NH2	461
124	Cl	H	H	Cl	H	CF3	NO2	H	H	CN	SCF3	NH2	541
125	Cl	H	H	Cl	OH	H	H	H	H	CN	SCF3	NH2	444
126	Cl	H	H	Cl	H	H	NH2	H	H	CN	SCF3	NH2	443
127	Cl	H	H	Cl	H	H	OH	H	H	CN	SCF3	NH2	444
128	Cl	H	H	Cl	H	CF3	H	NH2	NH2	CN	SCF3	NH2	526
129	Cl	H	H	Cl	H	F	Me	H	H	CN	SCF3	NH2	460
130	Cl	H	H	Cl	CN	F	H	H	H	CN	SCF3	NH2	471
131	Cl	H	H	Cl	H	Me	F	H	H	CN	SCF3	NH2	460
132	Cl	H	H	Cl	Me	H	H	F	H	CN	SCF3	NH2	460
133	Cl	H	H	Cl	Ph	H	H	H	H	CN	SCF3	NH2	504
134	Cl	H	H	Cl	OH	OMe	H	CHO	H	CN	SCF3	NH2	502
135	Cl	H	H	Cl	H	Cl	Cl	H	H	CN	S(O)2CH3	NH2	474
136	Cl	H	H	Cl	Cl	H	H	H	Cl	CN	S(O)2CH3	NH2	474
137	Cl	H	H	Cl	Cl	H	Cl	H	H	CN	S(O)2CH3	NH2	474
138	Cl	H	H	Cl	Cl	H	CF3	H	Cl	CN	S(O)2CH3	NH2	542
139	Cl	H	H	Cl	H	H	OCF3	H	H	CN	S(O)2CH3	NH2	490
140	Cl	H	H	Cl	CF3	H	H	H	H	CN	S(O)2CH3	NH2	474
141	Cl	H	H	Cl	NO2	H	CF3	H	H	CN	S(O)2CH3	NH2	519
142	Cl	H	H	Cl	Me	H	F	H	H	CN	S(O)2CH3	NH2	438
143	Cl	H	H	Cl	H	F	H	F	H	CN	S(O)2CH3	NH2	442
144	Cl	H	H	Cl	F	H	H	H	F	CN	S(O)2CH3	NH2	442
145	Cl	H	H	Cl	F	H	H	H	H	CN	S(O)2CH3	NH2	424
146	Cl	H	H	Cl	H	H	CF3	H	H	CN	C=NOH	NH2	439
147	Cl	H	H	Cl	H	H	CF3	H	H	CN	NO2	NHAc	483
148	Cl	H	H	Cl	H	H	CF3	H	H	CN	NO2	NH2	441
149	Cl	H	H	Cl	F	H	F	H	H	CN	S(O)2CH3	NH2	442
150	Cl	H	H	Cl	H	OCF3	H	H	H	CN	S(O)2CH3	NH2	490
151	Cl	H	H	Cl	H	F	H	Cl	H	CN	S(O)2CH3	NH2	458
152	Cl	H	H	Cl	H	H	CONH2	H	H	CN	S(O)2CH3	NH2	449
153	Cl	H	H	Cl	H	CN	H	H	H	CN	S(O)2CH3	NH2	431
154	Cl	H	H	Cl	H	H	SMe	H	H	CN	S(O)2CH3	NH2	452
155	Cl	H	H	Cl	H	OMe	H	H	H	CN	S(O)2CH3	NH2	436
156	Cl	H	H	Cl	H	H	CH2CN	H	H	CN	S(O)2CH3	NH2	445
157	Cl	H	H	Cl	H	H	Ms	H	H	CN	S(O)2CH3	NH2	484
158	Cl	H	H	Cl	H	H	SO2NH2	H	H	CN	S(O)2CH3	NH2	485
159	Cl	H	H	Cl	H	H	NMe2	H	H	CN	S(O)2CH3	NH2	449
160	Cl	H	H	Cl	H	OCH2O	H	H	H	CN	S(O)2CH3	NH2	450
161	Cl	H	H	Cl	H	Cl	CN	H	H	CN	S(O)2CH3	NH2	465
162	Cl	H	H	Cl	H	H	CN	H	H	CN	S(O)2CH3	NH2	431
163	Cl	H	H	Cl	H	H	CO2Me	H	H	CN	S(O)2CH3	NH2	464
164	Cl	H	H	Cl	H	CF3	H	H	H	CN	S(O)2CH3	NH2	474
165	Cl	H	H	Cl	F	F	F	F	F	CN	S(O)2CH3	NH2	496
166	Cl	H	H	Cl	H	H	OPh	H	H	CN	S(O)2CH3	NH2	498
167	Cl	H	H	Cl	Cl	H	CF3	H	H	CN	S(O)2CH3	NH2	508
168	Cl	H	H	Cl	Cl	H	SF5	H	Cl	CN	S(O)2CH3	NH2	600
169	Cl	H	H	Cl	Cl	H	OCF3	H	Cl	CN	S(O)2CH3	NH2	558
[170]	[Cl]	[H]	[H]	[Cl]	[H]	[OCF2O]	[H]	[H]	[CN]	[S(O)2CH3]	[NH2]	[486]	
170	Cl	H	H	Cl	H	OCF2O	H	H	CN	S(O)2CH3	NH2	486	
171	Cl	H	H	Cl	H	CO2H	OH	H	H	CN	S(O)2CH3	NH2	466
172	Cl	H	H	Cl	H	F	OH	H	H	CN	S(O)2CH3	NH2	440
173	Cl	H	H	Cl	H	F	NH2	H	H	CN	S(O)2CH3	NH2	439
[174]	[Cl]	[H]	[H]	[Cl]	[H]	[CF3]	[NH2]	[H]	[H]	[CN]	[S(O)2CH3]	[NH2]	[519]
174	Cl	H	H	Cl	H	CF3	NO2	H	H	CN	S(O)2CH3	NH2	519

TABLE 2




CMP. NO	R12	R13	R15	R16	R2	R22	R23	R24	R25	R26	R3	R4	R5	m/e
[175]	[Cl]	[H]	[H]	[Cl]	[H]	[(CH=CH)2]	[H]	[H]	[H]	[H]	[SCFCl2]	[CN]	[Cl]	[528]
175	Cl	H	H	Cl	H	(CH=CH)2	H	H	H	H	SCFCl2	CN	Cl	528
[176]	[Cl]	[H]	[H]	[Cl]	[Cl]	[H]	[(CH=CH)2]	[H]	[H]	[H]	[CN]	[SCFCl2]	[H]	[528]
176	Cl	H	H	Cl	Cl	H	(CH=CH)2	H	H	H	CN	SCFCl2	H	528
177	Cl	H	H	Cl	Cl	H	H	H	H	H	CN	SCFCl2	H	478
178	Cl	H	H	Cl	H	H	H	Me	H	H	SCFCl2	CN	Cl	492
179	Cl	H	H	Cl	H	Me	H	H	H	H	SCFCl2	CN	Cl	492
180	Cl	H	H	Cl	Cl	H	NH2	H	H	H	CN	SCFCl2	H	493
181	Cl	H	H	Cl	H	H	NO2	H	H	H	SCFCl2	CN	Cl	523
182	Cl	H	H	Cl	Cl	H	NHAc	H	H	H	CN	SCFCl2	H	535
183	Cl	H	H	Cl	H	CHO	H	H	H	H	SCFCl2	CN	C	506
184	Cl	H	H	Cl	H	H	CHO	H	H	H	SCFCl2	CN	Cl	506
185	Cl	H	H	Cl	Cl	H	H	CHO	H	H	CN	SCFCl2	H	506
186	Cl	H	H	Cl	H	OMe	H	H	H	H	SCFCl2	CN	Cl	508
187	Cl	H	H	Cl	H	H	OMe	H	H	H	SCFCl2	CN	Cl	508
188	Cl	H	H	Cl	Cl	H	H	OMe	H	H	CN	SCFCl2	H	508
189	Cl	H	H	Cl	Cl	H	H	SMe	H	H	CN	SCFCl2	H	524
190	Cl	H	H	Cl	H	H	H	F	H	H	SCFCl2	CN	Cl	496
191	Cl	H	H	Cl	Cl	H	F	H	H	H	CN	SCFCl2	H	496
192	Cl	H	H	Cl	Cl	H	Cl	F	H	H	CN	SCFCl2	H	530
193	Cl	H	H	Cl	H	H	H	Cl	H	H	SCFCl2	CN	Cl	512
194	Cl	H	H	Cl	H	H	H	Br	H	H	SCFCl2	CN	Cl	556
195	Cl	H	H	Cl	Cl	H	Cl	H	Cl	H	CN	SCFCl2	H	546
196	Cl	H	H	Cl	H	Cl	H	Cl	H	H	SCFCl2	CN	Cl	546
197	Cl	H	H	Cl	Cl	H	H	CF3	H	H	CN	SCFCl2	H	546
198	Cl	H	H	Cl	Cl	H	CF3	H	H	H	CN	SCFCl2	H	546
199	Cl	H	H	Cl	H	H	CF3	H	CF3	H	SCFCl2	CN	Cl	614
200	Cl	H	H	Cl	Cl	H	Cl	H	H	H	CN	SCFCl2	H	512
201	Cl	H	H	Cl	Cl	H	Br	H	H	H	CN	SCFCl2	H	556
202	Cl	H	H	Cl	H	H	H	CF3	H	H	S(O)CFC12	CN	Cl	562
203	Cl	H	H	Cl	H	H	H	CF3	H	H	S(O)CFC12	CN	Cl	578
204	H	H	H	H	H	H	H	CF3	H	H	CHO	H	H	315
205	H	H	H	Me	H	H	H	CF3	H	H	CHO	H	H	329
206	Me	H	H	Me	H	H	H	CF3	H	H	CHO	H	H	343
207	Cl	H	H	H	H	H	H	CF3	H	H	CHO	H	H	349
208	H	Cl	H	H	H	H	H	CF3	H	H	CHO	H	H	349
209	Cl	H	H	Me	H	H	H	CF3	H	H	CHO	H	H	353
210	Cl	H	H	Cl	H	H	H	CF3	H	H	CHO	H	H	383
211	F	H	H	H	H	H	H	CF3	H	H	CHO	H	H	333
212	F	H	H	F	H	H	H	CF3	H	H	CHO	H	H	351
213	H	H	H	CF3	H	H	H	CF3	H	H	CHO	H	H	383
214	H	H	CF3	H	H	H	H	CF3	H	H	CHO	H	H	383
215	F	F	F	F	H	H	H	CF3	H	H	CHO	H	H	387
216	Me	H	H	NO2	H	H	H	CF3	H	H	CHO	H	H	374
217	H	H	(CH=CH)2	H	H	H	H	CF3	H	H	CHO	H	H	365

TABLE 3




CMP. NO	R12	R13	R15	R16	R22	R23	R24	R25	R26	R3	R5	m/3
218	Cl	H	H	Cl	H	H	CF3	H	H	t-Bu	NH2	428
219	Cl	H	H	Cl	H	H	CF3	H	H	t-Bu	H	413
220	Cl	H	H	Cl	H	H	CF3	H	H	t-Bu	Br	491
221	Cl	H	H	Cl	H	H	CF3	H	H	CF3	NH2	440
222	Cl	H	H	Cl	H	NO2	H	H	H	CF3	NH2	417
223	Cl	H	H	Cl	H	F	H	H	H	CF3	NH2	390

TABLE 3-continued




CMP. NO	R12	R13	R15	R16	R22	R23	R24	R25	R26	R3	R5	m/3
224	Cl	H	H	Cl	H	H	CF3	H	H	CF3	Br	503
225	Cl	H	H	Cl	H	H	CF3	H	H	CF3	H	425
226	Cl	H	H	Cl	H	(CH=CH)2	H	H	H	CF3	NH2	422
227	Cl	H	H	Cl	(CH=CH)2	H	H	H	H	CF3	NH2	422
228	Cl	H	H	Cl	Me	H	H	H	H	CF3	NH2	386
229	Cl	H	H	Cl	H	OMe	H	H	H	CF3	NH2	402
230	Cl	H	H	Cl	H	H	OMe	H	H	CF3	NH2	402
231	Cl	H	H	Cl	H	H	SMe	H	H	CF3	NH2	418
232	Cl	H	H	Cl	H	H	F	H	H	CF3	NH2	390
233	Cl	H	H	Cl	H	H	Cl	H	H	CF3	NH2	406
234	Cl	H	H	Cl	H	H	Br	H	H	CF3	NH2	450
235	Cl	H	H	Cl	H	Cl	H	Cl	H	CF3	NH2	440
236	Cl	H	H	Cl	Cl	H	Cl	H	H	CF3	NH2	440
237	Cl	H	H	Cl	H	CF3	H	H	H	CF3	NH2	440
238	Cl	H	H	Cl	H	Cl	H	H	H	CF3	NH2	406
239	Cl	H	H	Cl	H	Br	H	H	H	CF3	NH2	450
240	Cl	H	H	Cl	(CH=CH)2	H	H	H	H	CF3	H	407
241	Cl	H	H	Cl	H	H	H	H	H	CF3	H	357
242	Cl	H	H	Cl	H	H	Me	H	H	CF3	H	371
243	Cl	H	H	Cl	H	NO2	H	H	H	CF3	H	402
244	Cl	H	H	Cl	H	CHO	H	H	H	CF3	H	385
245	Cl	H	H	Cl	H	OMe	H	H	H	CF3	H	387
246	Cl	H	H	Cl	H	H	F	H	H	CF3	H	375
247	Cl	H	H	Cl	H	H	F	H	H	CF3	H	375
248	Cl	H	H	Cl	H	F	H	H	H	CF3	H	375
249	Cl	H	H	Cl	H	Cl	F	H	H	CF3	H	409
250	Cl	H	H	Cl	H	(CH=CH)2	H	H	H	CF3	H	407
251	Cl	H	H	Cl	H	NHAc	H	H	H	CF3	H	414
252	Cl	H	H	Cl	CHO	H	H	H	H	CF3	H	385
253	Cl	H	H	Cl	H	H	CHO	H	H	CF3	H	385
254	Cl	H	H	Cl	OMe	H	H	H	H	CF3	H	387
255	Cl	H	H	Cl	H	H	CF3	H	H	SMe	H	403
256	Cl	H	H	Cl	H	H	CF3	H	H	SMe	NH2	418
257	Cl	H	H	Cl	H	H	CF3	H	H	SOMe	NH2	434
258	Cl	H	H	Cl	H	H	CF3	H	H	SOMe	H	419
259	Cl	H	H	Cl	H	H	CF3	H	H	SOMe	Br	497
260	Cl	H	H	Cl	H	H	CF3	H	H	SCCl2F	H	489
261	Cl	H	H	Cl	H	H	CF3	H	H	S(O)CCl2F	H	505
263	Cl	H	H	Cl	H	H	CF3	H	H	S(O)2CCl2F	H	521

TABLE 4



CMP. NO	R12	R13	R15	R16	R21	R23	R24	R25	R26	R4	R5	m/e
264	Cl	H	H	Cl	H	H	CF3	H	H	SCCl2F	H	488
265	Cl	H	H	Cl	H	H	CF3	H	H	SCCl2F	Br	566
266	Cl	H	H	Cl	H	H	CF3	H	H	SCCl2F	NH2	503
267	Cl	H	H	Cl	H	H	F3	H	H	SO2CCl2F	Br	537
268	Cl	H	H	Cl	H	H	CF3	H	H	SOCCl2F	H	504
269	Cl	H	H	Cl	H	H	CF3	H	H	SO2CCl2F	H	520
270	H	H	H	F	H	H	CF3	H	H	SOCCl2F	H	454
271	Cl	H	H	Cl	Cl	H	CF3	H	Cl	SOCCl2F	H	572
272	Cl	H	H	Cl	H	H	OCF3	H	H	SOCCl2F	H	520
273	Cl	H	H	Cl	H	H	OCF2O	H	H	SOCCl2F	H	516
274	Cl	H	H	Cl	H	H	H	H	H	SOCCl2F	H	436
275	Cl	H	H	Cl	Me	H	Me	Me	H	SOCCl2F	H	478

TABLE 4-continued



CMP. NO	R12	R13	R15	R16	R21	R23	R24	R25	R26	R4	R5	m/e
276	Cl	H	H	Cl	Me	H	H	H	Me	SOCCl2F	H	464
277	Cl	H	H	Cl	H	H	<i>t</i> -Bu	H	H	SOCCl2F	H	492
278	Cl	H	H	Cl	H	CF3	H	CF3	H	SOCCl2F	H	572
279	Cl	H	H	Cl	H	CH2CN	H	H	H	SOCCl2F	H	475
280	Cl	H	H	Cl	H	OCH2O	H	H	H	SOCCl2F	H	480
281	Cl	H	H	Cl	H	F	H	F	H	SOCCl2F	H	472
282	Cl	H	H	Cl	F	F	NH2	F	F	SOCCl2F	H	523
283	Cl	H	H	Cl	NO2	H	CF3	H	H	SOCCl2F	H	549
284	Cl	H	H	Cl	H	H	CH2CN	H	H	SOCCl2F	H	475
285	Cl	H	H	Cl	H	H	<i>n</i> -Bu	H	H	SOCCl2F	H	492
286	Cl	H	H	Cl	Cl	H	Cl	H	H	SOCCl2F	H	504
287	Cl	H	H	Cl	H	NO2	NH2	Me	Me	SOCCl2F	H	524
288	Cl	H	H	Cl	Cl	H	CF3	H	H	SOCCl2F	H	538
289	Cl	H	H	Cl	Cl	H	Cl	H	Cl	SOCCl2F	H	538
290	Cl	H	H	Cl	F	F	CF3	F	F	SOCCl2F	H	576
291	Cl	H	H	Cl	F	F	CN	F	F	SOCCl2F	H	533
292	Cl	H	H	Cl	5-tetrazolyl	H	H	H	H	SOCCl2F	H	504
293	Cl	H	H	Cl	H	H	CN	H	H	SOCCl2F	H	461
294	Cl	H	H	Cl	Cl	H	SF5	H	Cl	SOCCl2F	H	630
295	H	H	H	F	H	H	CF3	H	H	SCCl2F	H	438
296	Cl	H	H	Cl	H	H	CF3	H	H	SCF3	H	456
297	Cl	H	H	Cl	H	H	CF3	H	H	SOCF3	H	472
298	Cl	H	H	Cl	H	H	CF3	H	H	SO2CF3	H	488

The present invention provides also a method for controlling pests at a locus comprising applying to said locus a pesticidally effective amount of a compound of formula (I), or a pesticidal composition comprising a pesticidally effective amount of a compound of formula (I) and a pesticidally acceptable carrier therefor. In a preferred embodiment, the invention provides a method for controlling insects at a locus comprising applying to said locus an insecticidally effective amount of a compound of formula (I), or an insecticidally effective amount of an insecticidal composition comprising an insecticidally effective amount of a compound of formula (I) and an agriculturally acceptable inert carrier therefor. Preferably, the locus to which the pesticidally (especially insecticidally) effective amount is applied is a crop-growing area, that is, an area in which a crop is growing or in which a crop has been planted, or an area in which a crop will be planted/grown.

The compositions which can be used in the invention for the pesticidal/insecticidal treatment of the invention can comprise from about 0.001 to 95% of the compound of formula (I).

The diluted liquid formulations, as applied to the locus to be treated or crop, generally comprise from about 0.001 to about 3% of active ingredient of formula (I), preferably from about 0.1 to about 0.5%.

The solid formulations as applied to the locus or crop generally comprise from about 0.1 to about 8% of active ingredient of formula (I), preferably from about 0.5 to about 1.5%.

The concentrated compositions are the compositions which are commercialized or transported or stored. For application to plants, they are normally diluted in water and applied in such diluted form. The diluted forms are part of the invention as well as the concentrated forms.

The concentrated formulations generally comprise from about 5 to about 95% of active ingredient of formula (I), preferably from about 10 to about 50%.

The insecticidal compositions of the invention can be applied once, or more than once, throughout the whole insect season. Insecticidal compositions according to the invention are usually applied to the locus to be treated or crop area at a rate of from about 0.01 to about 2 kg/ha of active ingredient, preferably from about 0.1 to about 1 kg/ha.

The concentrated insecticidal compositions according to the invention can be in the form of a solid, e.g., dusts or granules or wettable powders, or, preferably, in the form of a liquid, such as an emulsifiable concentrate or a true solution.

The compositions according to the instant invention generally comprise from about 0.5 to about 95% of active ingredient of formula (I). The remainder of the composition up to 100% comprises a carrier as well as various additives such as those hereafter indicated.

By "carrier", there is meant herein an organic or inorganic material, which can be natural or synthetic, and which is associated with the active ingredient and which facilitates its application to the locus to be treated or crop. This carrier is thus generally inert and should be agriculturally acceptable, especially on the contemplated or treated locus or crop. The carrier can be solid (clay, silicates, silica, resins, wax, fertilizers, etc.) or liquid (water, alcohols, ketones, oil solvents, saturated or unsaturated hydrocarbons, chlorinated hydrocarbons, liquified petroleum gas, etc.).

Among the many additives, the compositions of the invention can comprise surfactants as well as other ingredients such as dispersants, stickers, antifoam agents, anti-freezing agents, dyestuffs, thickeners, adhesives, protective colloids, penetrating agents, stabilizing agents, sequestering agents, antiflocculating agents, corrosion inhibitors, pigments and polymers.

More generally, the compositions of the invention can comprise all kinds of solid or liquid additives which are known in the art of insecticides and insecticidal treatments.

The surfactants can be of the emulsifying or wetting type, ionic or non-ionic. Possible surfactants are salts of polyacrylic or lignosulfonic acids; salts of phenolsulfonic or naphthalenesulfonic acids; polycondensates of ethylene oxide with fatty alcohols or fatty acids or fatty amines or substituted phenols (particularly alkylphenols or arylphenols); ester-salts of sulfosuccinic acids; taurine derivatives, such as alkyl taurates; phosphoric esters; or esters of alcohols or polyoxyethylated phenols. When the spraying vehicle is water, the use of at least one surfactant is generally required because the active ingredients are not water-soluble.

The method of application of the compositions of the invention is generally the spraying of a mixture which has been previously made, by dilution of more concentrated formulations according to the invention.

Solid compositions can be powders for dusting or for dispersion (wherein the content of active ingredient can be up to 100%) and granules, especially extruded or compacted granules, or granules which have been made by impregnation of a powder (the content of active ingredient in such powders being between about 1 and about 80%).

Liquid compositions of compositions which have to be liquid when applied include solutions, water-soluble concentrates, emulsifiable concentrates, emulsions, wettable powders or pastes or water-dispersible granules.

Emulsifiable concentrates generally comprise from about 10 to about 80% of active ingredient; the emulsions when applied generally comprise from about 0.01 to about 20% of active ingredient.

For example, the emulsifiable concentrates can comprise the solvent and, to the extent needed, from about 2 to about 20% of suitable additives such as stabilizers, surfactants, penetrating agents, corrosion inhibitors or other additives already recited.

These concentrates are usually diluted in tank water so as to obtain the dilution appropriate for spraying.

The concentrated suspensions can also be applied by spraying and have to be fluid without allowing any solid to separate and fall to the bottom. Generally they comprise from about 1 to about 75% of active ingredient (preferably from about 2 to about 50%), from about 0.5 to about 15% of surfactant, from about 0.1 to about 10% of thickener, from 0 to about 10% of other suitable additives as already indicated, the remainder being water or an organic liquid wherein the active ingredient is insoluble or has a low solubility.

The wettable powders generally comprise the active ingredient (from about 1 to about 95%, preferably from about 2 to about 80%), the solid carrier, a wetting agent (from 0 to about 5%), a dispersing agent (from about 3 to about 10%) and, to the extent needed, from 0 to about 10% of other additives such as stabilizers and others as already listed.

In order to obtain these wettable powders or dusting powders, it is appropriate to intimately mix the active ingredients and the additives, as by grinding in a mill or similar device.

Dispersible granules are generally made by agglomeration of a powder, followed by an appropriate granulation process.

The emulsions herein described can be of the oil-in-water or water-in-oil types. Fluidity of the emulsions can range from low viscosities up to high viscosities approaching those of gels.

Among these many compositions or formulations, one skilled in the art can choose the one most appropriate, according to the specific conditions of the treatment problem.

The compounds and compositions of the invention can also be used in admixtures with another pesticide, e.g., an insecticide, acaricide or herbicide.

The compounds of the invention may also be used in controlling pests found in non-agricultural domains.

In the field of veterinary medicine or livestock husbandry or in the maintenance of public health against arthropods, helminths or protozoa which are parasitic internally or externally upon vertebrates, particularly warm-blooded vertebrates, for example man or domestic animals, e.g. cattle, sheep, goats, equines, swine, poultry, dogs or cats, for example Acarina, including ticks (e.g. *Ixodes* spp., *Boophilus* spp. e.g. *Boophilus microplus*, *Amblyomma* spp., *Hyalomma* spp., *Rhipicephalus* spp. e.g. *Rhipicephalus appendiculatus*, *Haemaphysalis* spp., *Dermacentor* spp., *Ornithodoros* spp. (e.g. *Ornithodoros moubata*) and mites (e.g. *Damalinia* spp., *Dermahyssus gallinae*, *Sarcoptes* spp. e.g. *Sarcoptes scabiei*, *Psoroptes* spp., *Chorioptes* spp., *Demodex* spp., *Eutrombicula* spp.); Diptera (e.g. *Aedes* spp., *Anopheles* spp., *Dermatobia* spp., *Haematobia* spp., *Musca* spp., *Hippoboscidae* spp., *Hypoderma* spp., *Gasterophilus* spp., *Simulium* spp.); Stomoxys spp., Hemiptera (e.g. *Triatoma* spp.); Phthirapter (e.g. *Damalinia* spp., *Linognathus* spp.); Siphonaptera (e.g. *Ctenocephalides* spp.); Dicotyoptera (e.g. *Periplaneta* spp., *Blattella* spp.); Hymenoptera (e.g. *Monomorium pharaonis*); for example against infections of the gastro-intestinal tract caused by parasitic nematode worms, for example members of the family *Trichostrongylidae*, *Nippostrongylus brasiliensis*, *Trichinella spiralis*, *Haemonchus contortus*, *Trichostrongylus colubriformis*, *Nematodirus batus*, *Ostertagia circumcincta*, *Trichostrongylus axei*, *Cooperia* spp. and *Hymenolepis nana*; in the control and treatment of protozoal diseases caused by, for example, *Eimeria* spp. e.g. *Eimeria tenella*, *Eimeria acervulina*, *Eimeria brunetti*, *Eimeria maxima* and *Eimeria necatrix*, *Trypanosoms cruzi*, *Leishmania* spp., *Plasmodium* spp., *Babesia* spp., *Trichomonadidae* spp., *Histomonas* spp., *Giardia* spp., *Toxoplasma* spp., *Entamoeba histolytica* and *Theileria* spp.

Furthermore, the compounds of the invention may be useful for coccidiosis, a disease caused by infections from protozoan parasites of the genus *Eimeria*.

Compositions

Solid or liquid compositions for application topically to animals, timber, stored products or household goods usually contain from about 0.00005% to about 90%, more particularly from about 0.001% to about 10%, by weight of one or more compounds of general formula (I). For administration to animals orally or parenterally, including percutaneously solid or liquid compositions, these normally contain from about 0.1% to about 90% by weight of one or more compounds of general formula (I). Medicated feedstuffs normally contain from about 0.001% to about 3% by weight of one or more compounds of general formula (I). Concentrates or supplements for mixing with feedstuffs normally contain from about 5% to about 90%, preferably from about 5% to about 50%, by weight of one or more compounds of general formula (I). Mineral salt licks normally contain from about 0.1% to about 10% by weight of one or more compounds of general formula (I).

Dusts or liquid compositions for application to livestock, persons, goods, premises or outdoor areas may contain from about 0.0001% to about 15%, more especially from about 0.005% to about 2.0%, by weight, of one or more compounds of general formula (I). Suitable concentrations in treated waters are between about 0.0001 ppm and about 20 ppm, more particularly about 0.001 ppm to about 5.0 ppm of

one or more compounds of general formula (I) and may be used therapeutically in fish farming with appropriate exposure times. Edible baits may contain from about 0.01% to about 5%, preferably from about 0.01% to about 1.0%, by weight, of one or more compounds of general formula (I).

When administered to vertebrates parenterally, orally or by percutaneous or other means, the dosage of compounds of general formula (I) will depend upon the species, age, or health of the vertebrate and upon the nature and degree of its actual or potential infestation by arthropod, helminth or protozoan pests. A single dose of about 0.1 to about 100 mg, preferably about 2.0 to about 20.0 mg, per kg body weight of the animal or doses of about 0.01 to about 20.0 mg, preferably about 0.1 to about 5.0 mg, per kg body weight of the animal per day, for sustained medication, are generally suitable by oral or parenteral administration. By use of sustained release formulations or devices, the daily doses required over a period of months may be combined and administered to animals on a single occasion.

The following methods were used to apply the compounds of the invention and to observe the results obtained therewith: a foliar/contact spray on sucking (aphids) or chewing (Lepidoptera) insects.

The species tested were as follows:

GENUS, SPECIES	COMMON NAME
<i>Aphis gossypii</i>	cotton leaf aphid
<i>Musca domestica</i>	housefly
<i>Diabrotica virgifera</i>	Western cornrootworm
<i>Periplaneta americana</i>	American Cockroaches
<i>Spodoptera eridania</i>	Southern armyworm

The Housefly Bait/Contact Test

About 25 four to six-day-old adult houseflies were anesthetized and placed in a cage with a sugar water bait solution containing the compound. The concentration of the selected compound of formula (I) in the bait solution was 50 ppm. After 24 hours, flies which showed no movement on stimulation were considered dead.

Foliar application (Contact Test) with Aphids

Aphid-infested cotton plants were placed on a revolving turntable, and sprayed to runoff with a 100 ppm formulation of the selected compound of formula (I). The treated, *A. gossypii*-infested plants were held for three days after treatment, after which the dead aphids were counted.

Application on Western cornrootworm

Jars are filled with 60 g dry soil homogeneously mixed with 5 ml aqueous solution containing the compound to be tested. After drying, four germinated corn seedlings are placed in the bottom of each jar and the soil is wetted. Then ten newly born larvae are placed in each jar. The jars are stored for six days at 27° C. under 70% of relative humidity. The number of surviving larvae are then counted.

Application by contact to Cockroaches

The inner wall of a 100 ml jar is covered by shaking 2 ml of pesticidal formulation. Furthermore, a pellet of dog chow is placed into a jar and the pesticidal formulation is absorbed on the pellet, using excess liquid. The jar is then left open up to evaporation of liquid. Cockroach nymphs are placed in the jar and mortality is assessed after five days.

Application to Southern armyworm

Leaves of soybeans were sprayed to run-off with a spray of various concentrations. Then the leaves were allowed to dry, excised, and placed into a container with five larvae. The containers are stored for five days at 25° C. under 50% relative humidity. Mortality ratings are then counted.

Application to *Caenorhabditis elegans* (nematode worm)

This test is significant for animal health applications of active ingredient.


Test formulation are prepared by mixing 2 microliters of dimethylsulfoxide with 100 microliters of a solution of *Escherichia coli* (which is food for *C. elegans*). 98 microliters of this mixture are then mixed with an inoculum solution which consists of mixed life stages of *C. elegans* (about 50 worms). This new mixture is stored 7 days at 20° C. Mortality is then visually assessed as well as the behavior of the worm. For the mortality, the rating is either 1 (large increase in number of the population, similar to the control which is trebling in 7 days) or 3 (slight increase of the population, but significantly less increase than the control) or 5 (little to no increase in the population). For the behavior effect, the rating is either 1 (normal motion) or 3 (motion is slower than the control) or 5 (little to no movement).

All the hereinabove prepared compounds show a positive activity in one or more of these tests (rather good activity on *C. elegans*).

While the invention has been described in terms of various preferred embodiments, the person skilled in the art will appreciate that various modifications, substitutions, omissions and changes can be made without departing from the spirit thereof. Accordingly, it is intended that the scope of the present invention be limited solely by the scope of the following claims, including equivalents thereof.

What is claimed is:

1. A compound of formula (I):



(I)

wherein:

X is N;

Y is C—R₃;

W is C—R₄;

R₃ is CN;

R₄ is —S(O)_nR₈;

R₅ is —NR₉R₁₀;

R₈ is alkyl, haloalkyl, alkenyl, or alkynyl, or R₈ is a cycloalkyl ring having 3 to 5 carbon atoms;

R₉ and R₁₀, independently of each other, are H, alkyl, haloalkyl, alkylcarbonyl, haloalkylcarbonyl, R₈S(O)_n, formyl, alkenyl, alkynyl, alkoxy carbonyl, alkylthiocarbonyl, or aroyl, the alkyl portions of R₉ and R₁₀ being optionally substituted by R₇;

R₇ is cyano, nitro, alkoxy, haloalkoxy, R₈S(O)_n, —C(O) alkyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, —CO₂H, halogen, hydroxy, aminosulfonyl, alkylaminosulfonyl or dialkylaminosulfonyl;

Z is C—R₁₆;

n is zero, one or two;

R₁₂, R₁₃, R₁₅ and R₁₆, independently of one another, are hydrogen, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, alkylsulfenyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfenyl, haloalkylsulfinyl, haloalkylsulfonyl, formyl, alkylcarbonyl, alkoxy carbonyl;

R₂₂, R₂₃, R₂₄, R₂₅, and R₂₆, independently of one another, are hydrogen, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, haloalkylcarbonyl, formyl, alkylcarbonyl, thioamide, amide, alkoxy carbonyl, SF₅, or R₈S(O)_{*n*};

or a pesticidally acceptable salt thereof.

2. A compound according to claim 1, wherein R₂₄ is halogen, haloalkyl or haloalkoxy.

3. A compound according to claim 1, wherein R₅ is —NH₂.

4. A compound according to claim 1, wherein R₈ is lower alkyl or haloalkyl.

5. A compound according to claim 1, having at least one feature selected from the group consisting of:

R₈ is methyl, ethyl, —CF₃, —CFCl₂, or —CF₂Cl;

R₁₂ and R₁₆, independently of each other, are F, Cl, Br or H;

R₁₃ and R₁₅ are H;

R₂₄ is —CF₃, —OCF₃, —CHF₂, —S(O)_{*n*}CF₃, —CFCl₂, —CF₂Cl, —OCF₂Cl, —OCFCl₂, Cl, Br or F; and

Z is CCl, CF, or CBr.

6. A compound according to [claim 2] claim 5, wherein R₅ is —NH₂.

7. A compound according to claim 1, wherein:

R₁₂ and R₁₆, independently of each other, are Cl or Br;

R₁₃ and R₁₅ are H;

R₂₄ is —CF₃, —OCF₃ or Br;

R₅ is amino;

R₉ and R₁₀, independently of each other, are H, alkyl or alkylcarbonyl; and

R₈ is methyl, ethyl, CF₃, CFCl₂ or CF₂Cl.

8. A compound according to [claim 2] claim 5, wherein:

R₁₂ and R₁₆, independently of each other, are Cl or Br;

R₂₄ is —CF₃, —OCF₃ or Br;

R₅ is amino; and

R₉ and R₁₀, independently of each other, are H, alkyl or alkylcarbonyl.

9. A compound according to claim 1, wherein the S(O)_{*n*}R₈ substituent of formula (I) is: methylthio, methylsulfinyl, methylsulfonyl, ethylsulfinyl, ethylsulfonyl, ethylthio, cyclopropylsulfinyl, cyclopropylthio, cyclopropylsulfonyl, isopropylsulfinyl, isopropylsulfonyl, isopropylthio, trifluoromethylthio, trifluoromethylsulfinyl, trifluoromethylsulfonyl, dichlorofluoromethylthio, dichlorofluoromethylsulfinyl, dichlorofluoromethylsulfonyl, chlorodifluoromethylthio, chlorodifluoromethylsulfonyl or chlorodifluoromethylsulfinyl.

10. A compound according to claim 5, wherein the S(O)_{*n*}R₈ substituent of formula (I) is: trifluoromethylthio, trifluoromethylsulfinyl, trifluoromethylsulfonyl, dichlorofluoromethylthio, dichlorofluoromethylsulfinyl, dichlorofluoromethylsulfonyl, chlorodifluoromethylthio, chlorodifluoromethylsulfonyl or chlorodifluoromethylsulfinyl.

11. A compound according to claim 7, wherein the S(O)_{*n*}R₈ substituent of formula (I) is: trifluoromethylthio,

trifluoromethylsulfinyl, trifluoromethylsulfonyl, dichlorofluoromethylthio, dichlorofluoromethylsulfinyl, dichlorofluoromethylsulfonyl, chlorodifluoromethylthio, chlorodifluoromethylsulfonyl or chlorodifluoromethylsulfinyl.

12. A compound according to claim 8, wherein the S(O)_{*n*}R₈ substituent of formula (I) is: trifluoromethylthio, trifluoromethylsulfinyl, trifluoromethylsulfonyl, dichlorofluoromethylthio, dichlorofluoromethylsulfinyl, dichlorofluoromethylsulfonyl, chlorodifluoromethylthio, chlorodifluoromethylsulfonyl or chlorodifluoromethylsulfinyl.


13. The compound of formula (I) according to claim 1, which is:

1-[2,6-dichloro-4-(4-trifluoromethylphenyl)phenyl]-3-cyano-4-trifluoromethylthio-5-aminopyrazole; or

1-[2,6-dichloro-4-(4-trifluoromethylphenyl)phenyl]-3-cyano-4-trifluoromethylsulfonyl-5-aminopyrazole.

14. A pesticidal composition comprising:

(a) a pesticidally effective amount of a compound of formula (I):



wherein:

X is N;

Y is C—R₃;

W is C—R₄;

R₃ is CN;

R₄ is —S(O)_{*n*}R₈;

R₅ is —NR₉R₁₀;

R₈ is alkyl, haloalkyl, alkenyl, or alkynyl, or R₈ is a cycloalkyl ring having 3 to 5 carbon atoms;

R₉ and R₁₀, independently of each other, are H, alkyl, haloalkyl, alkylcarbonyl, haloalkylcarbonyl, R₈S(O)_{*n*}, formyl, alkenyl, alkynyl, alkoxy carbonyl, alkylthiocarbonyl, or aroyl, the alkyl portions of R₉ and R₁₀ being optionally substituted by R₇;

R₇ is cyano, nitro, alkoxy, haloalkoxy, R₈S(O)_{*n*}, —C(O)alkyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, —CO₂H, halogen, hydroxy, aminosulfonyl, alkylaminosulfonyl or dialkylaminosulfonyl;

Z is C—R₁₆;

n is zero, one or two;

R₁₂, R₁₃, R₁₅ and R₁₆, independently of one another, are hydrogen, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, alkylsulfenyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfenyl, haloalkylsulfinyl, haloalkylsulfonyl, formyl, alkylcarbonyl or alkoxy carbonyl;

29


R_{22} , R_{23} , R_{24} , R_{25} , and R_{26} , independently of one another, are *hydrogen*, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, haloalkylcarbonyl, formyl, alkylcarbonyl, thioamide, amide, alkoxy carbonyl, SF_5 , or $R_8S(O)_n$;

or a pesticidally acceptable salt thereof.

(b) a pesticidally acceptable carrier therefor.

15. A composition according to claim 14, comprising from 0.001 to 95% of compound of formula (I).

16. A method for controlling pests at a locus comprising applying to said locus a pesticidally effective amount of a compound of formula (I):



wherein:

X is N;

Y is C—R₃;

W is C—R₄;

R₃ is [H, halogen, hydroxy, alkyl, cycloalkyl, haloalkyl, alkoxy, haloalkoxy,] CN;

R₄ is —S(O)_nR₈;

R₅ is —NR₉R₁₀;

R₈ is alkyl, haloalkyl, alkenyl, or alkynyl, or R₈ is a cycloalkyl ring having 3 to 5 carbon atoms;

R₉ and R₁₀, independently of each other, are H, alkyl, haloalkyl, alkylcarbonyl, haloalkylcarbonyl, $R_8S(O)_n$,

30

formyl, alkenyl, alkynyl, alkoxy carbonyl, alkylthiocarbonyl, or aroyl, the alkyl portions of R₉ and R₁₀ being optionally substituted by R₇;

R₇ is cyano, nitro, alkoxy, haloalkoxy, $R_8S(O)_n$, —C(O) alkyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, —CO₂H, halogen, hydroxy, aminosulfonyl, alkylaminosulfonyl or dialkylaminosulfonyl;

Z is C—R₁₆;

n is zero, one or two;

R₁₂, R₁₃, R₁₅ and R₁₆, independently of one another, are hydrogen, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, alkylsulfenyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfenyl, haloalkylsulfinyl, haloalkylsulfonyl, formyl, alkylcarbonyl or alkoxy carbonyl;

R₂₂, R₂₃, R₂₄, R₂₅, and R₂₆, independently of one another, are *hydrogen*, halogen, alkyl, haloalkyl, cyanoalkyl, cyano, nitro, amino, hydrazino, alkoxy, haloalkoxy, haloalkylcarbonyl, formyl, alkylcarbonyl, thioamide, amide, alkoxy carbonyl, SF_5 , or $R_8S(O)_n$;

or a pesticidally acceptable salt thereof.

17. A method according to claim 16, wherein said pests are insects and wherein said pesticidally effective amount is an insecticidally effective amount.

18. A method according to claim 17, comprising applying to said locus from about 0.01 to about 2 kg/ha of compound of formula (I).

19. A method according to claim 18, comprising applying to said locus from about 0.1 to about 1 kg/ha of compound of formula (I).

20. A method for controlling pests at a locus comprising applying to said locus a pesticidally effective amount of a composition as claimed in claim 14.

21. A method according to claim 20, wherein said pests are insects and wherein said pesticidally effective amount is an insecticidally effective amount.

* * * * *