



US00RE39220E

(19) **United States**
 (12) **Reissued Patent**
Gopal

(10) **Patent Number: US RE39,220 E**
 (45) **Date of Reissued Patent: *Aug. 1, 2006**

(54) **PEPTIDE-MEDIATED GENE TRANSFER** 5,482,858 A 1/1996 Huston et al.
 5,502,037 A 3/1996 Kondratyev
 (75) Inventor: **T. Venkat Gopal**, North Potomac, MD 5,503,833 A 4/1996 Redmond et al.
 (US) 5,514,546 A 5/1996 Kool et al.
 (73) Assignee: **Genetic Applications, LLC**, San 5,521,290 A 5/1996 Sivam et al.
 Diego, CA (US) 5,521,291 A 5/1996 Curiel et al.
 (*) Notice: This patent is subject to a terminal dis- 5,527,885 A 6/1996 Coughlin et al.
 claimer. 5,541,287 A 7/1996 Yau et al.
 (21) Appl. No.: **09/404,979** 5,547,932 A 8/1996 Curiel et al.
 5,574,142 A 11/1996 Meyer, Jr. et al.
 (22) Filed: **Sep. 22, 1999** 5,576,201 A 11/1996 Mason et al.
 5,578,287 A 11/1996 Theodore et al.
 5,583,020 A 12/1996 Sullivan
 5,585,468 A 12/1996 Coughlin et al.
 5,589,392 A * 12/1996 Short 435/320.1

(Continued)

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **5,670,347**
 Issued: **Sep. 23, 1997**
 Appl. No.: **08/240,514**
 Filed: **May 11, 1994**

(51) **Int. Cl.**
C12N 15/87 (2006.01)
C12N 15/63 (2006.01)
C12N 15/85 (2006.01)
C07K 14/00 (2006.01)

(52) **U.S. Cl.** **435/455**; 435/320.1; 435/467;
 530/300; 530/350

(58) **Field of Classification Search** 435/455,
 435/467, 468, 471, 320.1; 530/300, 350
 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,046,722 A 9/1977 Rowland
 4,671,958 A 6/1987 Rodwell et al.
 4,691,006 A 9/1987 Stevens
 4,847,240 A 7/1989 Ryser et al.
 4,867,973 A 9/1989 Goers et al.
 4,870,023 A 9/1989 Fraser et al.
 4,897,355 A 1/1990 Eppstein
 4,904,582 A 2/1990 Tullis et al.
 4,946,778 A 8/1990 Ladner et al.
 4,946,787 A 8/1990 Eppstein et al.
 4,950,599 A 8/1990 Bertling et al.
 5,024,939 A * 6/1991 Gorman 435/69.1
 5,047,227 A 9/1991 Rodwell et al.
 5,071,651 A 12/1991 Sabara et al.
 5,084,441 A 1/1992 Shaw et al.
 5,108,921 A 4/1992 Low et al.
 5,135,736 A 8/1992 Anderson et al.
 5,156,840 A 10/1992 Goers et al.
 5,166,320 A 11/1992 Wu et al.
 5,169,933 A 12/1992 Anderson et al.
 5,171,563 A 12/1992 Abrams et al.
 5,182,107 A 1/1993 Friden
 5,196,510 A 3/1993 Rodwell et al.
 5,229,490 A 7/1993 Tam et al.
 5,283,342 A 2/1994 Gustavson et al.
 5,298,422 A 3/1994 Schwartz et al.
 5,326,856 A 7/1994 Coughlin et al.
 5,354,844 A * 10/1994 Beug et al. 530/345
 5,420,105 A 5/1995 Gustavson et al.
 5,449,761 A 9/1995 Belinka, Jr. et al.

FOREIGN PATENT DOCUMENTS

EP 0 359 347 3/1990
 WO WO 91/08220 6/1991
 WO WO 91/16024 10/1991
 WO WO 92/13570 8/1992
 WO WO 92/15682 9/1992
 WO WO 93/07282 4/1993
 WO WO 93/07283 4/1993
 WO WO 93/18137 9/1993
 WO WO 93/18759 * 9/1993
 WO WO 93/19768 10/1993
 WO WO 93/20186 10/1993
 WO WO 94/04698 3/1994
 WO WO 94/23751 10/1994
 WO WO 95/02397 1/1995
 WO WO 95/09640 4/1995
 WO WO 95/11692 5/1995
 WO WO 95/14078 5/1995
 WO WO 95/31557 11/1995
 WO WO95/34647 12/1995
 WO WO 96/01841 1/1996
 WO WO 96/02662 2/1996
 WO WO 96/05218 2/1996
 WO WO 96/10038 4/1996

OTHER PUBLICATIONS

Moreland, R.B. et al. (1987) Amino Acid Sequences That Determine the Nuclear Localization of Yeast Histone 2B, Mol. Cell Biol. 7, 4048-4057.

(Continued)

Primary Examiner—Terry McKelvey
 (74) *Attorney, Agent, or Firm*—Patton Boggs LLP

(57) **ABSTRACT**

[A methodology that allows for highly efficient transfer and stable integration of DNA into both established eukaryotic cell lines and primary cells, including non-dividing cells such as human peripheral blood monocytes and macrophages, entails the use of a synthetic polypeptide comprised of a peptide domain which corresponds to a nuclear localization signal sequence and a DNA binding domain which is rich in basic amino acids, separated by a hinge region of neutral acid which prevents steric interference between the two domains.] *A synthetic polypeptide that allows for highly efficient transfer of DNA into eukaryotic cells, including, for example, non-dividing cells such as human peripheral blood monocytes and macrophages.*

14 Claims, No Drawings

US RE39,220 E

U.S. PATENT DOCUMENTS							
5,593,866	A	1/1997	Hancock et al.	5,908,623	A	6/1999	Baetge et al.
5,599,704	A	2/1997	Thompson et al.	5,914,265	A	6/1999	Roop et al.
5,607,691	A	3/1997	Hale et al.	5,925,564	A	7/1999	Schwartz et al.
5,608,060	A	3/1997	Axworthy et al.	5,945,100	A	8/1999	Fick
5,610,052	A	3/1997	Thompson et al.	5,955,365	A	9/1999	Szoka, Jr. et al.
5,616,490	A	4/1997	Sullivan et al.	5,955,647	A	9/1999	Fitchen et al.
5,622,854	A	4/1997	Draper	5,958,412	A	9/1999	Welt et al.
5,630,996	A	5/1997	Reno et al.	5,958,764	A	9/1999	Roop et al.
5,631,236	A	5/1997	Woo et al.	5,962,427	A	10/1999	Goldstein et al.
5,631,237	A	5/1997	Dzau et al.	5,965,371	A	10/1999	Marasco et al.
5,635,380	A	6/1997	Naftilan et al.	5,972,699	A	10/1999	Draper et al.
5,637,471	A	6/1997	Artavanis-Tsakonas et al.	5,972,704	A	10/1999	Draper et al.
5,637,481	A	6/1997	Ledbetter et al.	5,972,900	A	10/1999	Ferkol, Jr.
5,639,275	A	6/1997	Baetge et al.	5,977,084	A	11/1999	Szoka, Jr. et al.
5,639,655	A	6/1997	Thompson et al.	5,977,307	A	11/1999	Friden et al.
5,641,662	A	6/1997	Debs et al.	5,981,273	A	11/1999	Curiel et al.
5,648,248	A	7/1997	Zenke et al.	5,989,906	A	11/1999	Thompson et al.
5,652,122	A	7/1997	Frankel et al.	5,990,089	A	11/1999	Szoka, Jr. et al.
5,654,006	A	8/1997	Fernandez et al.	5,994,109	A	11/1999	Woo et al.
5,661,025	A	8/1997	Szoka, Jr. et al.	6,004,940	A	12/1999	Marasco et al.
5,668,255	A	9/1997	Murphy	6,008,336	A	12/1999	Hanson et al.
5,670,483	A	9/1997	Zhang et al.	6,010,908	A	1/2000	Gruenert et al.
5,670,617	A	9/1997	Frankel et al.	6,020,202	A	2/2000	Jessee
5,672,479	A	9/1997	Johnson et al.	6,022,735	A	2/2000	Curiel et al.
5,674,703	A	10/1997	Woo et al.	6,022,950	A	2/2000	Murphy et al.
5,674,704	A	10/1997	Goodwin et al.	6,022,966	A	2/2000	Gustavson et al.
5,674,835	A	10/1997	Androphy et al.	6,030,822	A	2/2000	Lechiner et al.
5,674,977	A	10/1997	Gariepy	6,033,884	A	3/2000	Woo et al.
5,674,980	A	10/1997	Frankel et al.	6,037,329	A	3/2000	Baird et al.
5,683,874	A	11/1997	Kool et al.	6,037,526	A	3/2000	Grimsley et al.
5,686,264	A	11/1997	Gaynor et al.	6,057,298	A	5/2000	Roop et al.
5,693,531	A	12/1997	Chiorini et al.	6,066,624	A	5/2000	Woo et al.
5,714,166	A	2/1998	Tomalia et al.	6,077,663	A	6/2000	Curiel et al.
5,717,058	A	2/1998	Matthews et al.	6,077,835	A	6/2000	Hanson et al.
5,747,641	A	5/1998	Frankel et al.	6,113,946	A	9/2000	Szoka, Jr. et al.
5,750,367	A	5/1998	Chan	6,117,657	A	9/2000	Usman et al.
5,750,390	A	5/1998	Thompson et al.	6,127,114	A	10/2000	Haseloff et al.
5,756,264	A	5/1998	Schwartz et al.	6,130,366	A	10/2000	Herrera-Estrella et al.
5,756,353	A	5/1998	Debs et al.	6,133,024	A	10/2000	Helene et al.
5,759,517	A	6/1998	Anderson et al.	6,136,601	A	10/2000	Meyer, Jr. et al.
5,763,209	A	6/1998	Sukhatme	6,140,466	A	10/2000	Barbas, III et al.
5,770,580	A	6/1998	Ledley	6,140,491	A	10/2000	Usman et al.
5,773,227	A	6/1998	Kuhn	6,143,727	A	11/2000	Roop et al.
5,773,583	A	6/1998	Sukhatme	6,150,168	A	11/2000	Woo et al.
5,780,444	A	7/1998	Kahne	6,171,859	B1	1/2001	Herrnstadt et al.
5,792,751	A	8/1998	Ledley	6,177,554	B1	1/2001	Woo et al.
5,795,870	A	8/1998	Kahne	6,184,356	B1	2/2001	Anderson et al.
5,798,209	A	8/1998	Chan	6,190,380	B1	2/2001	Abela
5,801,158	A	9/1998	Thompson et al.	6,200,801	B1	3/2001	Ferkol, Jr. et al.
5,804,604	A	9/1998	Frankel et al.	6,217,860	B1	4/2001	Woo et al.
5,807,746	A	9/1998	Lin et al.	6,238,914	B1	5/2001	Boyce
5,808,036	A	9/1998	Kool et al.	6,242,568	B1	6/2001	Barbas, III et al.
5,811,297	A	* 9/1998	Gopal 435/320.1	6,255,113	B1	7/2001	Zarling et al.
5,821,234	A	10/1998	Dzau et al.	6,262,034	B1	7/2001	Mathiowitz et al.
5,827,703	A	10/1998	Debs et al.	6,274,322	B1	8/2001	Curiel et al.
5,837,533	A	11/1998	Boutin	6,281,411	B1	8/2001	Adams et al.
5,840,674	A	11/1998	Yatvin et al.	6,287,863	B1	9/2001	Hodgson et al.
5,843,643	A	12/1998	Ratner	6,300,317	B1	10/2001	Szoka, Jr. et al.
5,844,088	A	12/1998	Hoffman et al.	6,303,582	B1	10/2001	Eljamal et al.
5,844,089	A	12/1998	Hoffman et al.	6,310,034	B1	10/2001	Woychik et al.
5,844,107	A	12/1998	Hanson et al.	6,312,699	B1	11/2001	Curiel et al.
5,847,121	A	12/1998	Yau	6,312,953	B1	11/2001	Meyer, Jr. et al.
5,851,829	A	12/1998	Marasco et al.	RE37,592	E	3/2002	Alden et al.
5,858,784	A	1/1999	Debs et al.	6,355,629	B1	3/2002	Kozak et al.
5,859,226	A	1/1999	Hunt et al.	6,362,323	B1	3/2002	Usman et al.
5,869,337	A	2/1999	Crabtree et al.	6,365,717	B1	4/2002	Middeldorp et al.
5,872,105	A	2/1999	Kool et al.	6,380,373	B1	4/2002	O'Malley et al.
5,877,302	A	3/1999	Hanson et al.	6,395,492	B1	5/2002	Manoharan et al.
5,888,762	A	3/1999	Joliot et al.	6,399,861	B1	6/2002	Anderson et al.
5,891,683	A	4/1999	Usman et al.	6,406,855	B1	6/2002	Patten et al.
				6,410,721	B1	6/2002	Hunt et al.

6,413,949	B1	7/2002	Kozak et al.
6,414,112	B1	7/2002	Buchardt et al.
6,416,998	B1	7/2002	O'Malley et al.
6,423,693	B1	7/2002	Schwartz et al.
6,468,798	B1	10/2002	Debs et al.
6,503,886	B1	1/2003	Baird et al.
6,514,747	B1	2/2003	Bultman et al.
6,528,631	B1	3/2003	Cook et al.
6,551,618	B1	4/2003	Baird et al.
6,555,668	B1	4/2003	Usman et al.
6,579,972	B1	6/2003	Lechiner et al.
6,620,617	B1	9/2003	Mathiowitz et al.
6,627,615	B1	9/2003	Debs et al.
6,683,170	B1	1/2004	Curiel et al.
6,710,035	B1	3/2004	Felgner et al.

OTHER PUBLICATIONS

SC35, accession #Q01130, NCBI database, entered Apr. 1, 1993.

Chilinae A, accession #P29022, NCBI database, entered Dec. 1, 1992.

Roget A to Z (1994). Harper Collins Pub.

Essentials of Molecular Biology (4th Ed.) Malacinski, Jones and Barlett, Eds. (2003).

Haensler, J., and Szoka, Jr., F.C. (1993) Synthesis and Characterization of a Trigalactosylated Bisacridine Compound to Target DNA to Hepatocytes. *Bioconjugate Chem.* 4, 85–93.

Rosenkranz, A.A., et al., (1992) *Experimental Cell Res.* 199, 323–329.

L. Stryer, *Biochemistry*, 3rd Ed. Freeman & Co., NY, 1988 (p. 18–20, and 28–29).

J. S. Huston et al, Protein Engineering of Single-Chain Fv Analogs and Fusion Proteins, *Methods in Enzymology*, 1991, 203:46–88.

P. Argos, An Investigation of Oligopeptides Linking Domains in Protein Tertiary Structures and Possible Candidates for General Gene Fusion, 1990, *Mol. Biol.*, 211: 943–958.

J. S. Huston et al, Protein Engineering of antibody binding sites: Recovery of specific activity in an anti-digoxin single-chain Fv analogue produced in *Escherichia coli*; 1988, *Proc. Natl. Acad. Sci USA*, 85: 5879–5883.

P. A. Silver, How Proteins Enter the Nucleus, 1991, *Cell*, 64:489–497.

Aumailley, M., et al. (1989) Cell attachment properties of collagen type VI and Arg–Gly–dependent binding to its $\alpha 2(VI)$ and $\alpha 3(VI)$ chains. *Experimental Cell Research* 181:463–474.

Berardi, A.C., et al. (1995) Functional isolation and characterization of human hematopoietic stem cells. *Science* 267:104–108.

Cameron, P.U., et al. (1992) Dendritic cells exposed to human immunodeficiency virus type-1 transmit a vigorous cytopathic infection to DC4 T cells. *Science* 257:383–387.

Carrasco, L., et al. (1982) Modification of membrane permeability in vaccinia virus-infected cells. *Virology* 117:62–69.

Caux, C., et al. (1992) GM-CSF and TNF- α cooperate in the generation of dendritic langerhans cells. *Nature* 360:258–261.

Caux, C., et al. (1994) Activation of human dendritic cells through CD40 cross-linking. *J. Exp. Med.* 180:1263–1272.

Citosky, V., et al. (1992) Nuclear localization of agrobacterium VirE2 protein in plant cells. *Science* 256:1802–1805.

Coulombel, L., et al. (1983) Enzymatic treatment of long-term human marrow cultures reveals the preferential location of primitive hemopoietic progenitors in the adherent layer. *Blood* 62:291–297.

Cotten, M., et al. (1992) High-efficiency receptor-mediated delivery of small and large (48 kilobase gene constructs using the endosome-disruption activity of defective or chemically inactivated adenovirus particles. *Proc. Natl. Acad. Sci.* 89:6094–6098.

Cotten, M., et al. (1990) Transferrin-polycation-mediated introduction of DNA into human leukemic cells: stimulation by agents that affect the survival of transfected DNA or modulate transferrin receptor levels. *Proc. Natl. Acad. Sci.* 87:4033–4037.

Curiel, D., et al. (1991) Adenovirus enhancement of transferrin-polylysine-mediated gene delivery. *Proc. Natl. Acad. Sci.* 88:8850–8854.

Dedhar, S., et al. (1987) A cell surface receptor complex for collagen type 1 recognizes the Arg–Gly–Asp sequence. *J. Cell. Biol.* 104:585–593.

DeRobertis, E., et al. (1978) Intracellular migration of nuclear proteins in xenopus oocytes. *Nature* 272:254–256.

Epanand, R., et al. (1992) Peptide models for the membrane destabilizing actions of viral fusion proteins. *Biopolymers* 32:309–314.

Eytan, G. (1982) Use of liposomes for reconstitution of biological functions. *Biochimica et Biophysica Acta* 694: 185–202.

Friedlander, D., et al. (1988) Functional mapping of cyto-tactin: proteolytic fragments active in cell-substrate adhesion. *The Journal of Cell Biology* 107(6):2329–2340.

Gardner, J., et al. (1985) Interaction of fibronectin with its receptor on platelets. *Cell* 42:439–448.

Gould-Fogerite, S., et al. (1989) Chimerasome-mediated gene transfer in vitro and in vivo. *Gene* 84:429–438.

Grant, D., et al. (1989) Two different laminin domains mediate the differentiation of human endothelial cells into capillary-like structures in vitro. *Cell* 58:933–943.

Haverstick, D., et al. (1985) Inhibition of platelet adhesion to fibronectin, fibrinogen, and von willebrand factor substrates by a synthetic tetrapeptide derived from the cell-binding domain of fibronectin. *Blood* 66(4):946–952.

Humphries, M., et al. (1987) Identification of two distinct regions of the type III connecting segment of human plasma fibronectin that promote cell type-specific adhesion. *The Journal of Biological Chemistry* 262(14):6886–6892.

Humphries M., et al. (1986) Identification of an alternatively spliced site in human plasma fibronectin that mediates cell type-specific adhesion. *The Journal of Cell Biology* 103(6):2637–2647.

Isaka, Y., et al. (1993) Glomerulosclerosis induced by in vivo transfection of transforming growth factor-B or platelet-derived growth factor gene into the rat kidney, *J. Clin. Invest.* 92:2597–2601.

Kalderson, D., et al. (1984) A short amino acid sequence able to specify nuclear location. *Cell* 39:499–509.

Kamata, H., et al. (1994) Amphiphilic peptides enhance the efficiency of liposome-mediated DNA transfection. *Nucleic Acids Research* 22(3):536–537.

Kaneda, Y., et al. (1989) Introduction and expression of the human insulin gene in adult rat liver. *The Journal of Biological Chemistry* 264(21):12126–12129.

- Kaneda, Y., et al. (1989) Increased expression of DNA cointroduced with nuclear protein in adult rat liver. *Science* 243:375–378.
- Kaneda, Y., et al. (1987) The improved efficient method for introducing macromolecules into cells using HVJ (sendai virus) liposomes with gangliosides. *Experimental Cell Research* 173:56–69.
- Kato, K., et al. (1991) Direct injection of hepatitis B virus DNA into liver induced hepatitis in adult rats. *The Journal of Biological Chemistry* 266:22071–22074.
- Kato, K., et al. (1991) Expression of hepatitis B virus surface antigen in adult rat liver. *The Journal of Biological Chemistry* 266:3361–3364.
- Klappe, K., et al. (1986) Parameters affecting fusion between sendai virus and liposomes. Role of viral protein, liposome composition, and pH. *Biochemistry* 25:8252–8260.
- Lanford, R., et al. (1986) Induction of nuclear transport with a synthetic peptide homologous to the SV40 T antigen transport signal. *Cell* 46:575–582.
- Lapidot, M., et al. (1990) Fusion-mediated microinjection of liposome-enclosed DNA into cultured cells with the aid of influenza virus glycoproteins. *Experimental Cell Research* 189:241–246.
- Lawler, J., et al. (1988) Cell attachment to thrombospondin: the role of ARG–GLY–ASP calcium and integrin receptors. *J. Cell. Biol.* 107:2351–2361.
- Liljestrom, P., et al. (1991) A new generation of animal cell expression vectors based on the semliki forest virus replicon. *Bio/Technology* 9:1356–1361.
- Marsh, M., et al. (1983) Interactions of semliki forest virus spike glycoprotein rosettes and vesicles with cultured cells. *The Journal of Cell Biology* 96:455–461.
- Maruyama, K., et al. (1990) Lipid composition is important for highly efficient target binding and retention of immunoliposomes. *Proc. Natl. Acad. Sci.* 87:5744–5748.
- Mason, P., et al. (1994) RGD sequence of foot-and-mouth disease virus is essential for infecting cells via the natural receptor but can be bypassed by an antibody-dependent enhancement pathway. *Proc. Natl. Acad. Sci.* 91:1932–1936.
- Mulligan, R.C. (1993) The basic science of gene therapy. *Science* 260:926–932.
- Neugebauer, J. (1990) Detergents: an overview. *Methods in Enzymology* 182:239–253.
- Otero, J., et al. (1987) Proteins are cointernalized with virion particles during early infection. *Virology* 160:75–80.
- Phalen, T., et al. (1991) Cholesterol is required for infection by semliki forest virus. *The Journal of Cell Biology* 112(4):615–623.
- Pierschbacher, M., et al. (1984) Cell attachment activity of fibronectin can be duplicated by small synthetic fragments of the molecule. *Nature* 309:30–33.
- Pierschbacher, M., et al. (1987) Influence of stereochemistry of the sequence Arg–Gly–Asp–Xaa on binding specificity in cell adhesion. *The Journal of Biological Chemistry* 262:17294–17298.
- Rafii, S., et al. (1994) Isolation and characterization of human bone marrow microvascular endothelial cells: hematopoietic progenitor cell adhesion. *Blood* 84:10–19.
- Rafii, S., et al. (1995) Human bone marrow microvascular endothelial cells support long-term proliferation and differentiation of myeloid and megakaryocytic progenitors. *Blood* 85:3353–3363.
- Remy, J., et al. (1995) Targeted gene transfer into hepatoma cells with lipopolyamine-condensed DNA particles presenting galactose ligands: A stage toward artificial viruses. *Proc. Natl. Acad. Sci.* 92:1744–1748.
- Roecklein, B.A., et al. (1995) Functionally distinct human marrow stromal cell lines immortalized by transduction with the human papilloma virus E6/E7 genes. *Blood* 85:997–1005.
- Roux, et al. (1991) *Oncogene* 6(11):2155–2160.
- Ruoslahti, E., et al. (1987) New perspectives in cell adhesion: RGD and integrins. *Science* 238:491–496.
- Sadler, et al. (1989) A yeast gene important for protein assembly into the endoplasmic reticulum and the nucleus has homology to DnaJ, and *Escherichia coli* heat shock protein. *J. Cell. Biol.* 109:2665–2675.
- Sands, J. (1986) Virucidal activity of cetyltrimethylammonium bromide below the critical micelle concentration. *FEMS Microbiology Letters* 36:261–263.
- Scheule, R. (1986) Novel preparation of functional sindbis virosomes. *Biochemistry* 25:4223–4232.
- Schlegel, R., et al. (1985) Biologically active peptides of the vesicular stomatitis virus glycoprotein. *Journal of Virology* 53(1):319–323.
- Schlegel, R., et al. (1983) Inhibition of VSV binding and infectivity by phosphatidylserine: is phosphatidylserine a VSV-binding site. *Cell* 32:639–646.
- Schwarzbaum, S., et al. (1984) The generation macrophage-like cell lines by transfection with SV40 origin defective DNA. *J. Immunol.* 132:1158–1162.
- Singh, P., et al. (1994) Overexpression of E2F-1 in rat embryo fibroblasts leads to neoplastic transformation. *EMBO Journal* 13:3329–3338.
- Steff, A-M., et al. (1996) Isolation and characterization of c-fos-expressing murine bone marrow stromal cell lines supporting myeloid differentiation. *Leukemia* 10:505–513.
- Stegmann, T., et al. (1989) Protein-mediated membrane fusion. *Annu. Rev. Biophys. Chem.* 18:187–211.
- Suzuki, S., et al. (1985) Complete amino acid sequence of human vitronectin deduced from cDNA. Similarity of cell attachment sites in vitronectin and fibronectin. *The EMBO Journal* 4:2519–2524.
- Tikhonenko, T., et al. (1988) Transfer of condensed viral DNA into eukaryotic cells using proteoliposomes. *Gene* 63:321–330.
- Tomita, N., et al. (1992) Direct in vivo gene introduction into rat kidney. *Biochemical and Biophysical Research Communications*. 186:129–134.
- Vaananen, P., et al. (1980) Fusion and haemolysis of erythrocytes caused by three togaviruses: semliki forest, sindbis and rubella. *J. Gen. Virol.* 46:467–475.
- Verfaillie, C.M. (1993) Soluble factor(s) produced by human bone marrow stroma increase cytokine-induced proliferation and maturation of primitive hematopoietic progenitors while preventing their terminal differentiation. *Blood* 82:2045–2053.
- Wagner, E., et al. (1992) Influenza virus hemagglutinin HA-2 N-terminal fusogenic peptides augment gene transfer by transferrin-polylysine-DNA complexes: toward a synthetic virus-like gene-transfer vehicle. *Proc. Natl. Acad. Sci.* 89:7934–7938.
- Wagner, E., et al. (1992) Coupling of adenovirus to transferrin-polylysine/DNA complexes greatly enhances receptor-mediated gene delivery and expression of transfected genes. *Proc. Natl. Acad. Sci.* 89:6099–6103.

- Wagner, E., et al. (1991) Transferrin-polycation-DNA complexes: the effect of polycations on the structure of the complex and DNA delivery to cells. *Proc. Natl. Acad. Sci.* 88:4255-4259.
- Walker, C., et al. (1992) Cationic lipids direct a viral glycoprotein into the class I major histocompatibility complex antigen-presentation pathway. *Proc. Natl. Acad. Sci.* 89:7915-7918.
- Wayner, E., et al. (1989) Identification and characterization of the T lymphocyte adhesion receptor for an alternative cell attachment domain (CS-1) in plasma fibronectin. *The Journal of Cell Biology* 109:1321-1330.
- Wickham, T.J., et al. (1995) Targeting of adenovirus penton base to new receptors through replacement of its RGD motif with other receptor-specific peptide motifs. *Gene Therapy* 2:750-756.
- Wu, G., et al. (1988) Evidence for targeted gene delivery to Hep G2 hepatoma cells in vitro. *Biochemistry* 27:887-892.
- Wu, G., et al. (1987) Receptor-mediated in vitro gene transformation by a soluble DNA carrier system. *The Journal of Biological Chemistry* 262:4429-4432.
- Wu, G., et al. (1988) Receptor-mediated gene delivery and expression in vivo. *The Journal of Biological Chemistry* 263:14621-14624.
- Young, J., et al. (1983) Interaction of enveloped viruses with planar bilayer membranes: observations on sendai, influenza, vesicular stomatitis, and semlike forest viruses. *Virology* 128:186-194.
- Yoshimura, K., et al. (1993) Adenovirus-mediated augmentation of cell transfection with unmodified plasmid vectors. *The Journal of Biological Chemistry* 268:2300-2303.
- Zhou, X., et al. (1994) DNA transfection mediated by cationic liposomes containing lipopolylysine: characterization and mechanism of action. *Biochimica et Biophysica Acta* 1189:195-203.
- "Nicholas B. La Thangue, "DRTF1/E2F: An Expanding Family Of Heterodimeric Transcription Factors Implicated In Cell-Cycle Control", [Clonexpress, Inc., pp. 108-114, (1994).]" *Trends Biochem Sci.*, pp. 108-114, (1994).
- Chaney, et al. (1986) High-frequency transfection of CHO cells using polybrene. *Somatic Cell and Molecular Genetics* 12:237-244.*
- Fink, D., et al. (1992) In vivo expression of B-galactosidase in hippocampal neurons by HSV-mediated gene transfer. *Human Gene Therapy* 3:11-19.*
- Garcia-Bustos, J., et al. (1991) Nuclear protein localization. *Biochimica et Biophysica Acta* 1071:83-101.*
- Gilboa et al. (1986) Transfer and expression of cloned genes using retroviral vectors. *Biotechniques* 4:504-512.*
- Hermonat, P., et al. (1984) Use of adeno-associated virus as a mammalian DNA cloning vector: transduction of neomycin resistance into mammalian tissue culture cells. *Proc. Natl. Acad. Sci.* 81:6466-6470.*
- Holter, W., et al. (1989) Efficient gene transfer by sequential treatment of mammalian cells with DEAE-dextran and deoxyribonucleic acid. *Experimental Cell Research* 184:546-551.*
- Jenster, G., et al. (1993) Nuclear import of the human androgen receptor. *Biochem. J.* 293:761-768.*
- Johnson, D., et al. (1993) Expression of transcription factor E2F1 induces quiescent cells to enter S phase. *Nature* 365:349-352.*
- La Thanque, N. (1994) DRTF1/E2F: an expanding family of heterodimeric transcription factors implicated in cell-cycle control. *Trends in Biochem. Sci.* 19:108-114.*
- Lau, Y., et al. (1984) Direct isolation of the functional human thymidine kinase gene with a cosmid shuttle vector. *Proc. Natl. Acad. Sci.* 81:414-418.*
- Mannino, R., et al. (1988) Liposome mediated gene transfer. *BioTechniques* 6:682-690.*
- Matheny, C., et al. (1984) The nuclear localization signal of NGFI-A is located within the zinc finger DNA binding domain. *The Journal of Biological Chemistry* 269:8176-8181.*
- McNally, M., et al. (1988) Optimizing electroporation parameters for a variety of human hematopoietic cell lines. *Biotechniques* 6:882-886.*
- Michael, S., et al. (1993) Binding-inefficient adenovirus facilitates molecular conjugate-mediated gene transfer by the receptor-mediated endocytosis pathway. *The Journal of Biological Chemistry* 268:6866-6869.*
- Miller et al. (1989) Improved retroviral vectors for gene transfer and expression. *Biotechniques* 7:980-988.*
- Miller, D., et al. (1986) Redesign of retrovirus packaging cell lines to avoid recombination leading to helper virus production. *Molecular and Cellular Biology* 6:2895-2902.*
- Morin, N., et al. (1989) Nuclear localization of the adenovirus DNA-binding protein: requirement for two signals and complementation during viral infection. *Molecular and Cellular Biology* 9(10):4372-4380.*
- Okada, C., et al. (1982) Introduction of macromolecules into cultured mammalian cells by osmotic lysis of pinocytotic vesicles. *Cell* 29:33-41.*
- Raikhel, N. (1992) Nuclear targeting in plants. *Plant Physiol* 100:1627-1632.*
- Rhim, J., et al. (1989) Neoplastic transformation of human keratinocytes by polybrene-induced DNA-mediated transfer of an activated oncogene. *Oncogene* 4:1403-1409.*
- Rothstein, L., et al. (1985) Amphotropic retrovirus vector transfer of the v-ras oncogene to human hematopoietic and stromal cells in continuous bone marrow cultures. *Blood* 65(3):744-752.*
- Schreiber, V., et al. (1992) The human poly(ADP-ribose) polymerase nuclear localization signal in a bipartite element functionally separate from DNA binding and catalytic activity. *The EMBO Journal* (9):3263-3269.*
- Stuhlmann, H., et al. (1989) Construction and properties of replication-competent murine retroviral vectors encoding methotrexate resistance. *Molecular and Cellular Biology* 9(3):100-108.*
- Takai, T., et al. (1990) DNA transfection of mouse lymphoid cells by the combination of DEAE-dextran-mediated DNA uptake and osmotic shock procedure. *Biochimica et Biophysica Acta* 1048:105-109.*
- Tratschin, J., et al. (1985) Adeno-associated virus vector for high-frequency integration, expression, and rescue of genes in mammalian cells. *Molecular and Cellular Biology* 5:3251-3260.*
- van der Krol, A., et al. (1991) The basic domain of plant B-ZIP proteins facilitates import of a reporter protein into plant nuclei. *The Plant Cell* 3:667-675.*
- Zwiebel, J., et al. (1989) High-level recombinant gene expression in rabbit endothelial cells transduced by retroviral vectors. *Science* 243:220-222.*

PEPTIDE-MEDIATED GENE TRANSFER

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.

RELATED APPLICATION

This application was filed as a reissue of Ser. No. 08/240, 514, filed May 14, 1994, now U.S. Pat. No. 5,670,347.

BACKGROUND OF THE INVENTION

The present invention is directed to a methodology for highly efficient, stable integration of DNA into a eukaryotic genome. More specifically, the present invention is directed to the use of a synthetic polypeptide, containing a nuclear localization signal to complex with a DNA molecule and to facilitate its transportation and integration into the nuclear genome of a mammalian or other eukaryotic cell, for example, in the context of producing cell lines with an extended life.

DNA-CaPO₄ co-precipitation was the first method developed to introduce genes into mammalian cells. ("Gene" in this regard denotes a structural DNA segment, i.e., a DNA that codes for a polypeptide, and comprehends oncogenes as well as DNAs coding for a known expression product.) The co-precipitation method was applicable only to certain cell types, however, and could not be used to introduce genes into a wide variety of cell lines, especially those of hematopoietic origin. Moreover, the stable gene transfer efficiency was rather low, on the order of 10⁻⁴ to 10⁻⁶. McNally, M. A., et al., *BioTechniques* 6: 8826 (1988); Yen, T. S. B., et al., loc. cit. 6: 413 (1988).

Limits on introducing and expressing genes in cultured mammalian cells motivated a search for other, more efficient approaches to gene transfer. Methods were developed, for example, that utilized chemical agents which were positively charged and, hence, able to complex with negatively charged DNA molecules. Examples of such agents include DEAE dextran and various cationic lipid molecules. Cells treated with DNA complexes comprised of such an agent can lead to the introduction of the DNA into different mammalian cell lines. Mannino, R. J. et al., *BioTechniques* 6: 682 (1988); Felgner, P. et al., *Proc. Nat'l Acad. Sci. USA* 84: 7413 (1987); Fraley, R. et al., *Trend Biochem. Sci.* 6: 77 (1981); Holter, W. et al., *Exp. Cell Res.* 184: 546 (1989); McCutchan, J. H. et al., *J. Nat'l Cancer Inst.* 41: 351 (1986); Chaney, W. C. et al., *Somatic Cell & Mol. Genet.* 12: 237 (1986).

The production of a gene product for only a short time period after transfection, usually from 48 to 72 hours, is called "transient expression." Many of the DNA-complexing agents reported heretofore, while useful in transferring a gene into mammalian cells, resulted in only transient expression of the introduced gene in a small fraction of the transfected cells. See, for example, Miller et al., *Proc. Nat'l Acad. Sci., USA*, 76: 949 (1979); Oi et al., loc. cit. 80: 825 (1983).

In addition to giving poor results with respect to stable gene expression, transfer methods based on such DNA-complexing agents often were effective only with established cell lines, and did not work very well with primary cells isolated from various mammalian species. Other techniques therefore were needed to enhance gene transfer efficiency, to increase the variety of cell types capable of being transfected, and to effect stable gene transfer. Stable

gene transfer is the ability of cells to maintain and express transfected DNAs in a stable manner, through integration of the transfected DNA into cell chromosomes.

Retroviral vectors, which were under development at about the same time seemed to be quite effective in transferring genes into different cell types. The use of such vectors was prompted by the elucidation of gene regulation in various murine and avian retroviruses. Two other developments led to the development of retrovirus-based gene transfer vehicles. The first development was the identification of minimal sequences required for efficient packaging of viral particles in a cell line which produced the coat proteins and other structural components of the viral particle in trans. The cell lines that provided the structural components for virus development are called "packaging" cell lines. The second significant step in the establishment of retroviral vectors was the development of both ecotropic and amphotropic packaging cell lines, which aided the design of recombinant retroviral particles which could infect both murine and human cell lines.

Additional modifications of retroviruses were deemed necessary to address concerns that retroviral vectors could recombine in vivo to generate wild-type virus. Developments in this regard yielded a number of safe retroviral vectors which have been used to transfer genes into a variety of established mammalian cell lines, as well as into certain primary cells in a few instances. E. Gilboa et al., *BioTechniques* 4: 504 (1986); A. D. Miller et al., *Mol. Cell. Biol.* 6: 2895 (1986); H. Stuhlmann et al., loc. cit. 9: 100 (1989); A. D. Miller et al., *BioTechniques* 7: 980 (1989); J. A. Zwiebel et al., *Science* 243: 220 (1989).

Even though these vectors were effective with respect to various mammalian cells, there were many restrictions on a wider application of the retroviral gene-transfer technique. These limitations included (1) the size of exogenous DNA that can be inserted into a retroviral vector and (2) the use of only dividing cells for retroviral gene transfer. E. Gilboa, *BioTechniques*, supra (1986); A. D. Miller, supra (1986); H. Stuhlmann, et al., *Mol. Cell. Biol.* supra, (1986); A. D. Miller et al., *BioTechniques*, supra (1986); J. A. Zwiebel et al., supra (1989).

Other viruses have been used to generate recombinant viral vectors for gene transfer studies. Adenovirus, adeno-associated virus, herpes simplex virus, and even HIV have been employed as vectors to introduce genes into both established cell lines and primary cells. Some of these viral vectors are capable of transferring genes into non-dividing cells. R. J. Samulski, et al., *EMBO J.* 10: 3941 (1981); J. D. Tratschin, et al., *Mol. Cell. Biol.* 5: 3251 (1985); P. L. Hermonat, et al., *Proc. Nat'l Acad. Sci. (USA)* 81: 6466 (1984); D. J. Fink, et al., *Human Gene Therapy* 3: 11 (1992).

Viral vectors capable of transferring genes into non-dividing cells usually require the generation of high-titer viral stock in order to achieve high efficiency gene transfer into different cell types. In addition, whenever a different regulatory sequence is to be tested for optimal level of gene expression into primary cells, a new viral stock must be made and tittered for every modification. All these involve very time-consuming experimental manipulations.

Still another concern relates to the application of viral vectors in human gene therapy. A number of studies have been carried out in primates to test the safety of retroviral vectors for introducing cells transduced with retroviral vectors into animals. Some of these animals have developed various forms of lymphoma. R. E. Donahue, et al., *J. Exp. Med.* 176: 1125 (1992). Additional safety features have been

introduced into some of the newer versions of retroviral vectors, yet are not available for all types of viral vectors.

SUMMARY OF THE INVENTION

It therefore is an object of the present invention to provide a method for high efficiency gene transfer to achieve expression, stable as well as transient, in a wide spectrum of cell types, including primary cells from various mammalian species.

It is also an object of the present invention to provide cell lines which, even if derived from primary mammalian cells, are characterized by an extended life in culture.

It is another object of the present invention to provide a readily implemented screening system for identifying sequences that influence in the expression of cloned genes in various primary cell types from different species.

In accomplishing these and other objectives, there has been provided, in accordance with one aspect of the present invention, a transfection vector comprising a synthetic polypeptide linked electrostatically to a DNA structural sequence, forming a polypeptide-DNA complex, where the polypeptide is comprised of (A) a polymeric chain of basic amino acid residues, (B) an NLS peptide and (C) a hinge region of neutral amino acids that connects the polymeric chain and the NLS peptide. The polymeric chain preferably is comprised of between 10 and 50 residues, which can be selected from lysine, arginine and ornithine, for example, while the hinge region is comprised of between 6 and 50 amino acid residues selected, for example, from glycine, alanine, leucine and isoleucine. The NLS peptide preferably is located at the amino terminus of said polypeptide and the polymeric basic amino acid chain at the carboxyl terminus. Among exemplary NLS peptides are the SV40 large T antigen NLS sequence, the polyoma large T antigen NLS sequence, the adenovirus E1a NLS sequence, and the adenovirus E1b NLS sequence.

In accordance with another aspect of the present invention an extended life cell line is provided that is the product of transfecting a mammalian cell with a vector as described above. The mammalian cell thus transfected can be selected, for example, from the group consisting of a human umbilical vein endothelial cell, a human dermal microvascular endothelial cell, a human peripheral blood monocyte/macrophage

cell, a human aortic smooth muscle cell, and a rabbit liver non-parenchymal cell.

Other object, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention allows for the highly efficient transfer and stable integration of DNA into eukaryotic cells, such as cells from established mammalian cell lines, primary cells from mammalian tissues, and plant cells. The present invention also can be applied to developing cell lines from non-dividing cells, such as human peripheral blood monocytes and macrophages.

In accordance with the present invention, a synthetic polypeptide is provided that can complex with a DNA molecule very efficiently by taking advantage of the high negative charge density on the polynucleotide. To this end, a synthetic polypeptide of the present invention comprises a DNA-binding sequence that is rich in basic amino acids, such as lysine, arginine and ornithine, and that is typically ten to fifty residues long. D-isomers of these basic amino acids are suitable so long as the length of the stretch of basic amino acids is within the prescribed length. The DNA-binding sequence can be a homopolymer of a basic amino acid, or it can comprise more than one kind of basic residue. The DNA binding sequence must be of adequate length to bind DNA, yet not so long that it precipitates out of the solutions employed in the present methodology, as discussed below.

A synthetic polypeptide of the present invention also contains an amino acid sequence corresponding to a nuclear localization signal (NLS) sequence. A representative sample from the diverse range of nuclear localization signals which have been identified are listed in Table I below. (SEQ ID NOS:1-54).

TABLE 1

Source	Nuclear Protein	Deduced Signal Sequence
Yeast	MAT α 2	(SEQ ID NO: 1) K--I--P--I--K (SEQ ID NO: 2) V--R--I--L--E--S--W--F--A--K--N--I
SV40	Large T	(SEQ ID NO: 3) P--K--K--K--R--K--V
Influenza Virus	Nucleoprotein	(SEQ ID NO: 4) A--A--F--E--D--L--R--V--R--S
Yeast	Ribosomal protein L3	(SEQ ID NO: 5) P--R--K--R
Polyoma virus	Large T	(SEQ ID NO: 6) V--S--R--K--R--P--R--P--A
SV40	VP1	(SEQ ID NO: 7) A--P--T--K--R--K
Adenovirus	E1a	(SEQ ID NO: 8) K--R--P--R--P
SV40	VP2 (VP3)	(SEQ ID NO: 9) P--N--K--K--K--R--K
Frog	Nucleoplasmin	(SEQ ID NO: 10) R--P--A--A--T--K--K--A--G--Q--A--K--K--K--L--D
Rat	Glucocorticoid receptor	(SEQ ID NO: 11) K--K--K--I--K
Monkey	v-sis (PDGF B)	(SEQ ID NO: 12) R--V--T--I--R--T--V--R--V--R--R--P--P--K--G--K--H--R--K
Yeast	Histone 2B	(SEQ ID NO: 13) [G--K--K--R--S--K--A]D--G--K--K--W--S
Chicken	v-re1	(SEQ ID NO: 14) K--A--K--R--[S--K--A]Q--R
Influenza	NS1	(SEQ ID NO: 15) D--R--L--R--R (SEQ ID NO: 16) P--K--Q--K--R--K

TABLE 1-continued

Frog	N1	(SEQ ID NO: 17) V--R--K--K--R--K--T (SEQ ID NO: 18) A--K--K--S--K--Q--E
Human	c-myc	(SEQ ID NO: 19) P--A--A--K--R--V--K--L--D (SEQ ID NO: 20) R--Q--R--R--N--E--L--K--[4]R--S--F
Human	lamin A	(SEQ ID NO: 21) T--K--K--R--K--L--E
HTLV-1	Rex(p27 ^{x-III})	(SEQ ID NO: 22) P--K--T--R--R--R--P (SEQ ID NO: 23) S--Q--R--K--R--P--P
Adenovirus	_F TP	(SEQ ID NO: 24) R--L--P--V--R--R--R--R--V--P
HIV-1	Tat	(SEQ ID NO: 25) G--R--K--K--R
Frog	Lamin L ₁	(SEQ ID NO: 26) V--W--T--T--K--G--K--R--K--R--I--D--V
Rabbit	Progesterone receptor	(SEQ ID NO: 27) R--K--F--K--K
HIV-1	Rev	(SEQ ID NO: 28) R--R--N--R--R--R--R--W
Human	PDGF A-chain	(SEQ ID NO: 29) P--R--[3]E--S--G--K--K--R--K--R--K--R--L--K--P--T
Mouse	c-ab1	(SEQ ID NO: 30) [K--K--K--K--K]S--A--L--I--K--K--K--K--K--M--A--P
Adenovirus	DBP	(SEQ ID NO: 31) P--P--K--K--R (SEQ ID NO: 32) P--K--K--K--K--K
Chicken	c-erb-A	(SEQ ID NO: 33) S--K--R--V--A--K--R--K--L
Human	c-myb	(SEQ ID NO: 34) P--L--L--K--K--I--[1]K--Q
Human	N-myc	(SEQ ID NO: 35) P--P--Q--K--K--I--K--S
Human	p53	(SEQ ID NO: 36) P--Q--P--K--K--K--P
Human	Hsp 70	(SEQ ID NO: 37) F--K--R--K--H--K--K--D--I--S--Q--N--K--R--A--V--R--R
Hepatitis B virus	Core protein	(SEQ ID NO: 38) S--K--C--L--G--W--L--W--G
Chicken	Ets1	(SEQ ID NO: 39) G--K--R--K--N--K--P--K
Yeast	Ribosomal protein L29	(SEQ ID NO: 40) K--T--R--K--H--R--G (SEQ ID NO: 41) K--H--R--K--H--P--G

Protein	Nuclear Localization Signals
TGA-1A (tobacco)	(SEQ ID NO: 42) R--R--L--A--Q--N--R--E--A--A--R--K--S--R--L--R--K--K
TGA-1B (tobacco)	(SEQ ID NO: 43) K--K--R--A--R--L--V--R--N--R--E--S--A--Q--L--S (SEQ ID NO: 44) R--Q--R--K--K
O2 NLS B (maize)	(SEQ ID NO: 45) R--K--R--K--E--S--N--R--E--S--A--R--R--S--R--Y--R--K
NIa (Polyvirus)	(SEQ ID NO: 46) K--K--N--Q--K--H--K--L--K--M-32aa-K--R--K
VirD2 (Agrobacterium)	(SEQ ID NO: 47) K--R--P--R--E--D--D--D--G--E--P--S--E--R--K--R--E--R
VirE2 NSE1 (Agrobacterium)	(SEQ ID NO: 48) K--L--R--P--E--D--R--Y--I--Q--T--E--K--Y--G--R--R
VirE2 NSE2 (Agrobacterium)	(SEQ ID NO: 49) K--T--K--Y--G--S--D--T--E--I--K--L--[L]K--S--K
O2 NLS A (maize)	(SEQ ID NO: 50) M--E--E--A--V--T--M--A--P--A--A--V--S--S--A--V--V--G--D--P (SEQ ID NO: 51) M--[3]E--Y--N--A--I--L--R--R--K--L--E--E--D--L--E
R NLS A (maize)	(SEQ ID NO: 52) G--D--R--R--A--A--P--A--R--P
R NLS M (maize)	(SEQ ID NO: 53) M--S--E--R--K--R--R--E--K--L
RNLS C (maize)	(SEQ ID NO: 54) M--I--S--E--A--L--R--K--A--I--G--K--R

See Garcia-Bustos et al., Biochem. Biophys. Acta 1071: 83 (1991). Raikhel, N., Plant Physiol. 100: 1627 (1992), and Citovsky, V. et al., Science 256: 1802 (1992), the contents of each of which are hereby incorporated by reference.

In the present invention, an NLS peptide, which typically is six to fifteen amino acids in length, facilitates transport of the associated DNA into the nucleus. Because the synthetic polypeptide promotes the transport of the transfected gene into the nucleus of the host cell, this method provides both highly efficient stable and transient gene expression. Once inside the nucleus, the introduced DNA is immediately available to the transcription machinery, and can be expressed transiently. Simultaneously, the introduced DNA is also in the process of getting integrated into the host chromosome to give rise to stable expression. Thus, the method of the instant invention can achieve both transient and stable expression of introduced DNA.

50

Transient gene expression results when the method of gene transfer results in the introduction of the DNA sequences into the nucleus in an non-integrated form. Transient transfection is measured 24 to 72 hours after transfection by assays that measure gene expression of the transfected gene(s). In contrast, stable expression of the encoded protein results when the transferred DNA sequences are stably integrated into the chromosomal DNA of the target cell. Stable transfectants remain capable of expressing the transfected DNA after two weeks or greater following the method of the invention. Commonly used assays monitor enzyme activities of chloramphenicol acetyltransferase (CAT), LAC-Z, β -galactosidase (β -gal), β -glucuronidase (GUS), luciferase, or human growth hormone, each of which may be contained in the present invention.

65

The NLS domain of the synthetic peptide is based on known endogenous peptide sequences that were identified by reference to two criteria: (1) sufficient to redirect a

cytoplasmic protein to the nucleus and (2) necessary for directing a nuclear protein to the nucleus. Methods for assessing an NLS peptide's ability to direct protein to the nucleus are known in the art. See Garcia-Bustos, et al., supra, Sandler et al., J. Cell Biol. 109: 2665 (1989), and Citovsky et al., supra, the respective contents of which are hereby incorporated by reference. For example, an NLS peptide or a natural protein containing an NLS is fused to an otherwise non-nuclear protein, by either synthetic or recombinant production. The hybrid protein is then assessed for its ability to target the non-nuclear protein to the nucleus.

The presence of the non-nuclear protein in the nucleus can be determined by a functional assay or immunofluorescence. An illustrative assay entails the histochemical determination of a product produced by the non-nuclear protein, such as a colorimetric marker produced by β -gal or GUS. (A "colorimetric marker" includes an enzyme that can catalyze a reaction with a substrate to elicit a colored product which can be detected or measured by a variety of means, such as standard fluorescence microscopy, flow cytometry, spectrophotometry or colorimetry. "Immunofluorescence" relates to detecting the presence of the non-nuclear protein in the nucleus by means of an antibody specific for the targeted protein.)

In the past NLS peptides have been studied to assess their ability to target reporter proteins to the nucleus. Also, endogenous proteins containing an NLS, such as the VirD2 and VirE2 of *Agrobacterium*, have been shown to mediate the transfer of the *Agrobacterium* single-stranded DNA intermediate T-strand to the plant cell nucleus endogenously. See Citovsky, et al., supra. There has been no suggestion heretofore, however, to use an NLS peptide to target a polynucleotide to the nucleus of a eukaryotic cell.

A preferred NLS domain contains a short stretch of basic amino acids like the NLS of the SV40 virus large T antigen (PKKKRKV) (SEQ ID NO:3), which is an NLS that has been shown to be effective in mammalian cells (basic residues are highlighted). Another preferred NLS domain consists essentially of short hydrophobic regions that contain one or more basic amino acids (KIPK) (SEQ ID NO:1), which is like the NLS of mating type α 2. The NLSs that transport DNA into the plant cell nucleus often are bipartite, which means that they are usually comprised of a combination of two regions of basic amino acids (*identified as stippled segments in Table I*) separated by a spacer of more than four residues [(see stippled segments in Table I)], such as the *Xenopus* nucleoplasmin (KRPAATKKAGQAKKKK) (SEQ ID NO:55).

The NLS peptide of the present invention can be designed to accommodate different host cells, both mammalian and plant cell hosts.

The method described here can suitably be modified to introduce genes into plant protoplasts using plant NLSs, such as those described by Raikhel (1989), supra.

The present gene transfer system is also capable of transferring foreign DNA into gymnosperms and angiosperms. Procedures for assessing the introduction of foreign DNA in plants are known to the art, such as those disclosed by Miki, B. L., et al., in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY, B. R. Glick et al., eds. (CRC Press, 1993), and Gruber, M. Y. et al., id.

A synthetic polypeptide of the present invention thus is comprised of a DNA binding domain and an NLS peptide domain which are separated by a third element, a hinge region of neutral amino acid, to minimize [stearic] steric interference between the two domains. For this purpose, the

hinge region ranges in length from about six to twenty-five amino acids, and contains a stretch of neutral small amino acids without any bulky hydrophobic or ionic side chains.

The NLS sequence can be located at either the amino terminus or the carboxy-terminus of the synthetic peptide. The arrangement of the two domains, basic amino acid sequence and NLS sequence can be interchanged without affecting the high gene transfer efficiency. As indicated previously, such a synthetic polypeptide binds electrostatically to the DNA that is to be introduced into the target cell. The weight ratio of polypeptide to DNA in the resulting complex generally is in the range of 1:1 to 1:10; for example, 1 μ g polypeptide to 1 to 10 μ g of polynucleotide.

In accordance with the present invention, entry of the DNA-polypeptide complex into cells can be promoted by treating target cells with a hypertonic solution, followed by hypotonic treatment of cells in the presence of gene-peptide complex. See, for example, Okada and Rechsteiner, Cell 29: 33 (1982). A suitable hypertonic solution can contain both polyethylene glycol (PEG) and sucrose, preferably in the concentration of 0.3M–0.6M and 10% to 25%, respectively, and is referred to as "primer" hereinafter. Okada et al., supra, and T. Takai, et al., Biochem. Biophys. Acta 1048: 105 (1990).

The methodology of the present invention has been used to develop stable transfectants of different established cell lines. It also has been employed to transfer genes into primary cells from different mammalian species, thereby to obtain cell lines that retain many of the characteristics of the cognate primary cells. Cell lines developed from primary cells via the methodology of the present invention are called "extended life" cell lines in this description, because the cell lines so developed retain almost all of the characteristics of their cognate primary cells even in their late passage. The range of cell types that can be converted to extended life cell lines, according to the present invention, is based on the availability of primary cells or the ability to isolate a primary cell from the organ in question. In this regard, the inventive methodology is not limited to cell types amenable to transformation. In addition to the cell types already mentioned, the present invention can be applied to pancreatic beta cells, human liver and kidney cells, and human hematopoietic stem cells, among others.

The methodology of the present invention has been used to develop an extended life cell line from human monocyte/macrophage cells, which are normally non-dividing. In all these instances, stable cell lines were obtained with a very high efficiency, either comparable to or better than the efficiency using retroviral vectors.

The present invention finds application as well in both ex vivo and in vivo gene therapies, where genetic material is transferred into specific cells of a patient. Ex vivo gene therapy entails the removal of the relevant target cells from the body, transduction of the cells in vitro, and subsequent reintroduction of the modified cells into the patient.

A gene therapy pursuant to the present invention could involve an ex vivo introduction, into a particular cell type from the patient, of a polynucleotide coding for a correcting protein which can be produced in functional form by the targeted cell type. Genes suitable for expression in this regard include an adenosine deaminase gene, a globin gene, an LDL receptor gene, and a glucose cerebrosidase gene.

Different kinds of gene-therapy applications require either stable or transient gene expression. The method of the present invention is advantageous in that it can be used in gene therapy requiring either stable gene expression or transient gene expression. Transient expression of a foreign

gene is preferred when expression of the exogenous product is needed only for a short period of time; thereafter, rapid clearance of the gene product and its vector is desirable. Transient expression is also desirable when the prolonged effects of the exogenous protein's expression are unknown. Stable expression in gene therapy is needed when the patient has a genetic defect that is incompatible with life. Such genetic defects include but are not limited to cystic fibrosis, Tay Sachs and cancer. Mulligan, *Science*, 260: 926 (1993).

A gene therapy pursuant to the present invention also could involve an in vivo introduction of a structural DNA into cells of a patient's body. For stable transfer of genes into a target tissue using this method, the ligand to the target receptor will be conjugated to the synthetic polypeptide. The polypeptide-ligand combination can be complexed to a polynucleotide coding for the needed protein and then introduced into the host organism through blood circulation. When this complex reaches the target tissue, the whole complex will be taken up by cells containing the corresponding receptor for the ligand through receptor mediated process. Because of the NLS in the polypeptide-ligand complex, the complex will enter into the nucleus, resulting in a stable integration of the introduced gene into the host chromosome and, thereby, a correction of the genetic defect in the host. Cell-specific receptors are well known to those of skill in the art, as are their ligands which can be used in complexes for receptor-mediated gene transfer. Michael, S. I, et al., *J. Biol. Chem.* 268: 6866 (1993). For example, when the liver is the tissue targeted for gene therapy, the DNA encoding corrective protein is complexed to a synthetic neoglycoprotein that will target the complex to the asialoglycoprotein receptor on hepatocytes. For example, a cell type specific [receptor] ligand such as asialoglycoprotein can be chemically linked to the transfection vector at the carboxyl terminal of the synthetic polypeptide molecule to deliver the foreign gene directly into liver cells. An additional hinge region can be incorporated into the molecule before chemically linking the polypeptide molecule to a cell-type specific ligand molecule, such as asialoglycoprotein or a cell-specific monoclonal antibody.

An example of a carrier useful for receptor-mediated gene transfer to liver is a synthetic glycoprotein in which bovine serum albumin (BSA) is covalently bound to poly L-lysine using 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC). Ferkol et al., *FASEB* 7: 1081 (1993). To produce a neoglycoprotein conjugate for use in targeting DNA to liver, a reaction mixture that contains about 170 mM galactose, 4 mM poly (L-lysine), 160 mM BSA and 10 mM EDC (pH 7.5) can be incubated for 48 hours at 22° C. DNA is complexed to the neoglycoprotein carrier in a 360:1 molar ratio. The carrier-DNA complexes are dialyzed against 150 mM sodium chloride before transfection.

Expression of a functional protein after transfection with DNA complexed to ligand alone is often transient. Ferkol et al., supra. The method of the present invention greatly improves the cell-specific targeting of receptor-mediated transfection by providing stable expression by increasing stable integration of a foreign DNA in the host cell using a synthetic polypeptide molecule of the present invention.

A variation of receptor-mediated gene transfer employs coupling a synthetic polypeptide as described above to monoclonal antibodies which recognize a cell surface antigen on the target cells. Maruyama et al., *Proc. Nat'l Acad. Sci. USA* 87: 5744 (1990). The coupled monoclonal antibody and synthetic polypeptide then are complexed with a DNA encoding the required or desired protein. This complex will target the DNA to the cells expressing the corresponding

cell surface antigen. Any tissue of the human body can be targeted for the gene therapy of the present invention using the disclosed methods. A target tissue is suitable in this context so long as it is susceptible to genetic modification according to the present invention.

The present invention is further described with reference to the following examples, which are only illustrative and not limiting of the invention.

EXAMPLE 1

Transfer of Genes into Established Cell Lines for Purposes Transient Gene Expression and Selection of Stable Transfectants, Respectively

Forming the DNA-Polypeptide Complex and Transfecting Cells Therewith

The DNA or other polynucleotide to be transfected, such as a plasmid containing a gene for a drug resistance marker or coding a protein needed for expression in the host cell, is complexed to a synthetic polypeptide molecule in different weight ratios in an isotonic buffer solution. For example the weight ratio of DNA:polypeptide can be between 1:1 and 10:1, although ratios outside of this range may be evaluated empirically for achieving the objects of the present invention. An isotonic buffer solution such as Hanks buffered salt solution or HEPES buffered saline may be used for complexing DNA to polypeptide.

While the complex is formed, the cells that are to be transfected either remain attached to a substratum, such as a tissue culture dish, or are pelleted (for cells that grow in suspension). The cells are treated with a hypertonic primer solution, such as a concentration of 0.3M–0.6M sucrose and 10% PEG in either Tris-HCl or HEPES (pH 7.2) buffered solution, for 3–5 mins at room temperature. The primer solution then is removed.

After the DNA-polypeptide complex is formed, it is made hypotonic. The complex solution is hypotonic when it has a lesser osmotic pressure than a 0.15M or 0.9% solution of NaCl. For example, the complex in isotonic buffer can be made 40–55% hypotonic or 0.075M simply by adding an amount of distilled water that is equal to the volume of the complex in isotonic buffer. The hypotonic complex solution then is added to the cells that have been treated with the primer solution. Cells remain in the hypotonic DNA-polypeptide solution for 3–4 minutes. Fresh medium then is added to the cells to rinse away excess DNA-polypeptide solution. Thereafter, the cells are grown normally.

Producing a Synthetic Polypeptide Molecule

An example of an synthetic polypeptide molecule of the present invention is one consisting of the amino acid sequence PPKKRKVS GGGGGK KKKKKKKKKKKK (SEQ ID NO:56). Such a peptide can be synthesized, using standard methods of peptide production, and purified by standard methods using high pressure liquid chromatography (HPLC).

Selection of Stably Transfected Cell Lines

Transfected cells are grown in regular growth medium for 48 hours, and then plated in selective medium containing 400 µg/ml of G418. Cells were plated at a density of 100–1000 cells per 60 cm² dish. The number of G418-resistant colonies was determined two weeks after the initiation of selection. Other selectable markers, such as pHyg, may be used to achieve the results of the instant invention. K. Blochlinger, et al., *Mol. Cell. Biol.* 4: 2929 (1984).

11

This method gave a stable-transfection efficiency of 5–10%. Similar results were obtained using either G418 or hygromycin selection. In general the stable transfection efficiency achieved by the method of the instant invention is a few orders of magnitude greater than prior art methods. The instant invention's 5–10% efficiency is several orders of magnitude better than the efficiency of the DNA-CaPO₄ co-precipitation method and at least equal or 5 times greater than the fairly high 1–10% level of stable transfection efficiency achieved by viral based methods. *See Table II.*

TABLE II

Method	Stable Transfection Efficiency
Peptide-Mediated Gene Transfer	5–10%
Retroviral vectors	1–10%
Non-viral methods (e.g., CaPO ₄)	<2%

That the transfectants of the instant invention are stable is shown by the following example. When G418 resistant colonies were grown without selection for variable period of times, and then tested for resistance to the antibiotic by plating the cells under clonal conditions, the same number of colonies were obtained both with and without G418. This result indicates that, once the cells are selected for the expression of the Neo gene, the resistance gene was retained stably in the chromosome.

Three different cell lines were used to test the efficiency of gene transfer of the new method. Mouse fibroblast cell line (L cells), mouse erythroleukemia cell line (C19TK), and COS cells. The COS cell line was used to establish conditions for transient gene expression. The eukaryotic expression vector, CH110, contains bacterial β -gal and was employed in these studies. The β -gal gene in CH110 is under the control of SV40 virus early promoter.

The COS cells were treated with primer and then exposed to DNA-polypeptide (2.5–5.0 μ g) complex under hypotonic conditions. After this treatment, cells were returned to the normal growth condition. Transfected cells were grown at 37° C. for 48 hrs, and stained for the expression of the β -gal reporter gene. Forty to fifty percent of the cells were positive for the expression of the reporter gene.

Mouse L cells were transfected with eukaryotic expression vector containing the Neo gene, which codes for the antibiotic G418 resistance gene. L cells are sensitive to G418 at 400 μ g/ml. Cells plated in 24-well tissue culture plates were then transfected with synthetic polypeptide complexed to the plasmid pRSV-Neo via the methodology of the present invention.

A mouse erythroleukemia cell line, C19TK, also was used as a representative cell line for testing the transfection efficiency of the present invention with respect to hematopoietic cells. The expression vector, pDR2, which carries a hygromycin-resistance gene, was used for these studies, C19TK cells are exquisitely sensitive for the antibiotic hygromycin. This cell line grows in suspension and, hence, was transfected in suspension.

Briefly, about million cells are spun down and the cell pellet is treated with primer. The cells are then exposed to DNA-polypeptide complex under hypotonic condition. Forty-eight hours after transfection, a known number of cells are plated in microtiter plates with hygromycin. The number of wells with growing population of cells was enumerated to determine the transfection efficiency. The

12

stable transfection efficiency was about 1–5%, as compared to most of the other non-virus-based methods that are very poor. Thus, the method described herein is very efficient for stable transfection efficiency both for hematopoietic and non hematopoietic cell lines. Only some retrovirus based vectors give a transfection efficiency comparable to the efficiency obtained with the current method for hematopoietic cell lines. See Gilboa, et al. (1986), Miller, et al. (1986), Stuhlmann, et al. (1989), Miller, et al. (1989), and Zwiebel, et al. (1989), each cited above.

EXAMPLE 2

Transfer of Genes into Human Primary Cells

The gene transfer method of the present invention was used to generate extended life cell lines from different human primary cells. Most of the primary cells have a limited in vitro life span. The following cell types were employed to test the efficacy of the inventive method to generate extended-life cell lines by transfer of various oncogenes, either singly, in pairs of combinations, or combinations of more than two oncogenes. Rhim, J. S., et al., *Oncogene 4: 1403 (1989)*.

EXAMPLE 3

Production of Extended Life Cell Lines

The method of introducing genes into primary cells is the same as that described above for introducing genes into established cell lines, such as the mouse fibroblast cell line L cells and the mouse erythroleukemia cell line C19TK. The main difference is that the host cell is a primary cell isolated from different species, human or other mammalian species, and the primary cells have only a limited in vitro life span. The isolation of primary cells from various tissue sources are well known to those of skill in the art.

In order to extend the life of primary cells that are endogenously incapable of extended growth in vitro, the cells are transfected with different oncogenes, such as SV40 large T antigen, polyoma large T antigen, adenovirus E1A and E1B, v-fms, Bc12, myc and ras. The oncogenes can be used either alone, in pairs of various combinations, or in combinations of more than two oncogenes.

In addition, other genes that do not come under the category of oncogenes may be used. For example, genes that are important for DNA synthesis and normally active during the S phase of the cell cycle, such as the dihydrofolate reductase gene (DHFR), thymidine kinase gene, thymidylate synthetase gene, a DRTF1/E2F transcription factor encoding DNA, or DNA encoding the E2F transcription factor can be complexed to synthetic polypeptide and used to extend the life of primary cells. The human DHFR gene complexed to synthetic polypeptide can be introduced into primary cells to produce extended life cell lines. DNA encoding a transcription factor that is active during the S phase of the cell cycle are particularly useful in the method of the instant invention. La Thangue, N. B. *Trends in Biochemical Sciences 19: 108 (1994)*; Johnson, D. G. et al., *Nature 365: 349 (1993)*, the respective contents of which are hereby incorporated by reference.

Because untreated primary cells have only a limited life span in vitro, their ability to grow continuously in culture after treatment with the present invention served to select for extended life cell lines. No other drug selection markers need to be used to select for extended life cell lines derived from primary cells.

To produce extended life cells lines from primary cells, newly cultured primary cells were treated by the method of the present invention employing synthetic polypeptide conjugated with various oncogenes, such as SV40 large T antigen and/or Adeno E1A. The treated cells were plated in their appropriate growth media and passed after the cells reached confluency. A parallel set of a control untreated primary cells were cultured under the same growth conditions. Typically, control primary cells stop growing after about 4–10 passages, depending upon the cell type (cell split ratio was usually 1:4 by surface area). In contrast, continuously growing cell lines were obtained from different primary cell types described in the following examples.

EXAMPLE 4

Analysis of Transformed (Extended Life) Cells

Extended life cell lines containing the oncogene are identified by restriction cleavage. Southern analysis and/or Northern analysis using appropriate DNA probes.

The DNA of each transformed extended life cell line is analyzed by Southern hybridization to determine whether the cell lines carry the oncogenes used to establish such extended life cell lines. DNA is extracted from the cell lines and the nucleic acid pellet is re-suspended in 200 μ l of 10 mM Tris-Cl pH 7.4, 0.1 mM EDTA, and 10 μ g is digested with a specific restriction enzyme, electrophoresed through 1.0% agarose, and transferred to nitrocellulose. Southern, J. Mol. Biol. 98: 503 (1975). Filters are hybridized to a radioactively labelled DNA, encoding each of the oncogenes that gave rise to the corresponding extended life cell line, in the presence of 10% dextran sulfate. After overnight hybridization, the filters were washed twice in 2 \times SSC, 0.1% SDS at 64° C.

Each transformed extended life cell line is analyzed by Northern hybridization to determine whether the cell lines transcribe the oncogenes. Cells not containing the oncogene of interest will not demonstrate transcripts in a Northern analysis whereas cells containing the DNA of interest will demonstrate a detectable transcript. Also, an ELISA method was used to detect the presence of oncogene products in some of the extended life cell lines, using publicly available antibodies that recognize the corresponding oncogene protein.

The presence of SV40 large T antigen and adenovirus E1A gene products in the HUVEC extended life cell line, as detected by ELISA, are shown in [t] Table [2] III. Briefly, the cell line grown in a 96 well tissue culture plate is fixed with glutaraldehyde and paraformaldehyde. The cells are then treated with antibodies to the corresponding oncogenes. Thereafter, the cells are washed and then treated with a secondary antibody linked with to β -galactosidase. The cells are washed and then treated with a substrate for β -galactosidase. The reaction develops a product which is then measured using a microplate reader.

To determine whether the extended life cell line has maintained the parental cell line phenotype may be determined by a number of ways. Extended life cells lines containing the oncogene are assessed by Northern analysis using a DNA probes encoding a cell-specific protein. The cell-specific DNA probe is labeled with ³²P-dCTP by nick translation pursuant, for example, to Rigby et al., J. Mol. Biol. 113: 237 (1977). Northern hybridization indicates that the extended life cell line is capable of transcribing the cell-specific protein.

Also, the maintenance of the parental phenotype in cells lines established according to the present invention can be

determined by a number of biochemical methods, such as ELISA and enzyme assays, that determine the presence or function of a protein specific to the parental cell line. An antibody recognizing a protein produced only [bythe] *by the* parental cell line can be used in an ELISA or immunofluorescence assay. Cell-specific markers are well known to those of skill in the art. For example, albumin is a marker for hepatocytes, insulin is a marker for pancreatic beta islet cells, factor VIII is a marker for endothelial cells, actin and myosin are markers for smooth muscle cells, and non-specific [esterass] *esterase* is a marker for brain microglial cells. In Table [II] III, the parental phenotype of the extended life endothelial cells produced by the present method of the invention was verified by several ELISAs to determine the expression of cell-specific endothelial markers. The parental phenotype of the monocyte/macrophage extended life cell lines produced by the present method was verified using a lysozyme enzyme assay to measure macrophage specific markers.

EXAMPLE 5

Human Umbilical Vein Endothelial Cells

Endothelial cells isolated from the human umbilical vein can only be cultured for a limited of passages, usually five to six. These cells were transfected with a combination of oncogenes, SV40 large T antigen and adenovirus E1A, or with another combination of genes. At least two oncogenes are needed to develop a truly transformed cell line. Ruley, H. E., et al., Nature 304: 602 (1983). For the instant invention, SV40 large T antigen combined with v-myc or ras or some an other oncogene can be used. When the gens encoding SV40 large T antigen is combined with either adenovirus E1A or E1B genes in the method of the instant invention, extended life cell lines may be produced from human umbilical vein endothelial cells. E1A or E1B or SV40 large T antigen alone did not give rise to established cell line with the high frequency obtained from using SV40 large T antigen in combination with E1A or E1B. Synthetic polypeptide complexed to DNA encoding either the SV40 large T antigen or polyoma large T antigen combined with the E2F1 transcription factor gens also produces extended life HUVEC cells lines.

Since the non-transfected primary cells normally grow in vitro only for a limited population doublings, cells that have taken up the oncogenes capable of generating extended life span cell lines were selected simply by repeated passage of the cells. When the transfected population of cells grows continuously, as compared to a control population of parental cells, it is reasonable to conclude that the oncogenes used are capable of generating extended life cells from a given cell type.

In HUVEC, for example, SV40 large T antigen and adenovirus E1A or E1B were effective in giving rise to a cell line. This cell line has now been growing in culture for 40 passages. In contrast, normal HUVECs stop growing by passage 7 or 8. Such cell lines arose with a high efficiency. It also is possible to generate cell lines using as few as a couple of hundred cells, grown either in a 24- or 48-well plate. These cells also have the same morphological appearance as the primary HUVEC and also display many of the biochemical properties characteristic of normal HUVEC.

Some of the properties that are characteristic of endothelial cells that were measured in the HUVEC extended life cell line are also listed in Table [2] III. These properties were also measured by ELISA using specific antibodies listed in the Table [2] III.

TABLE III

ELISA assay for the expression of ELAM-12, VCAM-1, ICAM-1, SV40 large T antigen and adenovirus EIA by extended life HUVEC line		
Antibody	O.D. _{490nm}	
	-IL-1	+IL-1
Control	0.071	0.069
Anti ELAM-1	0.212	1.016
Anti VCAM-1	0.146	0.520
Anti ICAM-1	0.422	1.524
Anti SV40 large T	0.618	—
Anti EIA	0.725	—

EXAMPLE 6

Human Cord Blood-Derived Monocyte Cell Line

Adherent cells from human cord blood cells were transfected with different combinations of oncogenes in suspension using the method of the present invention. The resulting cells are selected in Granulocyte-Macrophage Colony Stimulating Factor (G-CSF). Control cells did not grow in culture, whereas growing populations of monocytes were obtained with several combinations of oncogenes. One preferred combination of polyoma large T antigen and adenovirus E1B encoding DNA produced extended life monocyte cells lines with somewhat higher efficiency than other combinations. Another preferred combination of SV40 large T or polyoma large T antigen and the E2F1 transcription factor gene produces monocyte extended life cells with high efficiency. The monocyte extended life cells also display many of the properties of normal monocytes, which illustrates the utility of the present invention in generating cell lines of hematopoietic origin.

EXAMPLE 7

Extended Life Human Aortic Smooth Muscle Cells

The method of the instant invention has also been used to generate extended life cell lines using a specific combination of oncogenes. Human aortic smooth muscle cells were obtained from Clonetics Corporation (San Diego, Calif. U.S.A.) and transfected with several combination of oncogenes. The combination of polyoma large T antigen and E1B gave rise to a continuously growing population of smooth muscle cells. Another preferred combination of SV40 large T or polyoma large T antigen and the E2F1 transcription factor gene produces extended life human aortic smooth muscle cells with high efficiency. This cell line resembles the

early passage primary aortic smooth muscle cells morphologically. The extended life human aortic smooth muscle cells also express smooth cell actin and myosin well beyond passage 20.

EXAMPLE 8

Other Extended Life Cell Types

Primary cells from other species, such as rabbit and monkey, also have been used to generate cell lines. Transfection methods employed for primary cells from non-human species are similar to those used for human primary cells. When developing an extended life cell line from a new primary cell, several different combinations of available oncogenes should be tried. For example, at least five or six pairs of combinations of SV40 large T antigen, adenovirus E1A, adenovirus E1B, polyoma virus large T antigen or others available to those in the art. That combination of genes that gives rise to an extended life cell lines from a given primary cell type is determined as described in the above examples.

When the E2F1 transcription factor gene is complexed to synthetic polypeptide in combination with DNA encoding either the SV40 large T antigen or polyoma large T antigen, extended life cells lines can be produced from a variety of primary cell types, such as HUVEC, dermal microvascular endothelial cells, human aortic smooth muscle cells, and bone marrow monocyte/macrophage cells. Thus, the method of the present invention can identify a combination of oncogene DNAs that is highly efficient in producing extended life cells lines from the primary cells of various species. The present invention also comprehends a combination of an oncogene and an S-phase transcription factor gene which likewise is highly efficient in producing extended life cells lines from different types of primary cells.

EXAMPLE 9

Identification of Cell Type-Specific Transcriptional and Translational Regulatory Sequences

The present invention provides a screening system for identifying sequences that influence the expression of cloned genes in various primary cell types from different species. The instant invention can identify cell type specific transcription and translational regulatory sequences. The sequence in question typically will be cloned into a vector containing a reporter gene, such as chloramphenicol acetyl transferase or luciferase, and then transfected into various cell types using the method described herein. Expression of the reporter gene determines the tissue specificity of the regulatory sequence.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 56

<210> SEQ ID NO 1
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Yeast

<400> SEQUENCE: 1

Lys Ile Pro Ile Lys

-continued

1 5

<210> SEQ ID NO 2
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Yeast

<400> SEQUENCE: 2

Val Arg Ile Leu Glu Ser Trp Phe Ala Lys Asn Ile
 1 5 10

<210> SEQ ID NO 3
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: SV40

<400> SEQUENCE: 3

Pro Lys Lys Lys Arg Lys Val
 1 5

<210> SEQ ID NO 4
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Influenza virus

<400> SEQUENCE: 4

Ala Ala Phe Glu Asp Leu Arg Val Arg Ser
 1 5 10

<210> SEQ ID NO 5
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Yeast

<400> SEQUENCE: 5

Pro Arg Lys Arg
 1

<210> SEQ ID NO 6
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Polyoma virus

<400> SEQUENCE: 6

Val Ser Arg Lys Arg Pro Arg Pro Ala
 1 5

<210> SEQ ID NO 7
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: SV40

<400> SEQUENCE: 7

Ala Pro Thr Lys Arg Lys
 1 5

<210> SEQ ID NO 8
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Adenovirus

<400> SEQUENCE: 8

Lys Arg Pro Arg Pro
 1 5

-continued

<210> SEQ ID NO 9
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: SV40

<400> SEQUENCE: 9

Pro Asn Lys Lys Lys Arg Lys
 1 5

<210> SEQ ID NO 10
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Frog

<400> SEQUENCE: 10

Arg Pro Ala Ala Thr Lys Lys Ala Gly Gln Ala Lys Lys Lys Lys Leu
 1 5 10 15

Asp

<210> SEQ ID NO 11
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Rat

<400> SEQUENCE: 11

Lys Lys Lys Ile Lys
 1 5

<210> SEQ ID NO 12
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Monkey

<400> SEQUENCE: 12

Arg Val Thr Ile Arg Thr Val Arg Val Arg Arg Pro Pro Lys Gly Lys
 1 5 10 15

His Arg Lys

<210> SEQ ID NO 13
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Yeast

<400> SEQUENCE: 13

Gly Lys Lys Arg Ser Lys Ala
 1 5

<210> SEQ ID NO 14
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Chicken

<400> SEQUENCE: 14

Gly Lys Lys Arg Ser Lys Ala
 1 5

<210> SEQ ID NO 15
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Influenza

<400> SEQUENCE: 15

Asp Arg Leu Arg Arg

-continued

1 5

<210> SEQ ID NO 16
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Influenza

<400> SEQUENCE: 16

Pro Lys Gln Lys Arg Lys
 1 5

<210> SEQ ID NO 17
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Frog

<400> SEQUENCE: 17

Val Arg Lys Lys Arg Lys Thr
 1 5

<210> SEQ ID NO 18
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Frog

<400> SEQUENCE: 18

Ala Lys Lys Ser Lys Gln Glu
 1 5

<210> SEQ ID NO 19
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 19

Pro Ala Ala Lys Arg Val Lys Leu Asp
 1 5

<210> SEQ ID NO 20
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 20

Arg Gln Arg Arg Asn Glu Leu Lys Ser Phe
 1 5 10

<210> SEQ ID NO 21
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 21

Thr Lys Lys Arg Lys Leu Glu
 1 5

<210> SEQ ID NO 22
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: HTLV-1

<400> SEQUENCE: 22

Pro Lys Thr Arg Arg Arg Pro
 1 5

-continued

<210> SEQ ID NO 23
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: HTLV-1

<400> SEQUENCE: 23

Ser Gln Arg Lys Arg Pro Pro
 1 5

<210> SEQ ID NO 24
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Adenovirus

<400> SEQUENCE: 24

Arg Leu Pro Val Arg Arg Arg Arg Arg Val Pro
 1 5 10

<210> SEQ ID NO 25
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: HIV-1

<400> SEQUENCE: 25

Gly Arg Lys Lys Arg
 1 5

<210> SEQ ID NO 26
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Frog

<400> SEQUENCE: 26

Val Arg Thr Thr Lys Gly Lys Arg Lys Arg Ile Asp Val
 1 5 10

<210> SEQ ID NO 27
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Rabbit

<400> SEQUENCE: 27

Arg Lys Phe Lys Lys
 1 5

<210> SEQ ID NO 28
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: HIV-1

<400> SEQUENCE: 28

Arg Arg Asn Arg Arg Arg Arg Trp
 1 5

<210> SEQ ID NO 29
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 29

Pro Arg Ser Gly Lys Lys Arg Lys Arg Lys Arg Leu Lys Pro Thr
 1 5 10 15

<210> SEQ ID NO 30

-continued

<211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Mouse

<400> SEQUENCE: 30

Ser Ala Leu Ile Lys Lys Lys Lys Lys Met Ala Pro
 1 5 10

<210> SEQ ID NO 31
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Adenovirus

<400> SEQUENCE: 31

Pro Pro Lys Lys Arg
 1 5

<210> SEQ ID NO 32
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Adenovirus

<400> SEQUENCE: 32

Pro Lys Lys Lys Lys Lys
 1 5

<210> SEQ ID NO 33
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Chicken

<400> SEQUENCE: 33

Ser Lys Arg Val Ala Lys Arg Lys Leu
 1 5

<210> SEQ ID NO 34
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 34

Pro Leu Leu Lys Lys Ile Ile Gln
 1 5

<210> SEQ ID NO 35
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 35

Pro Pro Gln Lys Lys Ile Lys Ser
 1 5

<210> SEQ ID NO 36
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 36

Pro Gln Pro Lys Lys Lys Pro
 1 5

<210> SEQ ID NO 37
 <211> LENGTH: 18
 <212> TYPE: PRT

-continued

<213> ORGANISM: Human

<400> SEQUENCE: 37

Phe	Lys	Arg	Lys	His	Lys	Lys	Asp	Ile	Ser	Gln	Asn	Lys	Arg	Ala	Val
1				5					10					15	

Arg Arg

<210> SEQ ID NO 38

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Hepatitis B virus

<400> SEQUENCE: 38

Ser	Lys	Cys	Leu	Gly	Trp	Leu	Trp	Gly
1				5				

<210> SEQ ID NO 39

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Chicken

<400> SEQUENCE: 39

Gly	Lys	Arg	Lys	Asn	Lys	Pro	Lys
1				5			

<210> SEQ ID NO 40

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Yeast

<400> SEQUENCE: 40

Lys	Thr	Arg	Lys	His	Arg	Gly
1				5		

<210> SEQ ID NO 41

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Yeast

<400> SEQUENCE: 41

Lys	His	Arg	Lys	His	Pro	Gly
1				5		

<210> SEQ ID NO 42

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Tobacco

<400> SEQUENCE: 42

Arg	Arg	Leu	Ala	Gln	Asn	Arg	Glu	Ala	Ala	Arg	Lys	Ser	Arg	Leu	Arg
1				5					10					15	

Lys Lys

<210> SEQ ID NO 43

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Tobacco

<400> SEQUENCE: 43

Lys	Lys	Arg	Ala	Arg	Leu	Val	Arg	Asn	Arg	Glu	Ser	Ala	Gln	Leu	Ser
1				5					10					15	

<210> SEQ ID NO 44

-continued

<211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Tobacco

<400> SEQUENCE: 44

Arg Gln Arg Lys Lys
 1 5

<210> SEQ ID NO 45
 <211> LENGTH: 18
 <212> TYPE: PRT
 <213> ORGANISM: Maize

<400> SEQUENCE: 45

Arg Lys Arg Lys Glu Ser Asn Arg Glu Ser Ala Arg Arg Ser Arg Tyr
 1 5 10 15

Arg Lys

<210> SEQ ID NO 46
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Polyvirus

<400> SEQUENCE: 46

Lys Lys Asn Gln Lys His Lys Leu Lys Met Lys Arg Lys
 1 5 10

<210> SEQ ID NO 47
 <211> LENGTH: 18
 <212> TYPE: PRT
 <213> ORGANISM: Agrobacterium

<400> SEQUENCE: 47

Lys Arg Pro Arg Glu Asp Asp Asp Gly Glu Pro Ser Glu Arg Lys Arg
 1 5 10 15

Glu Arg

<210> SEQ ID NO 48
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Agrobacterium

<400> SEQUENCE: 48

Lys Leu Arg Pro Glu Asp Arg Tyr Ile Gln Thr Glu Lys Tyr Gly Arg
 1 5 10 15

Arg

<210> SEQ ID NO 49
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Agrobacterium

<400> SEQUENCE: 49

Lys Thr Lys Tyr Gly Ser Asp Thr Glu Ile Lys Leu Leu Ser Lys
 1 5 10 15

<210> SEQ ID NO 50
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Maize

<400> SEQUENCE: 50

Met Glu Glu Ala Val Thr Met Ala Pro Ala Ala Val Ser Ser Ala Val

-continued

1	5	10	15
---	---	----	----

Val Gly Asp Pro
 20

<210> SEQ ID NO 51
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Maize

<400> SEQUENCE: 51

Met Tyr Asn Ala Ile Leu Arg Arg Lys Leu Glu Glu Asp Leu Glu			
1	5	10	15

<210> SEQ ID NO 52
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Maize

<400> SEQUENCE: 52

Gly Asp Arg Arg Ala Ala Pro Ala Arg Pro			
1	5	10	15

<210> SEQ ID NO 53
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Maize

<400> SEQUENCE: 53

Met Ser Glu Arg Lys Arg Arg Glu Lys Leu			
1	5	10	15

<210> SEQ ID NO 54
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Maize

<400> SEQUENCE: 54

Met Ile Ser Glu Ala Leu Arg Lys Ala Ile Gly Lys Arg			
1	5	10	15

<210> SEQ ID NO 55
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Xenopus nucleoplasmin

<400> SEQUENCE: 55

Lys Arg Pro Ala Ala Thr Lys Lys Ala Gly Gln Ala Lys Lys Lys Lys			
1	5	10	15

<210> SEQ ID NO 56
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide with nuclear localization
 signal domain and DNA binding domain.

<400> SEQUENCE: 56

Pro Lys Lys Lys Arg Lys Val Ser Gly Gly Gly Gly Lys Lys Lys			
1	5	10	15

Lys Lys Lys Lys Lys Lys Lys Lys Lys			
20	25	30	35

What is claimed is:

1. A transfection vector comprising a synthetic polypeptide linked electrostatically to a DNA structural sequence, forming a polypeptide-DNA complex, wherein said polypeptide is comprised of (A) a polymeric chain of basic amino acid residues, (B) a nuclear localization signal (NLS) peptide and (C) a hinge region of neutral amino acids that connects said polymeric chain and said NLS peptide.

2. The vector of claim 1, wherein said polymeric chain is comprised of between 10 and 50 residues.

3. The vector of claim 1, wherein said basic amino acid residues are selected from the group consisting of lysine, arginine and ornithine.

4. The vector of claim 1, wherein said NLS peptide is selected from the group consisting of a Simian Virus 40 (SV40) large T antigen nuclear localization signal sequence, a polyoma large T antigen nuclear localization signal sequence, an adenovirus E1a nuclear localization signal sequence, and an adenovirus E1b nuclear localization signal sequence.

5. The vector of claim 1, wherein said hinge region is comprised of between 6 and 50 amino acid residues.

6. The vector of claim 1, wherein said neutral amino acids are selected from the group consisting of glycine, alanine, leucine and isoleucine.

7. The vector of claim 1, wherein said NLS peptide is located at the amino terminus of said polypeptide and said polymeric basic amino acid chain is located at the carboxyl terminus.

8. The transfection vector of claim 1, further comprising (D) a cell type-specific ligand molecule.

9. The transfection vector of claim 1, wherein said DNA structural sequence comprises (a) a segment coding for SV40 large T antigen or polyoma large T antigen and (b) a transcription factor gene.

10. A vector according to claim 1, wherein said DNA structural sequence comprises an oncogene.

11. A vector according to claim 10, wherein said oncogene is selected from the group consisting of SV40 large T antigen, polyoma large T antigen, adenovirus E1A, adenovirus E1B, v-fms, BC12, myc, and ras.

12. A vector according to claim 1, wherein said DNA structural sequence comprises a DNA sequence selected from the group consisting of a dihydrofolate reductase [gens] gene (DHFR), a thymidine kinase [gens] gene, a thymidylate synthetase [gene] gene, a DRTF1/E2F transcription factor-encoding DNA sequence, and an E2F transcription factor-encoding DNA sequence.

13. A process for producing a transformed mammalian cell line, comprising the step of transfecting a mammalian cell with a vector according to claim 1, wherein said DNA structural sequence comprises a DNA sequence selected from the group consisting of a dihydrofolate reductase gene (DHFR), a thymidine kinase gene, a thymidylate synthetase [gene] gene, a DRTF1/E2F transcription factor-encoding DNA sequence, and an E2F transcription factor-encoding DNA sequence.

14. A process for producing a transformed mammalian cell line, comprising the step of transfecting a mammalian cell with a vector according to claim 1, wherein said DNA structural sequence comprises an oncogene.

* * * * *



US00RE39220F1

(12) **EX PARTE REEXAMINATION CERTIFICATE (6350th)**
United States Patent
Gopal

(10) **Number:** **US RE39,220 F1**
(45) **Certificate Issued:** ***Aug. 12, 2008**

(54) **PEPTIDE-MEDIATED GENE TRANSFER**

(52) **U.S. Cl.** **435/455**; 435/320.1; 435/467;
530/300; 530/50

(75) **Inventor:** **T. Venkat Gopal**, North Potomac, MD
(US)

(58) **Field of Classification Search** None
See application file for complete search history.

(73) **Assignee:** **Genetic Applications, LLC**, San Diego,
CA (US)

(56) **References Cited**

Reexamination Request:

No. 90/008,557, Mar. 23, 2007

U.S. PATENT DOCUMENTS

5,342,921 A 8/1994 Cousens et al.
5,955,365 A 9/1999 Szoka, Jr. et al.

Reexamination Certificate for:

Patent No.: **1,039,220**
Issued: **Aug. 1, 2006**
Appl. No.: **09/404,979**
Filed: **Sep. 22, 1999**

OTHER PUBLICATIONS

Alberts, *Molecular Biology of the Cell*, c1994, pp. 56–57.*
Lanford, et al., “Effect of Basic and Nonbasic Amino Acid
Substitutions on Transport Induced by Simian Virus 40
T–Antigen Synthetic Peptide Nuclear Transport Signals,”
Molecular and Cellular Biology. Jul. 1988, vol. 8, No. 7, pp.
2722–2729.

(*) **Notice:** This patent is subject to a terminal dis-
claimer.

* cited by examiner

Related U.S. Patent Documents

Reissue of:

(64) **Patent No.:** **5,670,347**
Issued: **Sep. 23, 1997**
Appl. No.: **08/240,514**
Filed: **May 11, 1994**

Primary Examiner—Sharon L Turner

(57) **ABSTRACT**

(51) **Int. Cl.**
C12N 15/87 (2006.01)
C12N 15/63 (2006.01)
C12N 15/85 (2006.01)
C07K 14/00 (2006.01)

A synthetic polypeptide that allows for highly efficient trans-
fer of DNA into eukaryotic cells, including, for example,
non-dividing cells such as human peripheral blood mono-
cytes and macrophages.

1
EX PARTE
REEXAMINATION CERTIFICATE
ISSUED UNDER 35 U.S.C. 307

NO AMENDMENTS HAVE BEEN MADE TO
THE PATENT

2
AS A RESULT OF REEXAMINATION, IT HAS
BEEN DETERMINED THAT:

The patentability of claims 1-14 is confirmed.

5

* * * * *