



US00RE39300E

(19) **United States**
(12) **Reissued Patent**
Mayer et al.

(10) **Patent Number:** **US RE39,300 E**
(45) **Date of Reissued Patent:** **Sep. 19, 2006**

(54) **INHIBITING THE DEVELOPMENT OF TOLERANCE TO AND/OR DEPENDENCE ON AN ADDICTIVE SUBSTANCE**

(75) Inventors: **David J. Mayer**, Richmond, VA (US);
Jianren Mao, Boston, MA (US)

(73) Assignee: **Virginia Commonwealth University Medical College of Virginia**,
Richmond, VA (US)

4,126,684 A 11/1978 Robson et al. 424/254
4,316,888 A 2/1982 Nelson 424/127
4,362,870 A 12/1982 Portoghese 542/403
4,416,871 A 11/1983 Walter et al. 424/177
4,446,140 A 5/1984 Nelson 424/260
4,464,378 A 8/1984 Hussain 424/260
4,476,141 A 10/1984 Cormier 424/321
4,602,909 A 7/1986 Csillik 604/20
4,758,559 A 7/1988 Wasley 514/211

(Continued)

(21) Appl. No.: **09/880,881**

(22) Filed: **Jun. 15, 2001**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **5,556,838**
Issued: **Sep. 17, 1996**
Appl. No.: **08/245,970**
Filed: **May 19, 1994**

U.S. Applications:

(63) Continuation of application No. 08/043,280, filed on Apr. 6, 1993, now Pat. No. 5,321,012, which is a continuation-in-part of application No. 08/010,583, filed on Jan. 28, 1993, now abandoned.

(51) **Int. Cl.**
A61K 31/70 (2006.01)
A61K 31/54 (2006.01)
A61K 31/44 (2006.01)
A61K 31/445 (2006.01)

(52) **U.S. Cl.** **514/25**; 514/216; 514/223.5;
514/224.5; 514/225.5; 514/231.2; 514/282;
514/304; 514/305; 514/306; 514/307; 514/812

(58) **Field of Classification Search** 514/289,
514/647, 216, 25, 223.5, 224.5, 225.5, 231.2,
514/282, 304-307, 812; 536/55, 55.1
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,770,569 A 11/1956 Fromherz et al. 167/65

OTHER PUBLICATIONS

Pharmacol. Biochem. Behav. vol. 35, No. 4 (1990), pp. 829-832.*

Pharmacol. Biochem. Behav. vol. 43, No. 2 (1992), pp. 487-490.*

Disch. Apoth.-Zig. vol. 119, No. 21 (1979), p. 821, "Opiat-Rezeptoren und Endorphine".*

Truillo et al.; "Inhibition of Morphine . . . MIC801"; Science; vol. 251, pp. 84-87.*

Marek et al; "Brain Research, Delayed application of MIC801 . . . in rats."; vol. 558, pp. 163-165.*

Adams et al., "Emergency Intravenous Sedation of the Delirious, Medically Ill Patient," *J. Clin. Psychiatry*, 49:12 Suppl., pp. 23-27 (1988).

(Continued)


Primary Examiner—Shengjun Wang

(74) *Attorney, Agent, or Firm*—Morgan & Finnegan, LLP

(57) **ABSTRACT**

Nontoxic substances that block the N-methyl-D-aspartate (NMDA) receptor, e.g., a morphinan such as dextromethorphan or dextrorphan, or that block a major intracellular consequence of NMDA-receptor activation, e.g., a ganglioside such as GM₁ or GT_{1b}, a phenothiazine such as trifluoperazine or a naphthalenesulfonamide such as N-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide, inhibit the development of tolerance to and/or dependence on addictive drugs, e.g., narcotic analgesics such as morphine, codeine, etc.

15 Claims, 10 Drawing Sheets



U.S. PATENT DOCUMENTS

4,769,372 A	9/1988	Kreek	514/282
4,785,000 A	11/1988	Kreek et al.	514/282
4,788,055 A	11/1988	Fischer et al.	424/79
4,806,543 A	2/1989	Choi	514/464
4,816,264 A	3/1989	Phillips	424/468
4,816,450 A	3/1989	Bell et al.	514/25
4,828,836 A	5/1989	Elger et al.	424/419
4,834,965 A	5/1989	Martani et al.	424/419
4,834,985 A	5/1989	Elger et al.	424/488
4,876,276 A	10/1989	Mechoulam et al.	514/454
4,888,347 A	12/1989	Woodruff et al.	514/289
4,906,779 A	3/1990	Weber et al.	564/238
4,912,114 A	3/1990	Revesz	514/282
4,919,916 A	4/1990	Golwyn	424/10
4,924,008 A	5/1990	Abou-Gharbia et al.	849/75
4,937,232 A	6/1990	Bell et al.	514/26
4,942,182 A	7/1990	Weiss et al.	514/812
4,959,493 A	9/1990	Ohfume et al.	562/506
4,975,430 A	12/1990	Jahr et al.	514/255
4,990,519 A	2/1991	Jones et al.	514/314
4,994,446 A	2/1991	Sokolovsky et al.	514/75
4,994,467 A	2/1991	Zimmerman	514/284
4,996,047 A	2/1991	Kelleher	424/79
5,006,510 A	4/1991	Ellis	514/16
5,013,540 A	5/1991	Redburn	424/10
5,023,239 A	6/1991	Ogura et al.	514/26
5,028,611 A	7/1991	Halikas	514/277
5,028,707 A	7/1991	Nichols et al.	546/156
5,034,395 A	7/1991	Tamada et al.	514/277
5,034,400 A	7/1991	Olney	514/315
5,041,446 A	8/1991	Silvestrini	514/255
5,047,229 A	9/1991	Nahas et al.	424/10
5,051,426 A	9/1991	Parnell	514/419
5,051,442 A	9/1991	Salituro et al.	514/381
5,055,481 A	10/1991	Inukai et al.	514/282
5,057,519 A	10/1991	Suberg	514/114
5,068,228 A	11/1991	Kohler	424/497
5,071,646 A	12/1991	Malkowska	514/464
5,075,341 A	12/1991	Mendelson et al.	514/282
5,091,391 A	2/1992	Aizenman et al.	514/311
5,093,129 A	3/1992	Horovitz et al.	424/451
5,093,525 A	3/1992	Weber et al.	564/338
5,095,009 A	3/1992	Whitten et al.	514/85
5,106,847 A	4/1992	Salituro et al.	514/232.5
5,109,017 A	4/1992	Schmiesing et al.	514/438
5,118,675 A	6/1992	Jirovsky et al.	514/80
5,124,319 A	6/1992	Baudy et al.	514/80
5,124,340 A	6/1992	Jaffe et al.	514/356
5,126,330 A	6/1992	Ogura et al.	514/26
5,133,974 A	7/1992	Paradissis	424/480
5,137,889 A	8/1992	Tamada et al.	514/252
5,137,919 A	8/1992	Igarashi et al.	514/642
5,141,957 A	8/1992	Jiang et al.	514/510
5,145,842 A	9/1992	Driedger et al.	514/63
5,151,360 A	9/1992	Handa et al.	435/240.2
5,166,207 A	11/1992	Smith	514/270
5,171,752 A	12/1992	Caldirola et al.	514/648
5,183,807 A	2/1993	della Valle et al.	514/25
5,185,329 A	2/1993	Gawin et al.	514/159
5,190,925 A	3/1993	dalla Valle et al.	424/25
5,321,012 A	6/1994	Mayer et al.	564/25
5,352,683 A	10/1994	Mayer et al.	514/289
5,441,982 A	8/1995	Itzhak	514/55
5,654,281 A	8/1997	Mayer et al.	514/25

OTHER PUBLICATIONS

- Ben-Eliyahu et al., "The NMDA receptor antagonist MK-801 prevents long-lasting non-associative morphine tolerance in the rat," *Brain Research*, vol. 575, pp. 304-308 (1991).
- Bristow et al., "Subcutaneous ketamine analgesia: postoperative analgesia using subcutaneous infusions of ketamine and morphine," *Annals of the Royal College of Surgeons of England*, vol. 71 (1989).
- Chapman et al., "The Combination of NMDA Antagonism and Morphine Produces Profound Antinociception in the Rat Dorsal Horn," *Brain Research*, vol. 573, pp. 321-323 (1991).
- Dickenson et al., "Dextromethorphan and Levorphanol on Dorsal Horn Nociceptive Neurons in the Rat," *Neuropharmacology*, vol. 30, No. 12A, pp. 1303-1308 (1991).
- Haley et al., "Evidence for spinal N-Methyl-D-aspartate receptor involvement in prolonged chemical nociception in the rat," *Brain Research*, vol. 518, pp. 218-226 (1989).
- Hayes et al., "Pretreatment with gangliosides reduces abnormal nociceptive responses associated with a rodent peripheral mononeuropathy," *Pain*, vol. 48, pp. 391-396 (1992).
- Itzhak et al., "Sensitization to the Toxic Effects of Cocaine in Mice is Associated with the Regulation of N-Methyl-D-Aspartate Receptors in the Cortex," *The Journal of Pharmacology and Experimental Therapeutics*, vol. 262, No. 2, pp. 464-470 (1992).
- Kamei et al., "Subsensitivity to the Cough-Depressant Effects of Opioid and Nonopioid Antitussives in Morphine-Dependent Rats: Relationship to Central Serotonin Function," *Pharmacology Biochemistry & Behavior*, vol. 34, pp. 595-598 (1988).
- Kanamaru et al., "Ketamine infusion for Control of Pain in Patients with Advanced Cancer," pp. 1368-1371 (1990) (see English translation of abstract).
- Koyunchoglu et al., "The treatment of heroin addicts with dextromethorphan: A double-blind comparison of dextromethorphan with Chlorpromazine," *International Journal of Clinical Pharmacology, Therapy and Toxicology*, vol. 28, No. 4 pp. 147-152 (1990).
- Koyuncuoglu et al., "Previous Chronic Blockade of NMDA Receptors Intensifies Morphine Dependence in Rats," *Pharmacology Biochemistry & Behavior*, vol. 39, pp. 575-579 (1991).
- Koyuncuoglu et al., "Suppression by Ketamine and Dextromethorphan of Precipitated Abstinence Syndrome in Rats," *Pharmacology Biochemistry and Behavior*, vol. 35, No. 4, pp. 829-832 (1990).
- Koyuncuoglu et al., "Effects of MK 801 on Morphine Physical Dependence: Attenuation and Intensification," *Pharmacology Biochemistry & Behavior*, vol. 43, pp. 487-490 (1992).
- Mao et al., "Intrathecal MK-801 and local nerve anesthesia synergistically reduce nociceptive behaviors in rats with experimental peripheral mononeuropathy," *Brain Research*, vol. 576, pp. 254-262 (1992).
- Mao et al., "Post-injury treatment with GM1 ganglioside reduces nociceptive behaviors and spinal cord metabolic activity in rats with experimental peripheral mononeuropathy," *Brain Research*, vol. 584, pp. 18-27 (1992).
- Mao et al., "Intrathecal GM1 ganglioside and local nerve anesthesia reduce nociceptive behaviors in rats with experimental peripheral mononeuropathy," *Brain Research*, vol. 584, pp. 28-35 (1992).

Mao et al., "Pain-related increases in spinal cord membrane-bound protein kinase C following peripheral nerve injury," *Brain Research*, vol. 588, pp. 144-149 (1992).

Marek et al., "Excitatory amino acid antagonists (kynurenic acid and MK-801) attenuate the development of morphine tolerance in the rat", *Brain Research*, vol. 547, pp. 77-81 (1991).

Marek et al., "Delayed application of MK-801 attenuates development of morphine tolerance in rats," *Brain Research*, vol. 558, pp. 163-165 (1991).

Mankowitz et al., "Epidural Ketamine," *SA Medical Journal*, vol. 20, pp. 441-442 (1982).

Oshima et al., "Continuous subcutaneous injection of ketamine for cancer pain," *Can. J. Anaesth.*, vol. 37:3, pp. 385-392, (1994).

Pharma Projects, 1561 (May 1994).

Sadee, "Opiat-Rezeptoren und Endorphine," *Deutsche Apotheker-Zeitung*, vol. 119, p. 821 (1979).

Tanganelli et al., "Glutamate antagonists prevent morphine withdrawal in mice and guinea pigs," *Neuroscience-Letter.*, vol. 122, pp. 270-272, 1991.

Trujillo et al., "Inhibition of Morphine Tolerance and Dependence by the NMDA Receptor Antagonist MK-801," *Science*, vol. 251, pp. 85-87 (1991).

Ventafriidda et al., "Studies on the effects of antidepressant drugs on the antinociceptive action of morphine and on plasma morphine in rat and man," *Pain*, vol. 43, pp. 155-162 (1990).

* cited by examiner




FIG.1




FIG.2




FIG.3




FIG.4




FIG.5




FIG. 6




FIG.7




FIG. 8




FIG. 9




FIG.10

**INHIBITING THE DEVELOPMENT OF
TOLERANCE TO AND/OR DEPENDENCE ON
AN ADDICTIVE SUBSTANCE**

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.

CROSS REFERENCE TO RELATED
APPLICATION

This is a continuation of application Ser. No. 08/043,280 filed Apr. 6, 1993, now U.S. Pat. No. 5,321,012, which is a continuation-in-part of U.S. patent application Ser. No. 08/010,583, filed Jan. 28, 1993, abandoned.

BACKGROUND OF THE INVENTION

This invention relates to a composition containing an addictive substance and a component which inhibits the development of tolerance to and/or dependence on the addictive substance. More particularly, the invention relates to a composition containing an addictive substance such as morphine or codeine and at least one nontoxic substance that blocks the N-methyl-D-aspartate (NMDA) receptor, e.g., a morphinan such as dextromethorphan or dextrorphan, or that blocks at least one major intracellular consequence of NMDA receptor activation, e.g., a ganglioside such as ganglioside GM₁ or GT_{1b}, a phenothiazine such as trifluoperazine or a naphthalenesulfonamide such as N-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide.

Morphine is a rapid and effective drug for the treatment of severe pain but its long term administration has been limited due to its negative side effects, principally tolerance and dependence, which develop rapidly after administration. In an effort to make morphine of greater use in the treatment of pain, it has been combined with a variety of substances intended to inhibit one or more of its undesirable side effects. U.S. Pat. No. 2,770,569 describes the combination of morphine with the compound levo-d-hydroxy-N-allylmorphinan which is said to suppress or eliminate such undesirable side reactions of morphine as depression, nausea and vomiting. U.S. Pat. No. 4,126,684 discloses reducing either the addiction liability of an addictive substance such as a narcotic analgesic or a barbiturate or the withdrawal symptoms caused by deprivation of such a substance in an addicted subject by administering the addictive substance, e.g., morphine, with a 4-amino-3-p-halophenylbutyric acid. U.S. Pat. No. 4,415,871 describes the prevention of treatment tolerance and physical dependence in chronic morphine treatment by combining the morphine with any of the specific dipeptides indicated therein. U.S. Pat. No. 5,041,446 discloses inhibiting the development of tolerance to morphine by combining the morphine with dapiprazole. U.S. Pat. No. 5,057,519 achieves a reduction in morphine tolerance by combining the morphine with a benzamide antagonist for a subtype of the serotonin receptor, 5-HT₃. Trujillo et al., "Inhibition of morphine tolerance and dependence by the NMDA receptor antagonist MK-801", *Science*, 251 (4989), pp. 85-87, Jan. 4, 1991; Tanganelli et al., "Glutamate antagonists prevent morphine withdrawal in mice and guinea pigs", *Neuroscience Letters*, 122(2), pp. 270-272, Jan. 28, 1991; Marek et al., "Excitatory amino acid antagonists (kynurenic acid and MK-801) attenuate the development of morphine tolerance in the rat", *Brain Research*, 547(1), pp. 77-81, Apr. 26, 1991; and, Marek et al., "Delayed application of MK-801 attenuates develop-

ment of morphine tolerance in rats, *Brain Research*, 558(1), pp. 163-165, Aug. 30, 1991 discuss the role of MK-801 (the compound 5-methyl-10,11-dihydro-SH-dibenzo[a,d]cyclohepten-5,10-imine), an NMDA receptor antagonist or blocker, in reducing morphine dependence in laboratory animals. However, MK-801 has been found to be toxic and is therefore unsuitable for pharmaceutical use.

SUMMARY OF THE INVENTION

In accordance with the present invention, a composition is provided which comprises an addictive substance and at least one nontoxic substance that blocks the N-methyl-D-aspartate receptor or at least one major intracellular consequence of N-methyl-D-aspartate receptor activation.

Further in accordance with the present invention, a method of inhibiting the development of tolerance to and/or dependence on an addictive substance administered to a mammal which is liable to addiction thereto is provided which comprises administering the addictive substance to the mammal before, with or following administration to the mammal of a tolerance-reducing and/or dependence-reducing amount of at least one non-toxic substance that blocks the N-methyl-D-aspartate receptor or at least one major intracellular consequence of N-methyl-D-aspartate receptor activation.

Still further in accordance with this invention, a method of alleviating withdrawal symptoms in a mammal addicted to an addictive substance is provided which comprises administering to the addicted mammal the addictive substance before, with or following administration to the mammal of a dependence-reducing amount of at least one nontoxic substance that blocks the N-methyl-D-aspartate receptor or the intracellular consequences of N-methyl-D-aspartate receptor activation thereby alleviating withdrawal symptoms when the addictive substance is withdrawn from the mammal.

The term "nontoxic" as used herein shall be understood in a relative sense and is intended to designate any substance that has been approved by the United States Food and Drug Administration ("FDA") for administration to humans or, in keeping with established criteria, is susceptible to approval by the FDA for administration to humans.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1-10 are graphical representations of experimental data demonstrating the effectiveness of specific nontoxic substances that block the N-methyl-D-aspartate receptor or a major consequence of N-methyl-D-aspartate receptor activation for inhibiting morphine tolerance and dependence in rats.

DESCRIPTION OF THE PREFERRED
EMBODIMENTS

A particularly important category of addictive substances with which the present invention is concerned are the narcotic analgesics, e.g., opiates, opiate derivatives, opioids and their pharmaceutically acceptable salts. Specific examples of narcotic analgesics include alfentanil, alphaprodine, anileridine, bezitramide, codeine, dihydrocodeine, diphenoxylate, ethylmorphine, fentanyl, heroin, hydrocodone, hydromorphone, isomethadone, levomethorphan, levorphanol, metazocine, methadone, metopon, morphine, opium extracts, opium fluid extracts, powdered opium, granulated opium, raw opium, tincture of opium, oxycodone, oxymorphone, pethidine, phenazocine,

piminodine, racemethorphan, racemorphan, thebaine and pharmaceutically acceptable salts thereof. For a detailed discussion of these and other narcotic analgesics, reference may be made to Jaffe et al., "Opioid Analgesics and Antagonists" in "Goodman and Gillman's Pharmacological Basis of Therapeutics", Goodman et al., eds. 7th ed., 1985, Mac-Millan and Company, New York pp. 491-531.

Other addictive substances that can be utilized herein include acetorphan, acetyldihydrocodeine, acetylmethadol, allylprodine, alphracetylmethadol, alphameprodine, alphamethadol, benzethidine, benzylmorphine, betacetylmethadol, betameprodine, betamethadol, betaprodine, clonitazene, cocaine, codeine, methylbromide, codeine-N-oxide, cyprenorphine, desomorphine, dextromoramide, diampromide, diethylthiambutene, dihydromorphine, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, drotebanol, ethanol, ethylmethylthiambutene, etonitazene, etorphine, etoxeridine, furethidine, hydromorphanol, hydroxypethidine, ketobemidone, levomoramide, levophenacetylmorphan, methyl-desorphine, methyldihydromorphine, morpheridine, morphine methylpromide, morphine methylsulfonate, morphine-N-oxide, myrophine, nicocodeine, nicomorphine, nicotine, noracetylmethadol, norlevorphanol, normethadone, normorphine, norpiperanone, phenadoxone, phenampromide, phenomorphitan, phenoperidine, piritramide, pholcodine, proheptazone, properidine, propiran, racemoramide, thebacon, trimeperidine and the pharmaceutically acceptable salts thereof.

Still other addictive substances that can be utilized in the practice of the invention include the sedatives and hypnotics, e.g., benzodiazepines such as chlordiazepoxide, chlorazepate, diazepam, flurazepam, halazepam, ketazolam, borazepam, oxazepam, prazepam, temazepam, triazolam and the pharmaceutically acceptable salts thereof, barbiturates such as amobarbital, ambobarbital, barbital, butabartital, mephobarbital, methohexital, pentobarbital, phenobarbital, secobarbital, talbutal, thiamylal and thiopental and the pharmaceutically acceptable salts thereof and other sedatives and hypnotics such as chloral hydrate, meprobamate, methaqualone, methyprylon and the pharmaceutically acceptable salts thereof.

By way of inhibiting the development of tolerance to and/or dependence on any of the foregoing and similarly addictive substances, the addictive substance is administered before, with or following the administration of at least one nontoxic substance that blocks the N-methyl-D-aspartate (NMDA) receptor or the intracellular consequences of N-methyl-D-aspartate receptor activation. Activation of the NMDA receptor, a subtype of excitatory amino acid receptors, induces a number of changes in the functional activity of nerve cells, and in particular, their capacity for excitability or inhibition in the presence of an addictive substance, via an increase in intracellular Ca^{++} concentration. The major consequences of NMDA receptor activation include the following sequences, or cascades, of events occurring within nerve cells:

- a) translocation and activation of protein kinases such as protein kinase C \rightarrow phosphorylation of substrate proteins such as cytosolic enzymes, channel proteins, receptor proteins, etc. \rightarrow changes in functional activity;
- b) initiation of early gene (c-fos, c-jun, zif-268, etc.) expression by either increased intracellular Ca^{++} or Ca^{++} -activated protein kinases \rightarrow expression of functional genes responsible for production of cellular

enzymes (such as protein kinases), receptor proteins (such as the NMDA receptor), ion channel proteins (such as K^+ , Na^+ , Ca^{++} channels), neuropeptides (such as dynorphin), etc. \rightarrow changes in functional activity;

- c) Ca^{++} /calmodulin (or other Ca^{++} binding proteins) induced activation of enzymes and other cellular components \rightarrow activation of Ca^{++} /calmodulin-protein kinase systems such as Ca^{++} /calmodulin kinase II \rightarrow autophosphorylation of enzymes (e.g., Ca^{++} /calmodulin kinase II) or other functional proteins \rightarrow changes in functional activity;
- d) Ca^{++} /calmodulin induced activation of constitutive nitric oxide synthase as well as induction of inducible nitric oxide synthase \rightarrow production of nitric oxide \rightarrow i) production of cyclic guanosine monophosphate via activation of guanosine cyclase resulting in activation of protein kinases and early gene expression; ii) direct protein modification such as enzymes, receptor and/or channel proteins; iii) lipid membrane modification and/or nucleic acid modification via scavenge of free radicals; iv) induction of neurotoxicity at higher nitric oxide levels; v) retrograde actions in adjacent neurons or glial cells such as facilitation of glutamate release/NMDA receptor activation and/or inhibition of post-synaptic NMDA receptors \rightarrow changes in functional activity;
- e) interactions with the cyclic adenosine monophosphate/protein kinase A system, the phospholipase C-inositol triphosphate- Ca^{++} -diacylglycerol-protein kinase system, the phospholipase A2-arachidonic acid/prostanoids/leukotrienes system \rightarrow changes in functional activity induced by second messenger systems other than NMDA receptor/ Ca^{++} / Ca^{++} -calmodulin/protein kinase systems; and,
- f) interactions with other excitatory amino acid receptor subtypes including non-NMDA receptors and metabotropic receptors as well as intracellular events subsequent to the activation of these excitatory amino acid receptor subtypes \rightarrow changes in functional activity induced by the non-NMDA and metabotropic receptor activation.

A substance that blocks the NMDA receptor will effectively prevent all of the foregoing major intracellular sequences of events from taking place. However, even with activation of the NMDA receptor, it is still possible to inhibit the development of tolerance to and/or dependence on an addictive substance by combining the addictive substance with a substance that blocks at least one of the foregoing major intracellular sequences of events. Thus, e.g., a substance that interferes with translocation and activation of protein kinase C or with calmodulin induced activation of constitutive nitric oxide synthase as well as induction of inducible nitric oxide synthase is also useful for the practice of this invention.

Among the nontoxic substances that block the NMDA receptor and as such are useful in the practice of the present invention are morphinans such as dextromethorphan ((+)-3-hydroxy-N-methylmorphinan) and dextrorphan ((+)-3-hydroxy-N-methylmorphinan), their mixtures and the pharmaceutically acceptable salts thereof. Other useful nontoxic substances that block the NMDA receptor include ketamine (2-(2-chlorophenyl)-2-(methylamino)cyclohexanone), pyrrolizidine quinone and cis-4-(phosphonomethyl)-2-piperidinecarboxylic acid.

Nontoxic substances that block a major intracellular consequence of NMDA receptor activation and are therefore useful in the practice of the invention include inhibitors of

protein kinase C such as the gangliosides, in particular, ganglioside GM₁ (monosialoganglioside) and ganglioside GT_{1b} (trisialoganglioside); amphipathic long chain bases such as sphingosine, N,N,N-trimethylsphingosine, sphinganine and psychosine; quinolyloxazole-2-ones such as 4-methyl-5-(3-quinolinyl)-2-(3H)-oxazolone and phenyl-5-(2-quinolinyl)-2-(3H)-oxazolone; 1,4-bis-(amino-hydroxyalkylamino)anthraquinones such as 1,4-bis-(3-propylamino-2-hydroxypropylamino)-9,10 anthracenedione and 1,4-bis-(3-benzylamino-2-hydroxypropylamino)-9,10 anthracenedione; and, mixtures and pharmaceutically acceptable salts of any of the foregoing.

Additional nontoxic substances that block a major intracellular consequence of NMDA receptor activation and as such are useful in the practice of the invention include inhibitors of calmodulin such as the phenothiazines, in particular, chlorpromazine, chlorpromazine sulfoxide, prochlorperazine dimaleate, perphenazine, trifluoperazine, fluphenazine, fluphenazine enanthate, fluphenazine decanoate, thioridazine, mesoridazine besylate, piperacetazine, acetophenazine dimaleate, carphenazine dimaleate, butaperazine dimaleate and phenothiazine sulfoxide; naphthalenesulfonamides such as N-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide, N-(6-aminohexyl)-5-chloro-2-naphthalenesulfonamide and N-(6-aminohexyl)-5-bromo-2-naphthalenesulfonamide; 4-substituted-4H,6H-pyrrolo[1,2-a][4,1] benzoxazepines such as 1,3-dihydro-1-((1-[(4-methyl-4H,6H-pyrrolo[1,2-a][4,1] benzoxazepin-4-yl)methyl]-4-piperidinyl)-2H-benzimidazol-2-one; benzhydriyls such as N-[2-(diphenylmethylthio)ethyl]-2-(trifluoromethyl)benzeneethanamine, N-[2-(bis(4-fluorophenyl)methylthio)ethyl]-2-(trifluoromethyl)benzeneethanamine and N-[2-(bis(4-fluorophenyl)methylthio)ethyl]-3-(trifluoromethyl)benzeneethanamine; tricyclic antidepressant drugs such as imipramine, 2-chloroimipramine and amitriptyline; penfluridol; haloperidol; pimozide; clozapine; calmidazolium; and, mixtures and pharmaceutically acceptable salts of any of the foregoing.

Administration of the composition of this invention can be in the form of a single dosage unit containing both the addictive substance and the nontoxic substance that blocks the NMDA receptor or a major intracellular consequence of NMDA receptor activation or the two substances can be administered separately provided both are ultimately present in effective amounts in the patient. Introduction of the composition into the patient can be by way of oral administration or by intravenous, intramuscular, subcutaneous, intrathecal, epidural or intracerebroventricular injection.

The preferred dosage of addictive substance and the nontoxic substance that blocks the NMDA receptor or a major intracellular consequence of NMDA receptor activation can vary widely, e.g., from about 0.25 to about 250 mg/day, but actual amounts will vary according to the particular active substances being used, the particular formulation containing the active substances and the state and circumstances of the host being treated. As those skilled in the art recognize, many factors that modify the action of the active substances herein will be taken into account by the treating physician such as the age, body weight, sex, diet and condition of the subject, the time of administration, the rate and route of administration, and so forth. Optimal dosages for a given set of conditions can be ascertained by those skilled in the art using conventional dosage determination tests in view of the experimental data provided herein.

In alleviating withdrawal symptoms in addicted subjects deprived of the addictive substance, the substance that

blocks the NMDA receptor or a major intracellular consequence of NMDA receptor activation can be administered to the subject, together with the addictive substance, at a dosage rate of about 0.25 to about 250 mg/day, again, specific dosage levels and routes of administration being selected in accordance with the subject's circumstances. As a result of this treatment, the subject will experience a reduced level of dependence on the addictive substance eventually reaching the point where total withdrawal of the substance will result in at most mild withdrawal symptoms.

The composition herein can be formulated as a liquid, powder, elixir, injectable solution, etc. Formulations for oral use can be provided as hard gelatin capsules wherein the composition is mixed with an inert solid diluent such as calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the composition is mixed with an oleaginous medium, e.g., liquid paraffin or olive oil.

Aqueous suspensions can contain the composition in admixture with pharmaceutically acceptable excipients such as suspending agents, e.g., sodium carboxymethyl cellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, dispersing or wetting agents such as naturally occurring phosphatide, e.g., lecithin, or condensation products of an alkylene oxide with fatty acids, e.g., polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, e.g., heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol, e.g., polyoxyethylene sorbitol monooleate or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, e.g., polyoxyethylene sorbitan monooleate. Such aqueous suspensions can also contain one or more preservatives, e.g., ethyl- or n-propyl-p-hydroxy benzoate, one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as sucrose, saccharin or sodium or calcium cyclamate.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the composition in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, e.g., sweetening, flavoring and coloring agents, can also be present. Syrups and elixirs can be formulated with sweetening agents, for example glycerol, sorbitol or sucrose. Such formulations can also contain a demulcent, a preservative and flavoring and coloring agents.

The composition of this invention or either of its principal active ingredients can be provided in sustained release dosage form of which many kinds are known, e.g., as described in U.S. Pat. Nos. 4,788,055; 4,816,264; 4,828,836; 4,834,965; 4,834,985; 4,996,047; 5,071,646; and, 5,133,974, the contents of which are incorporated by reference herein.

The examples that follow are illustrative of the invention.

EXAMPLE 1

The effect of systemic dextrophan on prevention of the development of morphine tolerance and dependence was examined in Sprague-Dawley rats weighing 350-400 gm. Morphine tolerance was developed in the rats by twice daily subcutaneous injection of 10 mg/kg morphine sulfate. The analgesic effect of the morphine was examined by using the well known tail-flick test which measures the latency of tail-flick upon radiant heat stimulation. The latency of

tail-flick test is defined as the time elapsed from the onset of radiant heat to the flick of the rat's tail. In order to examine the effect of dextrorphan on the development of morphine tolerance, each morphine-treated rat also received intraperitoneal administration of either dextrorphan (1.56, 3.13, 6.25, 12.5 mg/kg, n=5/group) or saline (n=6) given 30 minutes prior to each morphine administration.

FIG. 1 shows the effects of systemic doses of dextrorphan (DEX) on tolerance to morphine analgesia produced by twice daily subcutaneous administration of 10 mg/kg morphine. Each symbol represents mean tail-flick latency scores (those above 4.5 seconds reflect analgesia) for each group of rats (N=5-6) and vertical bars are standard errors in this and the other figures. Baseline scores were between 4 and 5 seconds (at Day 0) and post-drug scores measured 1 hour after drug administration were close to 10 seconds for the first 5 days of daily drug administration. The control group (open triangles) show marked reduction in response to morphine (i.e., tolerance) at 7 and 9 days. In contrast, dextrorphan potently prevented the development of morphine tolerance as shown by no significant decreases in tail flick latencies, i.e., remaining analgesic during the whole course of repeated morphine administration. Asterisks indicate mean scores that were significantly different from those of the control group. All tested doses of dextrorphan were effective in preventing development of morphine tolerance with optimal doses ranging from 3.13 mg/kg to 12.5 mg/kg.

FIG. 2 shows the effects of systemic doses of dextrorphan on jumping, a withdrawal symptom produced by subcutaneous naloxone (2 mg/kg) in rats previously injected with morphine (10 mg/kg) twice daily for 9 days. Asterisks indicate median number of jumps in dextrorphan treatment groups (MOR+DEX) that were significantly less than that of the control group (MOR+SAL). Vertical bars refer to the range of the numbers of jumps. Thus, 3.13 and 6.25 mg/kg dextrorphan (but not 1.56 mg/kg) significantly reduced the incidence of jumping in morphine tolerant rats, a behavioral manifestation of morphine dependence, brought about following subcutaneous injection with 2 mg/kg naloxone. Thus, coadministration of dextrorphan with morphine greatly inhibits the development of both tolerance to and dependence on morphine while the analgesic effect of the morphine remains substantially unaffected.

EXAMPLE 2

The effects of ganglioside GM₁ in inhibiting morphine tolerance and dependence utilizing both systemic and intrathecal treatment were evaluated. The systemic treatment procedure, including both morphine and ganglioside GM₁ administration, was exactly the same as that used in the experimental work presented in Example 1 except that ganglioside GM₁ was given 1 hour before each morphine administration.

As shown in FIG. 3, the tail flick latency in ganglioside GM₁-treated (10, 30, 60 mg/kg, n=6/group) rats remained significantly longer than that of saline-treated rats on days 5, 7, 9, and 10 of repeated morphine administration, indicating the prevention of the development of morphine tolerance by ganglioside GM₁. Although all 3 doses of ganglioside GM₁ were effective, 30 and 60 mg/kg were more effective at days 9 and 10 than 10 mg/kg.

FIG. 4 shows the effects of systemic doses of ganglioside GM₁ on jumping, a withdrawal symptom produced by subcutaneous naloxone (2 mg/kg) in rats previously injected with morphine (10 mg/kg) twice daily for 9 days. Asterisks indicate median number of jumps in GM₁ treatment groups

(M+G) that were significantly less than that of the control group (MOR+SAL). Vertical bars refer to the range of the numbers of jumps. All three doses [10 mg/kg (10); 30 mg/kg (30); 60 mg/kg (60)] of GM₁ were effective with 60 mg/kg GM₁ being the most effective dose tested.

EXAMPLE 3

This example demonstrates the effectiveness of ganglioside GM₁ in preventing the development of morphine tolerance at the site of the spinal cord. Morphine sulfate 10 μg was delivered once daily through an intrathecal (spinal) canula implanted 5 days before the first morphine injection. Ganglioside GM₁ or saline also was delivered intrathecally 30 minutes before each morphine injection.

FIG. 5 shows the effects of the intrathecal doses of ganglioside GM₁ on tolerance to morphine analgesia produced by once daily intrathecal administration of 10 μg morphine. Intrathecal ganglioside GM₁ was given 30 min before each morphine administration. Each symbol represents mean scores for maximal possible effects (and hence analgesia) for each group of rats (n=5-6) measured at 15, 30, 60, 90, 120, 180, and 240 minutes after morphine injection on Day 8, i.e., 24 hours following 6 consecutive daily intrathecal morphine injections. Vertical bars are standard errors. Maximal possible effects (MPE) were calculated by the formula % MPE=[(TL-BL)/(10-BL)]×100. TL: actual tail-flick latency; BL: baseline latency obtained before the first morphine injection; 10: cut-off time for radiant heat stimulation. The control group (open circle) showed marked reduction in response to morphine (i.e., tolerance) at each tested time point. In contrast, treatment with ganglioside GM₁ effectively prevented the development of morphine tolerance as indicated by significantly higher maximal possible analgesia effects of morphine (160 nmol>80=40 nmol) as compared to those of saline-treated rats. Asterisks indicate mean scores that were significantly different from those of other groups.

EXAMPLE 4

The effects of ganglioside GM₁ and the toxic NMDA receptor antagonist MK 801 on morphine tolerance were evaluated.

As shown in FIG. 6, treatment with ganglioside GM₁ (60 mg/kg) inhibited the development of morphine tolerance to the degree equivalent to that induced by 0.3 mg/kg MK 801. However, 50% (3 out of 6) of the rats treated with 0.3 mg/kg MK 801 died before the completion of the experiment and the remaining rats in the group showed apparently poor grooming and weight loss indicating adverse effects of MK 801 on health. The rats treated with ganglioside GM₁ or dextrorphan continued to appear well groomed and did not show weight loss. None of the rats in the GM₁ or dextrorphan treatment groups died due to drug administration. Thus, dextrorphan and GM₁ are nontoxic in contrast to MK 801 which exhibits severe cytotoxic effects and as such, is unlikely to be approved by the FDA for administration to humans.

EXAMPLES 5 AND 6

These examples illustrate the effects of the phenothiazine trifluoperazine (Example 5) and the naphthalenesulfonamide N-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide hydrochloride (Example 6) in preventing the development of morphine tolerance in rats.

Calmodulin is an intracellular cofactor necessary for the nitric oxide pathway that can be initiated upon NMDA

receptor activation. FIGS. 7-10 demonstrate that trifluoperazine (TFP) and N-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide hydrochloride (W-7), both of which are calmodulin antagonists, effectively prevent the development of morphine tolerance in rats following intrathecal administration. In both cases, rats receiving once daily morphine sulfate (10 µg) and saline injection given intrathecally for 7 consecutive days developed tolerance to the analgesic effect or morphine as indicated by the reliable decrease in tail-flick latencies as compared to baseline latencies (day 1). In contrast, rats treated with TFP or W-7 (100 or 50 nmol) given immediately before each morphine administration showed a reliable analgesic effect of morphine (day 8) employing the same dose regimen used in the saline treatment group. The prevention of the development of morphine tolerance by TFP or W-7 is dose-related: 100=50 nmol>25=12.5 nmol. Each data point in FIGS. 7-10 represents the mean of a group of rats (n=6) and standard errors are shown by vertical lines. The asterisks refer to statistical differences ($\alpha=0.05$) between the saline group and each other group.

What is claimed is:

[1. A composition comprising an addictive substance and at least one nontoxic substance that blocks the N-methyl-D-aspartate receptor or a major intracellular consequence of N-methyl-D-aspartate receptor activation, the addictive substance being selected from the group consisting of alfentanil, alphaprodine, anileridine, bezitramide, codeine, dihydrocodeine, diphenoxylate, ethylmorphine, fentanyl, heroin, hydrocodone, hydromorphone, isomethadone, levomethorphan, levorphanol, metazocine, methadone, metopon, morphine, opium extracts, opium fluid extracts, powdered opium, granulated opium, raw opium, tincture of opium, oxycodone, oxymorphone, pethidine, phenazocine, piminodine, racemethorphan, racemorphan and pharmaceutically acceptable salts thereof.]

[2. The composition of claim 1 in sustained release dosage form.]

3. A formulated pharmaceutical composition comprising an addictive substance and at least one nontoxic synthetic substance that provides an improved effect for the addictive substance if used alone and that blocks the N-methyl-D-aspartate receptor and consists essentially of a morphinan or blocks a major intracellular consequence of N-methyl-D-aspartate receptor activation, the addictive substance being selected from the group consisting of alfentanil, alphaprodine, anileridine, bezitramide, codeine, dihydrocodeine, diphenoxylate, ethylmorphine, fentanyl, heroin, hydrocodone, hydromorphone, isomethadone, levomethorphan, levorphanol, metazocine, methadone, metopon, morphine, opium extracts, opium fluid extracts, powdered opium, granulated opium, raw opium, tincture of opium, oxycodone, oxymorphone, pethidine, phenazocine, piminodine, racemethorphan, racemorphan and pharmaceutically acceptable salts thereof.

4. A formulated pharmaceutical composition comprising an addictive substance and a non-toxic synthetic substance, the addictive substance being selected from the group consisting of alfentanil, codeine, dihydrocodeine, fentanyl, hydrocodone, hydromorphone, isomethadone, levorphanol, methadone, morphine, oxycodone, oxymorphone, pethidine, and pharmaceutically acceptable salts thereof, the non-toxic synthetic substance providing an improved effect for the addictive substance if used alone and being selected from the group consisting of dextromethorphan, dextrorphan, and pharmaceutically acceptable salts thereof.

5. A formulated pharmaceutical composition comprising an addictive substance and a non-toxic synthetic substance,

the addictive substance being selected from the group consisting of alfentanil, codeine, dihydrocodeine, fentanyl, hydrocodone, hydromorphone, isomethadone, levorphanol, morphine, oxycodone, oxymorphone, pethidine, and pharmaceutically acceptable salts thereof, the non-toxic synthetic substance being a blocker of the N-methyl-D-aspartate receptor and consisting essentially of morphinans, and providing an improved effect for the addictive substance if used alone.

6. A composition according to claims 3, 4 or 5 wherein the addictive substance is selected from the group consisting of alfentanil, codeine, dihydrocodeine, fentanyl, hydrocodone, hydromorphone, isomethadone, methadone, morphine, oxycodone, oxymorphone, pethidine, and pharmaceutically acceptable salts thereof.

7. A composition according to claims 3, 4 or 5 wherein the addictive substance is selected from the group consisting of alfentanil, codeine, dihydrocodeine, fentanyl, isomethadone, methadone, pethidine, and pharmaceutically acceptable salts thereof.

8. A composition according to claims 3, 4 or 5 wherein the addictive substance is selected from the group consisting of codeine, methadone, and pharmaceutically acceptable salts thereof.

9. A composition according to claims 3, 4 or 5 wherein the addictive substance includes morphine or a pharmaceutically acceptable salt thereof.

10. A composition according to claims 3, 4 or 5 wherein the addictive substance includes oxycodone or a pharmaceutically acceptable salt thereof.

11. A composition according to claims 3, 4 or 5 wherein the addictive substance includes hydrocodone or a pharmaceutically acceptable salt thereof.

12. A composition according to claims 3, 4 or 5 wherein the addictive substance includes oxymorphone or a pharmaceutically acceptable salt thereof.

13. A composition according to claim 3, 4 or 5 wherein the addictive substance includes hydromorphone or a pharmaceutically acceptable salt thereof.

14. A composition according to claims 3, 4 or 5, in oral dosage form.

15. A composition according to claims 3, 4 or 5, in sustained release dosage form.

16. A composition according to claims, 3, 4 or 5, in oral dosage and sustained release dosage form.

17. A formulated pharmaceutical composition comprising an addictive substance and at least one nontoxic synthetic substance that blocks the N-methyl-D-aspartate receptor or a major intracellular consequence of N-methyl-D-aspartate receptor activation, selected from pyrroloquinoline quinone and cis-4-(phosphonomethyl)-2-piperidinecarboxylic acid, the addictive substance being selected from the group consisting of alfentanil, alphaprodine, anileridine, bezitramide, codeine, dihydrocodeine, diphenoxylate, ethylmorphine, fentanyl, heroin, hydrocodone, hydromorphone, isomethadone, levomethorphan, levorphanol, metazocine, methadone, metopon, morphine, opium extracts, opium fluid extracts, powdered opium, granulated opium, raw opium, tincture of opium, oxycodone, oxymorphone, pethidine, phenazocine, piminodine, racemethorphan, racemorphan, racemorphan and pharmaceutically acceptable salts thereof.