

US00RE40789E

(19) **United States**
 (12) **Reissued Patent**
Elledge et al.

(10) **Patent Number:** **US RE40,789 E**
 (45) **Date of Reissued Patent:** **Jun. 23, 2009**

(54) **MAMMALIAN CHECKPOINT PROTEINS
 POLYPEPTIDES AND ENCODING
 SEQUENCES THEREOF**

(75) Inventors: **Stephen J. Elledge**, Houston, TX (US);
Yolanda Sanchez, Cincinnati, OH (US)

(73) Assignee: **Baylor College of Medicine**, Houston,
 TX (US)

(21) Appl. No.: **10/267,717**

(22) Filed: **Oct. 9, 2002**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **6,307,015**
 Issued: **Oct. 23, 2001**
 Appl. No.: **09/488,364**
 Filed: **Jan. 12, 2000**

U.S. Applications:

(62) Division of application No. 08/924,183, filed on Sep. 5,
 1997, now Pat. No. 6,218,109.

(51) **Int. Cl.**
C12N 9/12 (2006.01)
C12N 9/10 (2006.01)
A61K 38/00 (2006.01)
C07K 1/00 (2006.01)

(52) **U.S. Cl.** **435/194**; 435/193; 530/324;
 530/326; 530/350

(58) **Field of Classification Search** 530/350,
 530/326, 827; 536/23.5; 435/69.1, 252.3,
 435/320.1

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,655,516 A * 4/1972 Roon et al. 435/12
 3,962,037 A * 6/1976 Mitchell 435/188
 4,347,323 A * 8/1982 Imamura et al. 435/194
 5,302,706 A 4/1994 Smith 536/23.1
 5,504,000 A * 4/1996 Littman et al. 435/194
 5,518,911 A * 5/1996 Abo et al. 435/194
 6,383,744 B1 * 5/2002 Green et al. 435/6

OTHER PUBLICATIONS

Sanchez et al., *Science*, (Sep. 1997), vol. 277, pp.
 1497–1501.*

Walworth et al., “rad-Dependent Response of the
 chk1-Encoded Protein Kinase at the DNA Damage Check-
 point,” *Science, News Series*, vol. 271, No. 5247 (Jan. 19,
 1996) pp. 353–356.*

Sanchez et al., “Conservation of the Chk1 Checkpoint Path-
 way in Mammals: Linkage to DNA Damage to Cdk Regula-
 tion Through Cdc25,” *Science*, Sep. 5, 1997, pp. 197–1501.*

O’Connell et al., “Chk1 is a wee1 kinase in the G2 DNA
 Damage checkpoint inhibiting cdc2 by Y15 phosphoryla-
 tion,” *The EMBO Journal*, vol. 16, No. 3, Feb. 1997, pp.
 545–554.*

Borges et al., *Journal of Immunological Methods*, vol. 173
 (1994), pp. 253–263.*

Hideo Toyoshima and Tony Hunter, “p27, A Novel Inhibi-
 tors of G1 Cyclin-Cdk Protein Kinase Activity, is Related to
 p21,” *Cell*, vol. 78, Jul. 15, 1994, pp. 67–74.

Shou Waga, Gregory J. Hannon, David Beach & Bruce Still-
 man, “The p21 Inhibitor of Cyclin-Dependent Kinases Con-
 trols DNA Replication by Interaction With PCNA,” *Nature*,
 vol. 369, Jun. 16, 1994, pp. 574–578.

Stephen J. Elledge and J. Wade Harper, “Cdk Inhibitors: on
 the Threshold of Checkpoints and Development,” *Cell Biol-
 ogy*, 1994, pp. 847–850.

Jin Jen, J. Wade Harper, Sandra H. Bigner, Darell D. Bigner,
 Nickolas Papadopoulos, Sanford Markowitz, James K.V.
 Willson, Kenneth W. Kinzler and Bert Vogelstein, “Deletion
 of p16 and p15 Genes in Brain Tumors,” *Cancer Research*,
 Dec. 15, 1994, pp. 6353–6358.

Gregory J. Hannon and David Beach, “p15^{INK48} is a Potent-
 ial Effector of TGF- β -Induced Cell Cycle Arrest,” *Nature*,
 vol. 371, Sep. 15, 1994, pp. 257–261.

Jamison Nourse, Eduardo Firpo, W. Michael Flanagan,
 Steve Coats, Kornelia Polyak, Mong-Hong Lee, Joan Mas-
 sague, Gerald R. Crabtree and James M. Roberts,
 “Interleukin-2-Mediated Elimination of the p27^{Kip1}
 Cyclin-Dependent Kinase Inhibitor Prevented by Rapamy-
 cin,” *Nature*, vol. 372, Dec. 8, 1994, pp. 570–573.

Jun-Ya Kato, Masaaki Matsuoka, Kornelia Polyak, Joan
 Massague and Charles J. Sherr, “Cyclin Amp-Induced G1
 Phase Arrest Mediated by an Inhibitor (p27^{Kip1}) of
 cyclin-Dependent Kinase 4 Activation,” *Cell*, vol. 79, Nov.
 4, 1994, pp. 487–496.

Hideo Toyoshima and Tony Hunter, “p27, A Novel Inhibitor
 of G1 Cyclin-CDK Protein Kinase Activity, is Related to
 p21,” *Cell*, vol. 78, Jul. 15, 1994, pp. 67–74.

Kornelia Polyak, Mong-Hong Lee, Hediye Erdument-Bro-
 mage, Andrew Koff, James M. Roberts, Paul Tempst and
 Joan Massague, “Cloning of P27^{Kip1}, A Cyclin-Dependent
 Kinase Inhibitor and a Potential Mediator of Extracellular
 Antimitogenic Signals,” *Cell*, vol. 78, Jul. 15, 1994, pp.
 59–66.

Richard A. Steinman, Barbara Hoffman, Augustine Iro,
 Christel Guillouf, Dan A. Lieberman and Motawa E.
 El-Houseini, “Induction of p21 (WAF-1/CIP1) During Dif-
 ferentiation,” *Oncogene*, 1994, pp. 3389–3396.

Hongping Jiang, Jian Lin, Zao-Zhong, Su, Frank R. Collart,
 Eliezer Huberman and Paul B. Fisher, “Induction of Differ-
 entiation in Human Promyelocytic HL-60 Leukemia Cells
 Activates p21, WAF1/CIP1, Expression in the Absence of
 p53,” *Oncogene*, 1994, pp. 3397–3406.

(Continued)

Primary Examiner—Bradley L Sisson

(74) *Attorney, Agent, or Firm*—Jennifer S. Sickler;
 Monique Vander Molen; Gardere Wynne Sewell

(57) **ABSTRACT**

The present invention relates to the isolation of gene
 sequences encoding mammalian cell cycle checkpoints, as
 well as the expression of the encoded proteins using recom-
 binant DNA technology. The expressed proteins are used to
 generate specific antibodies and to inhibit the growth of
 cells. The human checkpoint gene sequences are used as a
 probe for a portion of the chromosome associated with
 tumors and other malignancies, as well as growth and/or
 development deficiencies.

7 Claims, 8 Drawing Sheets

OTHER PUBLICATIONS

- M. Saeed Sheikh, Xiao-Su Li, Jian-Chyi Chen, Zhi-Ming Shao, Jose V. Ordonex and Joseph A. Fontana, "Mechanisms of Regulation of WAF1/Cip1 Gene Expression in Human Breast Carcinoma: Role of P53-Dependent and Independent Signal Transduction Pathways," *Oncogene*, 1994, pp. 3407-3415.
- Paolo Michieli, Marcio Chedid, David Lin, Jacalyn H. Pierce, W. Edward Mercer and David Givol, "Induction of WAF1/CIP1 by a P53-Independent Pathway," *Cancer Research*, Jul. 1, 1994, pp. 3391-3395.
- Wafik S. El-Deiry, J. Wade Harper, Patrick M. O'Connor, Victor E. Velculescu, Christine E. Canman, Joany Jackman, Jennifer A. Pierenpol, Marilee Burrell, David E. Hill, Yisong Wang, Klas G. Wiman, W. Edward Mercer, Michael B. Kastan, Kurt W. Kohn, Stephen J. Elledge, Kenneth W. Kinzler and Bert Vogelstein, "WAF1/CIP1 is Introduced in P53-Mediated G1 Arrest and Apoptosis," *Cancer Research*, Mar. 1, 1994, pp. 1169-1174.
- Susan B. Parker, Gregor Eichele, Pumin Zhang, Alan Rawls, Arthur T. Sands, Allan Bradley, Eric N. Olson, J. Wade Harper and Stephen J. Elledge, "P53-Independent Expression of p21^{CIP1} in Muscle and Other Terminally Differentiating Cells," *Science*, vol. 267, Feb. 17, 1995, pp. 1024-1027.
- Vjekoslav Dulic, William K. Kaufmann, Sandra J. Wilson, Thea D. Tisty, Emma Lees, J. Wade Harper, Stephen J. Elledge and Steven I. Reed, "P53-Dependent Inhibitor of Cyclin-Dependent Kinase Activities in Human Fibroblasts During Radiation-Induced G1 Arrest," *Cell*, vol. 76, Mar. 25, 1994, pp. 1013-1023.
- J. Wade Harper, Stephen J. Elledge, Khandan Keyomarsi, Brian Dynlacht, Li-Huei Tsai, Pumin Zhang, Steven Dobrowolski, Chang Bai, Lisa Connell-Crowley, Eric Swindell, M. Pat Fox and Nan Wei, "Inhibition of Cyclin-Dependent Kinases by p21," *Molecular Biology of the Cell*, Apr. 1995, vol. 6, pp. 387-400.
- Shou Waga, Gregory J. Hannon, David Beach and Bruce Stillman, "The p21 Inhibitor of Cyclin-Dependent Kinases Controls DNA Replication by Interaction with PCNA," *Nature*, vol. 369, Jun. 16, 1994, pp. 574-578.
- Hernan Flores-Rozas, Zvi Kelman, Frank B. Dean, Zhen-Qiang Pan, J. Wade Harper, Stephen J. Elledge, Michael O'Donnell and Jerard Hurwitz, "Cdk-Interacting Protein 1 Directly Binds with Proliferating Cell Nuclear Antigen and Inhibits DNA Replication Catalyzed by the DNA Polymerase Holoenzyme," *Proc. Natl. Acad. Sci. USA*, vol. 91, Aug. 1994, pp. 8655-8659.
- Stephen J. Elledge and J. Wade Harper, "Cdk Inhibitors: on the Threshold of Checkpoints and Development," *Biology*, 1994, pp. 847-852.
- Manuel Serrano, Gregory J. Hannon and David Beach, "A New Regulatory Motif in Cell-Cycle Control Causing Specific Inhibition of Cyclin D/CDK4," *Nature*, vol. 366, Dec. 16, 1993, pp. 704-707.
- Asao Noda, Yi Ning, Susan F. Venable, Olivia M. Pereira-Smith and James R. Smith, "Cloning of Senescent Cell-Derived Inhibitors of DNA Synthesis Using an Expression Screen," *Experimental Cell Research* 1994, pp. 90-98.
- Yue Xiong, Gregory J. Hannon, Hui Zhang, David Casso, Ryuji Kobayashi and David Beach, "p21 is a Universal Inhibitor of Cyclin Kinases," *Nature*, vol. 366, Dec. 16, 1993, pp. 701-704.
- Wafik S. El-Deiry, Takashi Tokino, Victor E. Velculescu, Daniel B. Levy, Ramon Parsons, Jeffrey M. Trent, David Lin, W. Edward Mercer, Kenneth W. Kinzler and Bert Vogelstein, "WAF1, A Potential Mediator of p53 Tumor Suppression," *Cell*, vol. 75, Nov. 19, 1993, pp. 817-825.
- Mark J. Solomon, "Activation of the Various Cyclin/cdc2 Protein Kinases," *Cell Biology*, 1993, pp. 180-186.
- Charles J. Sherr, "G1 Phase Progression: Cycling on Cue," *Cell*, vol. 79, Nov. 18, 1994, pp. 551-555.
- Dawn Coverely, Mark Jackman and Julia Turner, "Cyclins and Cyclin-Dependent Kinases: Take Your Partners," *TIBS*, Jun. 1993, pp. 195-197.
- Tsutomu Nobori, Kaoru Miura, David J. Wu, Augusto Lois, Kenji Takayashi and Dennis A. Carson, "Deletions of the Cyclin-Dependent Kinase-4 Inhibitor Gene in Multiple Human Cancers," *Nature*, vol. 368, Apr. 21, 1994, pp. 753-756.
- Kun-Liang Guan, Christopher W. Jenkins, Yan Li, Michael A. Nicols, Xiaoyu Wu, Christine L. O'Keefe, A. Gregory Matera and Yue Xiong, "Growth Suppression by p18, a p16^{INK4/MTS1} and p14^{INK4B/MTS2}-Related CDK6 Inhibitor, Correlates with Wild-Type pRb Function," *Genes and Development*, 1994, pp. 2939-2952.
- Jacob W. Udo, Elizabeth A. Lindsay, Richard A. Wells and Antonio Baldini, "Multiple Variants in Subtelomeric Regions of Normal Karyotypes," *Genomics*, 1992, pp. 1019-1025.
- Olof H. Sudin, Heinz G. Busse, Melissa B. Rogers, Lorraine J. Gudas and Gregor Eichele, "Region-Specific Expression in Early Chick and Mouse Embryos of Ghox-Lab and Hox 1.6, Vertebrate Home,box-Containing Genes Related to Drosophila Labial," *Development*, 1990, pp. 47-58.
- Beat Lutz, Shigeru Kuratani, Austin J. Cooney, Stefan Wawersik, Sophia Y. Tsai, Gregor Eichele and Ming-Jer Tsai, "Developmental Regulation of the Orphan Receptor Coup-RFII Gene in Spinal Motor Neurons," *Development*, 1994, pp. 25-36.
- Tim Durfee, Kathleen Becherer, Phang-Lang Chen, Shiou-Hwei Yeh, Yanzhu Yang, April E. Kilburn, Wen-Hwa Lee and Stephen J. Elledge, "The Retinoblastoma Protein Associates with the Protein Phosphatase Type 1 Catalytic Subunit" Dec. 18, 1992, pp. 555-569.
- Charles H. Spruck III, Mirella Gonzalez-Zulueta, Atsuko Shibata, Anne R. Simoneau, Ming-Fong Lin, Felicidad Gonzales, Yvonne C. Tsai and Peter A. Jones, "p16 Gene in Uncultured Tumours," *Nature*, vol. 370, Jul. 21, 1994, pp. 183-184.
- Alexander Kamb, Qingyun Liu, Keith Harshman and Sean Tavtigian, "Rates of p16 (MTS1) Mutations in Primary Tumors with 9p Loss," *Science*, vol. 165, Jul. 15, 1994, pp. 415-417.
- M.J. Pettenati, J.L. Haines, R.R. Higgins, R.S. Wappner, C.G. Palmer and D.D. Weaver, "Wiedemann-Beckwith Syndrome: Presentation of Clinical and Cytogenetic Data on 22 New Cases and Review of the Literature," *Human Genetics*, 1986, pp. 143-154.
- N.D. Hastie, "The Genetics of Wilms' Tumor—A Case of Disrupted Development," *Annu. Rev. Genet.*, 1994, pp. 523-558.
- B.R. Seizinger, H.P. Klinger, C. Junien, Y. Nakamura, M. Le Beau, W. Cavenee, B. Emanuel, B. Ponder, S. Naylor, F. Mitelman, D. Louis, A. Menon, I. Newsham, J. Decker, M. Kaelbling, I. Henry, A.v. Deimling, "Report of the Committee on Chromosome and Gene Loss in Human Neoplasia," *Cytogenet Cell Genet*, 1991, pp. 1080-1096.

- Ueli Gubler and Beth J. Hoffman, "A Simple and Very Efficient Method for Generating cDNA Libraries," Elsevier Science Publishers, 1983, pp. 263–269.
- F. L. Graham and A.J. Van Der Eb, "A New Technique for the Assay of Infectivity of Human Adenovirus 5 DNA," *Virology* 52, 1973, pp. 456–467.
- Dan Y. Wu and R. Bruce Wallace, "The Ligation Amplification Reaction (LAR)—Amplification of Specific DNA Sequences Using Sequential Rounds of Template-Dependent Ligation," *Genomics*, 1989, pp. 560–569.
- Stephen J. Elledge and J. Wade Harper, "Cdk Inhibitors: on the Threshold of Checkpoints and Development," *Cell Biology*, 1994, pp. 847–852.
- Mark G. Alexandrow and Harold L. Moses, "Transforming Growth Factor B and Cell Cycle Regulation," *Cancer Research*, Apr. 1, 1995, pp. 1452–1457.
- Kim Nasmyth and Tim Hunt, "Dams and Sluices," *Nature*, vol. 366, Dec. 16, 1993, pp. 634–635.
- Tony Hunter and Jonathon Pines, "Cyclins and Cancer II: Cyclin D and CDK Inhibitors Come of Age," *Cell*, vol. 79, Nov. 18, 1994, pp. 573–582.
- Jesse Martinex, Ingo Georgoff, Joanne Martinez and Arnold J. Levine, "Cellular Localization and Cell Cycle Regulation by a Temperature-Sensitive p53 Protein," *Genes and Development*, 1991, pp. 151–159.
- Sander van den Heuvel and Ed Harlow, "Distinct Roles for Cyclin-Dependent Kinases in Cell Cycle Control," *Science*, vol. 262, Dec. 24, 1993, pp. 2050–2054.
- Liang Zhu, Sander van den Heuvel, Kristian Helin, Ali Fataey, Mark Ewen, David Livingston, Nicholas Dyson and Ed Harlow, "Inhibition of Cell Proliferation by p107, A Relative of the Retinoblastoma Protein," *Genes and Development*, 1993, pp. 1111–1125.
- Stephen J. Elledge, Ronald Richman, Frederick L. Hall, Richard T. Williams, Naomi Lodgson and J. Wade Harper, "CDK2 Encodes A 33-KDA Cyclin A-Associated Protein Kinase and is Expressed Before CDC2 in the Cell Cycle," *Proc. Natl. Acad. Sci. USA*, vol. 89, Apr. 1992, pp. 2907–2911.
- Hui Zhang, Gregory J. Hannon and David Beach, "p-21 Containing Cyclin Kinases Exist in Both Active and Inactive States," *Genes & Development*, 1994, pp. 1750–1758.
- Nat Sternberg, "Bacteriophage P1 Cloning System for the Isolation, Amplification, and Recovery of DNA Fragments as Large as 100 Kilobase Pairs," *Proc. Natl. Acad. Sci. USA*, vol. 87, Jan. 1990, pp. 103–107.
- Matthew Meyerson and Ed Harlow, "Identification of G1 Kinase Activity for cdk6, A Novel Cyclin D Partner," *Molecular and Cellular Biology*, Mar. 1994, vol. 14, No. 3, pp. 2077–2086.
- Hitoshi Matsushime, Mark E. Ewen, David K. Strom, Jun-Ya Kato, Steven K. Hanks, Martine F. Roussel and Charles J. Sherr, "Identification and Properties of an Atypical Catalytic Subunit (P34^{psk-j3}/CDK4) For Mammalian D Type G1 Cyclins," *Cell*, vol. 71, Oct. 16, 1992, pp. 323–334.
- Giulio Draetta, "CDC2 Activation: The Interplay of Cyclin Binding and Thr161 Phosphorylation," *Trends in Cell Biology*, vol. 3, Sep. 3, 1993, pp. 287–289.
- Claudine Junien, "Beckwith-Wiedemann Syndrome, Tumorigenesis and Imprinting," *Current Biology*, 1992, pp. 431–438.
- John Wiley & Sons, Inc., "Current Protocols in Molecular Biology, vol. 1," 1994, Section III, 8 pages.
- Orna Halevy, Bennett G. Novitch, Douglas B. Spicer, Stephen X. Skapek, James Rhee, Gregory J. Hannon, David Beach and Andrew B. Lassar, "Correlation of Terminal Cell Cycle Arrest of Skeletal Muscle With Induction of p21 by MyoD," *Science*, vol. 267, Feb. 17, 1995, pp. 1018–1021.
- L. Breeden and K. Nasmyth, "Regulation of the Yeast Ho Gene," pp. 643–650 (no date).

* cited by examiner

ggcc gga cag tcc gcc gag glg etc ggt gga gtc atg gca gtg ccc ttt gtg gaa gac tgg
D L V Q T L G E G A Y G E V Q L A V N R
gac ttg gtg cao occ ctg ggo gao ggt gcc tat ggo gaa gtt caa ctt gct glg aat aga
D L V Q T L G E G A Y G E V Q L A V N R
gta act gaa gao gca gtc gca gtg aag att gta gat atg aag cgt gcc gta gac tgt cca
V T E E A V A V K I V D M K R A V D C P
gaa aat att aag aoo gag atc tgt atc aat aaa atg cta aat cat gaa aat gla gla aoo
E N I K K E I C I N K M L N H E N V V K
llc tat ggt cac agg oga gaa ggc aat atc cao tat tta ttt ctg gag lac tgt agt gga
F Y G H R R E G N I Q Y L F L E Y C S G
gga gag ctt ttt gac oga ata gag cca gac ata ggc atg cct gaa cca gat gct cag oga
G E L F D R I E P D I G M P E P D A Q R
llc ttc cat cao ctc atg gca ggg gtg gtt tat ctg cat ggt att gga ata act cac agg
F F H Q L M A G V V Y L H G I G I T H R
gat att aoo cca gaa aat ctt ctg ttg gat gaa agg gat aac ctc aoo atc tca gac ttt
D I K P E N L L L D E R D N L K I S D F
ggc ttg gca aca gla ttt cgg tat aat aat cgt gag cgt ttg ttg aac aag atg tgt ggt
G L A T V F R Y N N R E R L L N K M C G
oct tta cca tat gtt gct cca gaa ctt ctg aag oga oga gaa ttt cat gca gga cca gtt
T L P Y V A P E L L K R R E F H A E P V
gat gtt tgg tcc tgt gga ata gla ctt oct gca atg ctc gcl gga gaa ttg cca tgg gac
D V W S C G I V L T A M L A G E L P W D
cao ccc agt gac agc tgt cog gag tot tct gac tgg aaa gaa aoo aoo aca tac ctc aac
Q P S D S C Q E Y S D W K E K K T Y L N
cct tgg aoo aoo atc gat tct gct cct cta gct ctg ctg cat aoo atc tta gtt gag aat
P W K K I D S A P L A L H K I L V E N
cca tca gca oga att occ att cca gac atc aoo aoo gat oga tgg tac aac aoo ccc ctc
P S A R I T I P D I K K D R W Y N K P L
aag aoo ggg gca aoo agg ccc cgo gtc act tca ggt ggt gtg tca gag tct ccc agt gga
K K G A K R P R V T S G G V S E S P S G
ttt tct aag cac att cao tcc aot ttg gac ttc tct cca gla aac agt gct tct agt gao
F S K H I Q S N L D F S P V N S A S S E
gaa aat gtg aag tac tcc agt tct cog cca gaa ccc cgc aca ggt ctt tcc tta tgg gat
E N V K Y S S S Q P E P R T G L S L W D
acc agc ccc tca tac att gat aoo ttg gla cao ggg atc agc ttt tcc cag ccc aca tgt
T S P S Y I D K L V Q G I S F S Q P T C
cct gat cat atg ctt llg aat agt cog tta ctt ggc occ cca gga tcc tca cog aac ccc
P D H M L L N S Q L L G T P G S S O N P
tgg cog cgg ttg gtc aoo oga atg aco cga ttc ttt occ aoo ttg gat gca gac aoo tct
W Q R L V K R M T R F F T K L D A D K S
tat cao tgc ctg aoo gag act tgt gag aag ttg ggc tat cao tgg aag aoo agt tgt atg
Y Q C L K E T C E K L G Y Q W K K S C M
aat cog gtt act ata tca aca act gat agg oga aac aat aoo ctc att ttc aoo gtg aot
N Q V T I S T T D R R N N K L I F K V N
llg tta gaa atg gat gat aoo ata ttg gtt gac ttc cgg ctt tct aag ggt gat gga ttg
L L E M D D K I L V D F R L S K G D G L
gag ttc aag oga cac ttc ctg aag att aoo ggg aag ctg att gat att gtg agc agc cog
E F K R H F L K I K G K L I D I V S S Q
aag gtt tgg ctt cct gcc oca tga tca gac cat cgg ctc tgg gga atc ctg glg aot ota
K V W L P A T
glg ctg cta tgt tga cat tot tct tcc tag oga oga tta tcc tgt cct gca aac tgc aoo
tag tag ttc ctg aag tgt tca ctt ccc tgt tta tcc aoo cat ctt cca att tot ttt gtt
tgt tag gca tac aoo tao tac cta tat ctt aot tgt aag caa aac ttt ggg gaa agg atg
aot oga att cat ttg att att tct tca tgt glg ttt agt atc tga att tga aac tca tct
ggt gga aac cao gtt tca ggg gac atg agt ttt cca gcl ttt ata cac acg tot ctc att
ttt atc aoo aco ttt tgt tt

FIG. 1

gctt	glc	gct	glg	ctt	gga	gtc	atg	gca	glg	ccl	lft	gtg	gaa	goc	lgg	gat	llg	gtg	caa
							M	A	V	P	F	V	E	D	W	D	L	V	Q
act	llg	gga	gaa	ggt	gcc	tat	gga	gaa	ggt	caa	ctt	gct	gtg	aat	aga	ala	act	gaa	caa
T	L	G	E	G	A	Y	G	E	V	Q	L	A	V	N	R	I	T	E	Q
gct	ggt	gca	gtg	aaa	att	gta	gac	atg	aag	cgg	gcc	ota	goc	tgt	cca	caa	aat	att	aag
A	V	A	V	K	I	V	D	M	K	R	A	I	D	C	P	Q	N	I	K
aaa	gag	atc	tgc	atc	aat	aaa	atg	lta	agc	cac	gag	aat	gta	gtg	aaa	ttc	tat	ggc	coc
K	E	I	C	I	N	K	M	L	S	H	E	N	V	V	K	F	Y	G	H
aag	agg	gaa	ggc	cat	atc	cag	tat	ctg	ttt	ctg	gag	lac	lgt	ogt	gga	gga	gaa	ctt	ttt
R	R	E	G	H	I	Q	Y	L	F	L	E	Y	C	S	G	G	E	L	F
gat	aga	att	gag	cca	gac	ata	ggg	atg	cct	gaa	caa	gat	gct	cag	agg	ttc	ttc	cac	caa
D	R	I	E	P	D	I	G	M	P	E	Q	D	A	Q	R	F	F	H	Q
ctc	atg	gca	ggg	gtg	ggt	tat	ctt	cat	gga	att	gga	alo	act	cac	agg	gat	att	aaa	cca
L	M	A	G	V	V	Y	L	H	G	I	G	I	T	H	R	D	I	K	P
gaa	ooc	ctc	ctc	llg	gat	gaa	agg	gat	aac	ctc	aaa	alc	lcl	gac	ttt	ggc	llg	gca	acg
E	N	L	L	L	D	E	R	D	N	L	K	I	S	D	F	G	L	A	T
gta	ttt	cgg	cat	aat	aat	cgt	gaa	cgc	lta	ctg	aac	aag	atg	lgt	ggg	act	lta	cct	tat
V	F	R	H	N	N	R	E	R	L	L	N	K	M	C	G	T	L	P	Y
glt	gct	ccg	gag	ctt	cta	aag	aga	aaa	gaa	ttt	cat	gca	gaa	cca	glt	gat	glt	tgg	lcc
V	A	P	E	L	L	K	R	K	E	P	H	A	E	P	V	D	V	W	S
tgt	gga	ala	gta	ctt	act	gca	atg	llg	gct	gga	gaa	llg	ccg	lgg	gac	cag	ccc	agt	gat
C	G	I	V	L	T	A	M	L	A	G	E	L	P	W	D	Q	P	S	D
agc	tgt	cag	gaa	tat	tct	gat	tgg	aaa	gaa	aaa	aaa	acc	tal	ctc	aat	cct	lgg	aaa	aaa
S	C	Q	E	Y	S	D	W	K	E	K	K	T	Y	L	N	P	W	K	K
att	got	tct	gct	cct	ctg	gct	ttg	ctt	cat	aaa	att	cto	gll	gag	act	cca	tca	gca	agg
I	D	S	A	P	L	A	L	L	H	K	I	L	V	E	T	P	S	A	R
atc	acc	atc	cca	gac	att	aag	aaa	gat	aga	tgg	lac	aac	aaa	cca	ctt	aac	aga	gga	gca
I	T	I	P	D	I	K	K	D	R	W	Y	N	K	P	L	N	R	G	A
aag	agg	cca	cgc	gcc	oca	tca	ggt	ggt	atg	tca	gag	tct	lct	agt	gga	ttc	tct	aag	cac
K	R	P	R	A	T	S	G	G	M	S	E	S	S	S	G	F	S	K	H
att	cat	tcc	aat	llg	gac	llt	tct	cca	gta	aat	aat	ggt	tcc	agt	gaa	gaa	acc	glg	aag
I	H	S	N	L	D	F	S	P	V	N	N	G	S	S	E	E	T	V	K
ttc	tct	ogt	tcc	cag	cca	gag	ccg	aga	aca	ggg	ctt	tcc	llg	lgg	gac	act	ggt	ccc	tcg
F	S	S	S	Q	P	E	P	R	T	G	L	S	L	W	D	T	G	P	S
aac	gtg	gac	aaa	ctg	ggt	cag	ggc	atc	agt	ttt	tcc	cag	cct	acg	tgt	cct	gag	cat	otg
N	V	D	K	L	V	Q	G	I	S	F	S	Q	P	T	C	P	E	H	M
ctt	gto	ooc	ogt	cag	lta	ctc	ggt	occ	cct	gga	llt	tca	cag	aac	ccc	lgg	cag	cgc	ttg
L	V	N	S	Q	L	Q	G	T	P	G	F	S	Q	N	P	W	Q	R	L
gtc	aaa	agg	atg	aca	cga	ttc	llt	act	aaa	ttg	gat	gcg	gac	aaa	lct	loc	caa	tgc	ctg
V	K	R	M	T	R	F	F	T	K	L	D	A	D	K	S	Y	O	C	L
aaa	gag	occ	ttc	gag	aag	ttg	ggc	tat	cag	tgg	aag	aag	agt	tgt	atg	aat	cag	glt	act
K	E	T	F	E	K	L	G	Y	Q	W	K	K	S	C	M	N	Q	V	T
gta	tca	oca	oct	gat	oga	ogo	aac	aat	aag	ttg	att	ttc	aaa	ota	aat	ttg	gta	gaa	atg
V	S	T	T	D	R	R	N	N	K	L	I	F	K	I	N	L	V	E	M
got	gag	oog	ata	ctg	ggt	gac	ttc	cga	ctt	tct	aag	ggt	gat	gga	lta	gag	ttc	aog	ago
D	E	K	I	L	V	D	F	R	L	S	K	G	D	G	L	E	F	K	R
cac	ttc	ctg	aag	att	aaa	ggg	aag	ctc	agc	gat	ggt	glg	agc	ogc	cag	aag	glt	tgg	ttt
H	F	L	K	I	K	G	K	L	S	D	V	V	S	S	Q	K	V	W	F
cct	ggt	aco	tga	gga	agc	tgt	cag	ctc	tgc	aco	ttc	ctg	gtg	aat	ogo	glg	ctg	cta	tgt
P	V	T	*																
gac	att	ttt	ctt	cct	aga	gaa	gat	tat	cta	ttc	tgc	aaa	ctg	caa	aca	ata	glt	glt	gaa
gag	ttc	tct	tcc	cat	tac	cca	aac	atc	ttc	cga	ttt	gta	gtg	ttt	ggc	ata	caa	ata	cta
atg	tat	ttt	aat	tgt	atg	taa	tgc	ttt	ggg	gaa	agg	atg	gat	caa	att	cat	lag	gta	ttt
gtc	cag	ctg	tct	lta	aat	tgt	ctg	gat	ttg	aaa	cca	agt	lat	ggg	ata	ctt	gag	ttt	gcc
agc	ttt	tat	acc	cat	gta	gta	gta	tca	ctt	llg	aaa	aat	caa	aag	ctt	glt	tca	tcc	caa
gca	aaa	tat	ttt	ctt	ctc	tgc	cta	ttt	aat	tgt	aag	gat	gaa	tca	aca	cag	acc	ata	tac
agt	tga	ttg	ggt	cat	gaa	tga	ggc	cag	cca	caa	aaa	tgt	gta	tgt	tca	tgt	atg	tac	tgt
att	ttc	ogt	ttg	ggt	ata	tgt	gct	gca	caa	ggg	ctt	goc	co						

FIG. 2

FIG. 3




FIG. 4

Hs	1	-----KAVPFVED--MDLVOTLGEGAYGEVOLAVNRVT-ERAVAVKIVDNK-----RAVOCF
Dm	1	MAATLTAQTGPAATREFVEG--WTLAOTLGEGAYGEVLLINROT-GCGCCNKIVDLK-----KIPDAA
Ce	1	MSAASTTSTPAAAAVAPOQPESLYRVVOTLGEGAFGEVLLIVNTKPEVVAANKIINIAN-----KSKDFI
Sp	1	-----MAQKLDNF PYHIGREIGTGAFAEVRLCYDDNA--RITAVKFEVKKHATSCKMAGVWA
Hs	50	ENIKREICINKMLN--NENWKFYGRREGNI CYLFLEYCSGGELFDRIEPDIGKPEPDAQFFHQLNAG
Dm	63	NSVKEVCIQKMLQ---DNVILRF GARSQGSVYLFLEYAAGGELFDRIEPDVGNPQHEAORYFTOLLG
Ce	67	DNIRREYLLQKRVSAVGNNDVIRIGMRNDPQFYLFLEYADGGELFDKIEPDCGSEPVTAQFYKQLICG
Sp	54	RRTASEIQLEKLCN--GNKNIHFYNTAENFQWVWVLEFAQGGDLFDKIEPDVGLDEDVAQFYEAQLREG
Hs	118	VVYLH-GIGIHRDIKPENLLLDEHDNLKISDFGLATVFRY--FRERLLNKCGTLPYVAPELLKRRE--E
Dm	131	LVYLH-QEGIAHRDIKPENLLLDEHDNVKISDFGHATMFRCK-CERLLDRCGTLPYVAPEVLOKAY--Q
Ce	138	LKFIH-DNDYVHRDIKPENLLLGTTEVLKISDFGHATLYRCK-CERLLDLEGGTIPYAAPELCAGKK--YR
Sp	125	ISEFN-SIGYANRDLKPENILLDYNONLKISDFGASLFSYK-GSRLLNSPVGSPRYAPEITQOYD---
Hs	186	AEPVDVWSCGIVLTANLAGELFWDPQSDSCOEYSDKKEKTYCN--PWKKIDSAPLALLKILVENPESARZ
Dm	199	PCPADLWSCGVILVTHLAGELFWDPQSTNCTEPTNRDDEWLOTPKSKIEDTLAISLDRKILLATSPGTR
Ce	206	GEPVDVWSEGIIVLIANLTGELFWDRASDASOSYNGC-IISNTSLDER-PWKKIDVVALCHLRKIVTDTKRA
Sp	191	GSVVDVWSCGIIIFALLGNTPWDEAISNTQDYLLYKQCEKPSYHPNLLSPGAYSLITQELREDFKRY
Hs	255	FIPDLKDRNYYNPLKKGAKRPRVTEGGVSESPSGYSKHIOENLDSPVNSAASSEAVKYSSSQIPEPTOL
Dm	270	TEKTLDERVCMQFADNERSYDLEDSAAALEYCSPEANRORLQSSAMENGLDSEIEGNYCSQPMPTMRT
Ce	277	FICQIQADPHYCH-----NFGQV-ETPNGRFUKRANNE-----EHTICTQDAECSAK
Sp	262	SVKHYVQEPHLETSSTPRTKNGCADPVALASRLNLELRIDLD-KPRLASSRASQNDGQSMTOPAKKND
Hs	326	SDMTSPKIDKIV-----QDISFSOPTCFDEMLNSQLLQTPQSSQNPQRLVK
Dm	341	DDDFNVRLGSGRIQ-----GCWRRPQTLAQEARLEYSPFSQPALLODLLLATQDQCTQWASQNTQRLVR
Ce	335	RR-RLTPNEXSSLAER-----QVASEFSOPTKEDLLTQHLIDHSQMS-NLQRMVC
Sp	332	QKELDRVSVYGLSQPV-----QLAKNIDVTEILKDPSSLQFCNEQFI-----KRLAK
Hs	376	R-----ITRFFETK--LDADKSYQCLKETCSLGYQNKASCNNQVTISTDRRNKLIKVNILLE
Dm	405	R-----ITRFFVTRWDQTI-KRLVOTIERLGGYTCYQDDGVTVSTVDROKLRLLVFKANIEE
Ce	374	R-----ITRFQVV--TDIRSTYQKVARASEKAGFOLRITDDYRQVLTIREVSNVSLYTDQDIP
Sp	371	KAKNEYEICFERTREYSR-----ASRETLIDHLYDSLRLLAISVTDIVRN-OTILYVNLHD
Hs	433	VDCKIL-----VDFRLSKGDCLEFKRFFLKIKKLLID--VSSCAVWLPAI----- 476
Dm	463	VDCKIL-----VDCRLSKGCGLEFKRFFIKIKKNALEDIVLQKPTTNP IAIATNSVP 513
Ce	431	DAPRVK-----VDFRSSRGDGIQPKKMFMDVRMRMHEMCTOSDILDMCQEIRK-- 479
Sp	440	KKXQAL-QGVLELTMLQHNLELINFIKENGDPLENKVFKNVVSQIKKPLVQDQVSN----- 496




FIG. 5A




FIG. 5B




FIG. 6




FIG. 7A




FIG. 7B




FIG. 8A




FIG. 8B




FIG. 9A




FIG. 9B




FIG. 9C



FIG. 9D




FIG. 10




FIG. 11




FIG. 12A




FIG. 12B

**MAMMALIAN CHECKPOINT PROTEINS
POLYPEPTIDES AND ENCODING
SEQUENCES THEREOF**

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.

This application is a Division Application of U.S. patent application Ser. No. 08/924,183 filed on Sep. 5, 1997, now U.S. Pat. No. 6,218,109.

FIELD OF THE INVENTION

The present invention relates to mammalian proteins and gene sequences involved in cellular responses to DNA damage. In particular, the present invention provides checkpoint genes and proteins.

BACKGROUND OF THE INVENTION

The proper development of a multicellular organism is a complex process that requires precise spatial and temporal control of cell proliferation. Cell proliferation in the embryo is controlled via an intricate network of extracellular and intracellular signaling pathways that process growth regulatory signals. This signaling network is superimposed upon the basic cell cycle regulatory machinery that controls particular cell cycle transitions.

Cell cycle checkpoints are regulatory pathways that control the order and timing of cell cycle transitions, and ensure that critical events such as DNA replication and chromosome segregation are completed with high fidelity. For example, proliferating eukaryotic cells arrest their progression through the cell cycle in response to DNA damage. This arrest is critical to the survival of the organism, as failure to repair damaged DNA can result in the formation and transfer of mutations, damaged chromosomes, cancer, or other detrimental effects. The mechanism responsible for monitoring the integrity of the organism's DNA and preventing the progression through the cell cycle when DNA damage is detected is referred to as the "DNA damage checkpoint."

In response to DNA damage, cells activate a checkpoint pathway that arrests the cell cycle, in order to provide time for repair, and induces the transcription of genes that facilitate the needed repair. In yeast, this checkpoint pathway consists of several protein kinases including phosphoinositide (Pt)-kinase homologs hATM (human), scMec1 (*Saccharomyces cerevisiae*), spRad3 (*Schizosaccharomyces pombe*), and protein kinases scDun1 (*Saccharomyces cerevisiae*), scRad53 (*Saccharomyces cerevisiae*), and spChk1 (*Schizosaccharomyces pombe*) (See e.g., S. Elledge, *Science* 1664 [1996])

Indeed, the ability to coordinate cell cycle transitions in response to genotoxic and other stressors is critical to the maintenance of genetic stability and prevention of uncontrolled cellular growth. Loss of a checkpoint gene leads to genetic instability and the inability of the cells to deal with genomic insults such as those suffered as a result of the daily exposure to ultraviolet radiation. The loss of negative growth control and improper monitoring of the fidelity of DNA replication are common features of tumor cells. When checkpoints are eliminated (e.g., by mutation or other means), cell death, infidelity in chromosome transmission, and/or increased susceptibility to deleterious environmental factors (e.g., DNA-damaging agents) result. A variety of abnormal cells arising due to infidelity during mitoses have been

detected in humans, including aneuploidy, gene amplification, and multipolar mitoses (See, L. H. Hartwell and T. A. Weinert, *Science* 245:629 [1989]).

Accordingly, elucidation of checkpoint function, as well as the disruption of checkpoint function, will further the understanding of the process of cellular transformation (i.e., the conversion of normal cells to a state of unregulated growth), as well as cell differentiation and organismal development.

SUMMARY OF THE INVENTION

The present invention provides mammalian proteins and gene sequences involved in cellular responses to DNA damage. In particular, the present invention provides Chk1 genes and proteins.

In one embodiment, the present invention provides the nucleotide sequence set forth in SEQ ID NO:1. In alternative embodiments, the present invention provides SEQ ID NO:1, wherein it further comprises 5' and 3' flanking regions. In yet another embodiment, the sequence further comprises intervening regions. In a further embodiment, the present invention also provides a polynucleotide sequence which is complementary to SEQ ID NO:1 or variants thereof. In a preferred embodiment, the present invention provides a vector comprising the nucleotide sequence of claim 1. The present invention also provides host cell(s) containing the vector of claim 4.

The present invention also provides a purified Chk1 protein encoded by the nucleotide sequence of claim 1, as well as a purified protein comprising the amino acid sequence set forth in SEQ ID NO:3. In addition, the present invention provides fusion proteins comprising a least a portion of the human Chk1 protein, as well as non-Chk1 protein sequences. It is not intended that the fusion proteins of the present invention be limited to any particular portion of the Chk1 portion or any particular non-Chk1 protein sequences. In preferred embodiments, the fusion the Chk1 protein portion of the fusion protein comprises at least a portion of SEQ ID NO:3. In an alternative embodiment, the non-Chk1 protein sequence comprises an affinity tag. In particularly preferred embodiment, the affinity tag comprises a histidine tract.

In yet another embodiment, the present invention provides the sequence set forth in SEQ ID NO:2. In an alternative embodiment, the nucleotide sequence further comprises 5' and 3' flanking regions. In another alternative embodiment, the nucleotide sequence further comprises intervening regions. In yet another embodiment, the present invention provides a polynucleotide sequence that is complementary to SEQ ID NO:2 or variants thereof.

The present invention also provides a vector comprising the nucleotide sequence set forth in SEQ ID NO:2. In one preferred embodiment, the present invention provides a host cell containing the vector comprising this nucleotide sequence.

The present invention further provides a purified Chk1 protein encoded by the nucleotide sequence of SEQ ID NO:2. In yet another embodiment, the present invention provides a purified protein comprising the amino acid sequence set forth in SEQ ID NO:4.

The present invention also provides fusions proteins comprising at least a portion of the murine Chk1 protein and a non-Chk1 protein sequence. It is not intended that the fusion proteins of the present invention be limited to any particular portion of the Chk1 portion or any particular non-Chk1 protein sequences. In preferred embodiments, the fusion the

Chk1 protein portion of the fusion protein comprises at least a portion of SEQ ID NO:4. In an alternative embodiment, the non-Chk1 protein sequence comprises an affinity tag. In particularly preferred embodiment, the affinity tag comprises a histidine tract.

The present invention also provides methods for detecting Chk1 protein. In one embodiment, the method comprises the steps of providing in any order: a sample suspected of containing the Chk1 protein; an antibody capable of specifically binding to a Chk1 protein; mixing the sample and the antibody under conditions wherein the antibody can bind to the Chk1 protein; and detecting the binding. In one alternative embodiment, the sample comprises one or more cells suspected of containing Chk1 protein. In yet another embodiment, the cells contain an abnormal Chk1 protein. In a further embodiment, the cells are selected from the group consisting of human cells and murine cells.

The present invention also provides antibodies capable of recognizing at least a portion of human and/or murine Chk1 protein. In one embodiment, the present invention provides an antibody, wherein the antibody is capable of specifically binding to at least one antigenic determinant on the proteins encoded by an amino acid sequence selected from the group comprising SEQ ID NOS:3, 4, 7, 8, 9, and 10. In one preferred embodiment, the antibody is a polyclonal antibody, while in an alternative embodiment, the antibody is a monoclonal antibody.

The present invention also provides methods for producing antibodies comprising the steps of providing in any order: an antigen comprising at least a portion of Chk1 protein; and an animal having immunocompetent cells; and exposing the animal to the Chk1 protein under conditions such that the immunocompetent cells produce anti-Chk1 antibodies. In one alternative embodiment, the method further comprises the step of harvesting the antibodies. In another alternative embodiment, the antigen comprises at least a portion of Chk1 protein is a fusion protein. In yet another embodiment, the method further comprises the step of fusing the immunocompetent cells with an immortal cell line under conditions such that a hybridoma is produced.

The present invention also provides methods for detection of polynucleotides encoding human and/or murine Chk1 in biological samples. It is not intended that the method be limited to any particular sequence contained within SEQ ID NOS:1 or 2. Indeed, it is contemplated that any sequence be used in the method, including degenerate primers that are based on the sequence of [chk1] *CHK1* and that are capable of recognizing at least a portion of the [chk1] *CHK1* gene.

In one embodiment, the method comprises the steps of hybridizing a nucleotide comprising at least a portion of the nucleotide of SEQ ID NO:1 to nucleic acid material of a biological sample, thereby forming a hybridization complex; and detecting the hybridization complex, wherein the presence of the complex correlates with the presence of a polynucleotide encoding human Chk1 in the biological sample.

In an alternative embodiment of the method, the biological sample is amplified by the polymerase chain reaction before hybridization. In yet another alternative method, polymerase chain reaction is conducted using primers selected from the group consisting of SEQ ID NOS:5, 6, 12, 13, 14, and 15.

DESCRIPTION OF THE FIGURES

FIG. 1 shows the cDNA sequence of human [chk1] *CHK1* (SEQ ID NO:1), as well as the predicted amino acid sequence (SEQ ID NO: 3).

FIG. 2 shows the cDNA sequence of murine [chk1] *Chk1* (SEQ ID NO:2), as well as the predicted amino acid sequence (SEQ ID NO: 4).

FIG. 3 shows the domain structure of the predicted human Chk1.

FIG. 4 shows the alignment of human, *D. melanogaster*, *C. elegans*, and *S. pombe* Chk1 homologs.

FIG. 5A shows the expression pattern of human [Chk1] *CHK1* mRNA in different adult tissues, as determined by Northern analysis.

FIG. 5B shows the expression pattern of murine [Chk1] *Chk1* mRNA in different adult tissues, as determined by Northern analysis.

FIG. 6 shows the reactivity of a 54 kD protein from different cell lines, with purified antibodies directed to a peptide (anti-PEP), or full-length Chk1 protein anti-FL).

FIG. 7A shows that Chk1 is modified in response to DNA damage in HeLa cells.

FIG. 7B shows that Chk1 is modified in response to DNA damage in Jurkat cells.

FIG. 8A is an autoradiograph showing radiolabeled phosphate incorporation due to phosphorylation of Chk1, Cdc25A, Cdc25B, and Cdc25C.

FIG. 8B is a Coomassie stained gel of FIG. 8A, showing the amount of each protein that is present in each lane.

FIG. 9A is an autoradiograph showing radiolabeled phosphate incorporation due to phosphorylation of Chk1 and wild-type Cdc25C(200-256).

FIG. 9B is a Coomassie-stained gel of FIG. 9A, showing the amount of each protein present in each lane.

FIG. 9C is an autoradiograph showing radiolabeled phosphate incorporation due to phosphorylation of Chk1 and wild-type Cdc25C.

FIG. 9D is a Coomassie-stained gel of FIG. 9C, showing the amount of each protein present in each lane.

FIG. 10 shows Chk1 binding to GST-Cdc25A, GST-Cdc25B and GST-Cdc25C by immunoblotting with anti Chk1 antibodies.

FIG. 11 shows the proteolytic sites surrounding serine 216 on Cdc25C.

FIG. 12A shows the results of mapping of the phosphorylation site on Cdc25C by Chk1 by proteolytic cleavage and sequencing of phosphorylated Cdc25C, as resolved by HPLC.

FIG. 12B provides identification of phosphorylated residue by manual Edman degradation of tryptic peptide present in fraction 57 shown in FIG. 12A.

DESCRIPTION OF THE INVENTION

The cell cycle comprises a collection of highly ordered processes that lead to the duplication of cells. As cells move through the cell cycle, they undergo several discrete transitions (i.e., an unidirectional change of state in which a cell that was performing one set of processes shifts its activity to perform a different set of processes). Although the mechanism of how these transitions are coordinated to occur at precise times and in a defined order remains unknown, in principle, the ordering of cell cycle events may be accomplished by requiring the next event to physically require the completion of the previous event. This pathway has been referred to as a "substrate-product relationship" (Hartwell and Weinert, *Science* 246:629 [1989]). However, other research has shown that the predominant mechanism for dependency relies upon positive or negative regulatory circuits.

These regulatory circuits are surveillance mechanisms that monitor the completion of critical cell cycle events, and allow the occurrence of subsequent cell cycle transitions. Two classes of regulatory circuits have been described “Intrinsic” mechanism act in each cell cycle to order events, while “extrinsic” mechanisms are induced to act only when a defect is detected. Both of these mechanisms may use the same components to enforce cell cycle arrest. These pathways are particularly important, as their loss leads to reduced fidelity of cell cycle events such as chromosome duplication and segregation. Also, such alterations decrease the reproductive fitness of unicellular organisms, and may lead to uncontrolled cellular proliferation and cancer in multi-cellular organisms.

The term “checkpoint” is used to refer to particular subsets of these intrinsic and extrinsic mechanisms. As used herein, a “checkpoint” is a biochemical pathway that ensures dependence of one process upon another process that is otherwise biochemically unrelated. As a null allele in a checkpoint gene results in a loss of this dependency, checkpoints are inhibitory pathways. This definition is broad, and can apply to many situations that occur in multicellular organisms, particularly during development. However, it is often used in reference to control of cell cycle transitions. In preferred embodiments, the term refers to the biochemical pathway that ensures dependency. For example, the DNA-damage checkpoint is the mechanism by which damaged DNA is detected and a signal is generated that arrests cells in the G1 phase of the cell cycle, slows down S phase (i.e., DNA synthesis), arrests cells in the G2 phase, and induces the transcription of repair genes. The position of arrest within the cell cycle varies, depending upon the phase in which the damage is determined. Whether the loss of a checkpoint has an immediate consequence for an organism during a normal cell cycle depends upon the particular pathway involved and the inherent timing of the processes. Thus, timing and checkpoints can act as redundant controls, in order to ensure the proper order of events. Therefore, there are no constraints on whether checkpoints are essential or inducible (extrinsic).

To address the conservation of checkpoint function, a search for human homologs of yeast checkpoint genes was conducted using a degenerate polymerase chain reaction (PCR) strategy. This search identified a human gene very similar to [spChk1] *spchk1* (See, FIG. 3). Using [hChk1] *hCHK1* cDNA (SEQ ID NO:1) as a probe, the [mChk1] *Chk1* gene from mouse (SEQ ID NO:2) was isolated. The sequence of the longest human cDNA (1891 bp) predicted a translation product of 476 amino acids with a molecular size of 54 kD (FIG. 3). No in-frame stop codon was found upstream of the first methionine, which is within the Kozak consensus sequence, and is likely to be the bone fide initiation codon because its encoded protein is the same size as that observed in cells (see below). [hChk1] *hCHK1* was found to be related to a [C. elegans] *C. elegans* gene in the database and a [D. melanogaster] *D. melanogaster* gene, *grp*, with a role in the cell cycle control and development (See e.g., FIG. 3). The predicted [hChk1] *hCHK1* is 29% identical and 44% similar to [spChk1] *spchk1*, 40% identical and 56% similar to the [ceChk1] *ceCHK1*, and 44% identical and 56% similar to [dmChk1] *dmCHK1*. Sequence analysis revealed several COOH-terminal domains that are highly conserved in the Chk1 family of kinases (See e.g., FIG. 4).

The chromosomal location of [hChk1] *hChk1* was then mapped to 11q24 by fluorescence in situ hybridization. This site is adjacent to the ATM gene at 11q23. This was of interest as ATM is mutated in patients with ataxia telangiectasia, a

fatal disease characterized by autosomal recessive inheritance, immunological impairment, ataxia related to progressive cerebellar Purkinje cell death, and a high incidence of cancer (See e.g., L. S. Cox and D. P. Lane, *Bioessays* 17:501 [1995]; Y. Shiloh et al., *J. Hum. Genet.*, 3:116 [1995]; K. Savitsky et al., *Science* 268:1749 [1995]); and K. Savitsky et al., *Hum. Mol. Genet.*, 4:2025 [1995]). Approximately 1% of humans are heterozygotic for ATM defects, and show an increased incidence of cancer (See, M. Swift et al., *N. Eng. J. Med.*, 316:1289 [1987]; and M. Swift et al., *N. Eng. J. Med.*, 325:1831 [1991]).

Northern blot analysis revealed ubiquitous expression with large amounts of Chk1 expression in human thymus, testis, small intestine and colon (FIG. 5). In this analysis, blots containing the polyadenylated RNA from the indicated tissues were probed with human *CHK1* (FIG. 5A) or mouse *Chk1* (FIG. 5B) [chk1] cDNAs. As shown in FIG. 5B, in adult mice, [mChk1] *Chk1* was detected in all tissues examined and in large amounts in the testis, spleen, and lung. In addition, mouse embryos revealed ubiquitous expression, with large amounts detected in the brain, liver, kidney, pancreas, intestines, thymus and lung. This was of interest as testis, spleen and thymus also express large amounts of ATM (G. Chen and E. Y. H. P. Lee, *J. Biol. Chem.*, 271:33693 [1996]; N. D. Lakin et al., *Oncogene* 13:2707 [1996]).

Affinity purified antibodies to hChk1 protein (GST fusion hChk1 protein) made in baculovirus (anti-FL) or to its COOH-terminal 15 amino acids (anti-PEP) recognized a 54-kD protein (FIG. 6) that comigrates with hChk1 expressed in baculovirus. As shown in this Figure, the anti-PEP, but not anti-FL signal is competed by addition of excess peptide, indicating that the two sera are recognizing different hChk1 epitopes, further confirming identity of the 54-kD band as endogenous hChk1. A 70 kD protein was also specifically recognized by anti-PEP.

When mChk1 was expressed from the cytomegalovirus (CMV) promoter in baby hamster kidney (BHK) cells, a 54 kD nuclear protein was detected only in transfected cells using antibodies directed against the C-terminal peptide of mChk1. This endogenous mChk1 was found to comigrate with endogenous mChk1 from mouse lung tissue.

To determine whether hChk1 is modified in response to DNA damage like spChk1, hChk1 protein in extracts from cells treated with ionizing radiation was examined. hChk1 from extracts obtained from damaged cells showed a minor but reproducible reduction in mobility compared to hChk1 from untreated cells (See, FIG. 7). The mobility alteration observed in response to DNA damage for spChk1 was also slight, as previously reported (See e.g., N. C. Walworth et al., *Nature* 363:368 [1993]; Al-Khodairy et al., *Mol. Biol. Cell* 5:147 [1994]; and N. C. Walworth and R. Bernards, *Science* 271:353 [1996]). This modification was confirmed by 2-dimensional gel analysis, which clearly demonstrated the generation of a more negatively charged hChk1 species 2 hours after γ -irradiation (See, FIG. 7). These results indicate that like spChk1, hChk1 may participate in transduction of the DNA damage signal. Indirect immunofluorescence revealed that hChk1 is localized to the nucleus in a punctate staining pattern, similar to that observed for ATM (See, G. Chen and E. Y. H. P. Lee, *J. Biol. Chem.*, 271:33693 [1996]; and N. D. Lakin et al., *Oncogene* 13:2702 [1996]). mChk1 expressed in BHK cells confirmed the nuclear localization.

To test for the ability of hChk1 to regulate the cell cycle, hChk1 or hChk1 (D130A) (a catalytically inactive mutant), were transfected under the control of the CMV promoter, or the CMV vector alone into HeLa cells treated with and with-

out 6 Gy of ionizing radiation. No perturbation of the cell cycle by either kinase relative to vector alone was observed, suggesting that overproduction alone was insufficient to deregulate the system.

Tyrosine phosphorylation of Cdc2 has been implicated in cell cycle arrest in response to DNA damage and replication blocks in both *S. pombe* (T. Enoch and P. Nurse, *Cell* 60:665 [1990]) and humans (P. Jin et al., *J. Cell. Biol.*, 134:963 [1996]). In *S. pombe*, Cdc2 mutants that cannot be phosphorylated on tyrosine display an inability to arrest the cell cycle in response to blockade of DNA replication. Although it was originally thought that the DNA damage checkpoint did not operate through tyrosine phosphorylation, recent experiments have shown that tyrosine phosphorylation is required for *S. pombe* cells to arrest in response to DNA damage (P. Jin et al., *J. Cell. Biol.*, 134:963 [1996]). While it is now clear that tyrosine phosphorylation is required for proper checkpoint control, the experiments implicating tyrosine phosphorylation in this pathway do not distinguish between a regulatory role in which tyrosine phosphorylation rates are manipulated by the checkpoint pathways, or a passive role in which tyrosine phosphorylation is required to allow cell cycle arrest, but is not the actual target of the checkpoint pathway (S. Elledge, *Science* 274:1664 [1996]; and D. J. Lew and S. Kombluth, *Curr. Opin. Cell. Biol.*, 8:795 [1996]).

To address this issue, the ability of hChk1 to phosphorylate key regulators of Cdk tyrosine phosphorylation was examined. For these experiments, the Cdc25 dual-specificity phosphatases, hCdc25A, hCdc25B, and hCdc25C were analyzed. These regulators were chosen for several reasons. First, overproduction of hChk4 mutants in which the inhibitory tyrosine is changed to phenylalanine abrogates G1 arrest in response to UV light (Y. Terada et al., *Nature* 376:358 [1995]). Secondly, the UV-sensitivity of chk1⁻ mutants in *S. pombe* is suppressed by inactivating cdc25 with a Ts (i.e., temperature sensitive) mutation (N. C. Walworth et al., *Nature* 363:368 [1993]). Finally, in *S. pombe* wee1mik1 mutants, DNA damage still causes a partial cell cycle delay that could be due to regulation of spCdc25 activity. GST-hChk1 and GST-hChk1 (D130A) were introduced into baculovirus, purified from baculovirus-infected insect cells, and incubated with GST-hCdc25A, GST-hCdc25B, and GST-hCdc25C, as described in Example 5. GST-hChk1 phosphorylated all three Cdc25 proteins, but GST alone did not (See, FIG. 8). Although GST-hCdc25C co-migrated with GST-hChk1 (which autophosphorylates), increased phosphorylation was observed at that position relative to phosphorylation in the presence of kinase alone, and phosphorylation of a GST-hCdc25C breakdown product was visible. In separate experiments using a His₆-tagged hChk1 derivative, there was clear phosphorylation of GST-hCdc25C (FIG. 9). A catalytically inactive mutant failed to phosphorylate itself or any of the Cdc25 proteins (See, FIG. 8).

Protein kinases often form complexes with their substrates. To examine whether this occurs with hChk1 and the Cdc25 proteins, GST-hCdc25 proteins on glutathione beads were incubated together with baculovirus extracts expressing His₆-tagged hChk1, and precipitated. GST-hCdc25A, GST-hCdc25B and GST-hCdc25C, each specifically bound hChk1, while GST alone did not (See, FIG. 10). Furthermore, two other GST fusion proteins, GST-Dun1 and GST-Skp1, all failed to bind hChk1. These results indicate that Cdc25 can form complexes with hChk1.

To establish the significance of the Cdc25 phosphorylation, the site of hChk1 phosphorylation on Cdc25C was mapped. It was determined that Ser²¹⁶ is the

main site of phosphorylation of hCdc25C in vivo. hChk1 phosphorylated a 56 amino acid region of the hCdc25C protein fused to GST, but not GST alone (FIGS. 8 and 9). This 56 amino acid motif contains four possible sites of phosphorylation. Peptide analysis of proteolytic fragments of full length His₆-hCdc25C phosphorylated with GST-hChk1 revealed a single phosphorylated tryptic peptide by high pressure liquid chromatography. Edman degradation of this peptide indicated release of radioactivity in the third cycle (FIG. 12B). Further degradation of this tryptic fragment with proline endopeptidase resulted in production of a peptide that released radioactivity in the first cycle. The Serine²¹⁶ residue is the only site on hCdc25C that is consistent with this phosphorylation pattern (FIG. 11). To confirm this, the Cdc25C S216A mutation in GST-Cdc25C and Cdc25C (200-256) were constructed. It was determined that both were poor substrates for hChk1, confirming that Serine²¹⁶ is the site phosphorylation (FIG. 9). Serine 216 is also phosphorylated by spChk1, demonstrating phylogenetic conservation of this regulatory relationship.

The present invention provides evidence that the Chk1 kinase family is conserved throughout eukaryotic evolution and that hChk1, like its *S. pombe* counterpart, is modified in response to DNA damage. This, together with the fact that ATM-related kinases are conserved members of checkpoint pathways and act upstream of chk1 in *S. pombe*, suggests that this entire checkpoint pathway may be conserved in all eukaryotes. Nonetheless, the present invention provides the first mammalian Chk1 sequences.

hChk1 directly phosphorylates a regulator of Cdc2 tyrosine phosphorylation, hCdc25C, on a physiologically significant residue, Serine²¹⁶. In addition, overexpression of the hCdc25C S216A protein reduces the ability of cells to arrest in G2 in response to DNA damage, as observed previously for the Cdc2AF mutants. However, the overexpression studies alone do not prove that the DNA damage checkpoint pathway operates through tyrosine phosphorylation, because hyperactive Cdc2 may be able to bypass checkpoint control. Nonetheless, in combination with the fact that this inhibitory serine is directly phosphorylated by the DNA damage-responsive checkpoint kinase hChk1, these results strongly imply that DNA damage regulates the G2-to-mitosis transition through control of Cdc2 tyrosine phosphorylation. These results suggest a model whereby in response to DNA damage, hChk1 phosphorylates hCdc25C on Serine²¹⁶, leading to binding of 14-3-3 protein and inhibition of Cdc25C's ability to dephosphorylate and activate Cdc2. This model does not preclude a role for other cell cycle regulators such as Wee1 in the damage response. Although an understanding of the mechanism is not necessary in order to use the present invention, the facts that hChk1 phosphorylated hCdc25A and hCdc25B, and that Serine²¹⁶ is conserved among these Cdc25 proteins, suggests that hChk1 may regulate other DNA damage checkpoints, such as those controlling the G1 to S phase transition, through a similar mechanism.

DEFINITIONS

To facilitate understanding of the invention, a number of terms are defined below.

“Nucleic acid sequence” as used herein refers to an oligonucleotide, nucleotide, or polynucleotide, and fragments of portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense strand. Similarly, “amino acid sequence” as used herein refers to an

oligopeptide, peptide, polypeptide, or protein sequence, and fragments or portions thereof, and to naturally occurring or synthetic molecules.

A "composition comprising a given polynucleotide sequence" as used herein refers broadly to any composition containing the given polynucleotide sequencer. The composition may comprise an aqueous solution. Compositions comprising polynucleotide sequences encoding human or murine Chk1 (SEQ ID NOS:3 and 4) or fragments thereof, may be employed as hybridization probes. In this case, the human and murine chk1-encoding polynucleotide sequences are typically employed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., SDS) and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

Where "amino acid sequence" is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms, such as "polypeptide" or "protein" are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

"Peptide nucleic acid," as used herein, refers to a molecule which comprises an oligomer to which an amino acid residue, such as lysine, and an amino group have been added. These small molecules, also designated anti-gene agents, stop transcript elongation by binding to their complementary strand of nucleic acid (Nielsen, P. E. et al., *Anticancer Drug Des.* 8:53-63 [1993]).

Chk1 as used herein, refers to the amino acid sequences of substantially purified Chk1 obtained from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and human, from any source whether natural, synthetic, semi-synthetic, or recombinant. In particularly preferred embodiments, Chk1 is human or murine.

"Consensus," as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, or which has been extended using any suitable method known in the art, in the 5' and/or the 3' direction and resequenced, or which has been assembled from the overlapping sequences of more than one clone using any suitable method known in the art, or which has been both extended and assembled.

A "variant" of human or murine Chk1, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties (e.g., replacement of leucine with isoleucine). More rarely, a variant may have "nonconservative" changes (e.g., replacement of a glycine with a tryptophan). Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTAR® software (*DNASTAR, Inc. Madison, Wis.*).

A "deletion," as used herein, refers to a change in either amino acid or nucleotide sequence in which one or more amino acid or nucleotide residues, respectively, are absent.

An "insertion" or "addition," as used herein, refers to a change in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid or nucleotide residues, respectively, as compared to the naturally occurring molecule.

A "substitution," as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

The term "biologically active," as used herein, refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic human or murine Chk1, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "modulate," as used herein, refers to a change or an alteration in the biological activity of human or murine Chk1. Modulation may be an increase or a decrease in protein activity, a change in binding characteristics, or any other change in the biological, functional, or immunological properties of human or murine Chk1.

The term "mimetic," as used herein, refers to a molecule, the structure of which is developed from knowledge of the structure of Chk1, or portions thereof and, as such, is able to effect some or all of the actions of human or murine Chk1-like molecules.

The term "derivative," as used herein, refers to the chemical modification of a nucleic acid encoding human or murine Chk1, or the encoded human or murine Chk1 protein. Illustrative of such modifications would be replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would encode a polypeptide which retains essential biological characteristics of the natural molecule.

The term "substantially purified," as used herein, refers to nucleic or amino acid sequences that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components with which they are naturally associated.

The term "hybridization," as used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing. Hybridization and the strength of hybridization (i.e., the strength of the association between the nucleic acids) is impacted by such factors as the degree of complementarity between the nucleic acids, stringency of the conditions involved, the T_m of the formed hybrid, and the G:C ratio within the nucleic acids.

As used herein, the term " T_m " is used in reference to the "melting temperature." The melting temperature is the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single strands. The equation for calculating the T_m of nucleic acids is well known in the art. As indicated by standard references, a simple estimate of the T_m value may be calculated by the equation: $T_m = 81.5 + 0.41(\% G+C)$, when a nucleic acid is in aqueous solution at 1 M NaCl (see e.g., Anderson and Young, *Quantitative Filter Hybridization, in Nucleic Acid Hybridization* (1985). Other references include more sophisticated computations which take structural as well as sequence characteristics into account for the calculation of T_m .

The term "hybridization complex," as used herein, refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., C_{ot} or R_{ot} analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., membranes, filters, chips, pins or glass slides to which cells have been fixed for in situ hybridization).

The terms "complementary" or "complementarity," as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, for the sequence "A-G-T" binds to the complementary sequence "T-C-A". Complementarity between two single-stranded molecules may be "partial", in which only some of the nucleic acids bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands.

The term "homology," as used herein, refers to a degree of complementarity. There may be partial homology or complete homology (i.e., identity). A partially complementary sequence is one that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid; it is referred to using the functional term "substantially homologous." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding (i.e., the hybridization) of a completely homologous sequence or probe to the target sequencer under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (i.e., selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% identity); in the absence of non-specific binding, the probe will not hybridize to the second non-complementary target sequences. When used in reference to a single-stranded nucleic acid sequence, the term "substantially homologous" refers to any probe which can hybridize (i.e., it is the complement of) the single-stranded nucleic acid sequence under conditions of low stringency as described.

As known in the art, numerous equivalent conditions may be employed to comprise either low or high stringency conditions. Factors such as the length and nature (DNA, RNA, base composition) of the sequence, nature of the target (DNA, RNA, base composition, presence in solution or immobilization, etc.), and the concentration of the salts and other components (e.g., the presence or absence of formamide, dextran sulfate and/or polyethylene glycol) are considered and the hybridization solution may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions. As used herein the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds such as organic solvents, under which nucleic acid hybridizations are conducted. With "high stringency" conditions, nucleic acid base pairing will occur only between nucleic acid fragments that have a high frequency of complementary base sequences. Thus, conditions of "weak" or "low" stringency are often required with nucleic acids that are derived from organisms that are genetically diverse, as the frequency of complementary sequences is usually less.

Low stringency conditions comprise conditions equivalent to binding or hybridization at 42° C. in a solution consisting of 5×SSPE (43.8 g/l NaCl, 6.9 g/l NaH₂PO₄·H₂O and

1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.1% SDS, 5×Denhardt's reagent (50×Denhardt's contains per 500 ml; 5 g [Ficoll] *FICOLL* (Type 400, [Pharmacia] *GE Healthcare Bio-Sciences AB LTD, Uppsala, Sweden*), 5 g BSA [Fraction V; [Sigma] *SIGMA-ALDRICH BIOTECHNOLOGY L.P., St. Louis, Mo.*]) and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 5×SSPE, 0.1% SDS at 42° C. when a probe of about 500 nucleotides in length is employed.

The art knows well that numerous equivalent conditions may be employed to comprise low stringency conditions; factors such as the length and nature (DNA, RNA, base composition) of the probe and nature of the target (DNA, RNA, base composition, present in solution or immobilized, etc.) and the concentration of the salts and other components (e.g., the presence or absence of formamide, dextran sulfate, polyethylene glycol) are considered and the hybridization solution may be varied to generate conditions of low stringency hybridization different from, but equivalent to, the above listed conditions. In addition, the art knows conditions which promote hybridization under conditions of high stringency (e.g., increasing the temperature of the hybridization and/or wash steps, the use of formamide in the hybridization solution, etc.).

The term "antisense," as used herein, refers to nucleotide sequences which are complementary to a specific DNA or RNA sequence. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. Antisense molecules may be produced by any method, including synthesis by ligating the gene(s) of interest in a reverse orientation to a viral promoter which permits the synthesis of a complementary strand. Once introduced into a cell, this transcribed strand combines with natural sequences produced by the cell to form duplexes. These duplexes then block either the further transcription or translation. In this manner, mutant phenotypes may be generated. The designation "negative" is sometimes used in reference to the antisense strand, and "positive" is sometimes used in reference to the sense strand.

The term also is used in reference to RNA sequences which are complementary to a specific RNA sequence (e.g., mRNA). Included within this definition are antisense RNA ("asRNA") molecules involved in gene regulation by bacteria. Antisense RNA may be produced by any method, including synthesis by splicing the gene(s) of interest in a reverse orientation to a viral promoter which permits the synthesis of a coding strand. Once introduced into an embryo, this transcribed strand combines with natural mRNA produced by the embryo to form duplexes. These duplexes then block either the further transcription of the mRNA or its translation. In this manner, mutant phenotypes may be generated. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. The designation (-) (i.e., "negative") is sometimes used in reference to the antisense strand, with the designation (+) sometimes used in reference to the sense (i.e., "positive") strand.

A gene may produce multiple RNA species which are generated by differential splicing of the primary RNA transcript. cDNAs that are splice variants of the same gene will contain regions of sequence identity or complete homology (representing the presence of the same exon or portion of the same exon on both cDNAs) and regions of complete non-identity (for example, representing the presence of exon "A" on cDNA 1 wherein cDNA 2 contains exon "B" instead). Because the two cDNAs contain regions of sequence identity they will both hybridize to a probe derived from the entire gene or portions of the gene containing sequences

found on both cDNAs; the two splice variants are therefore substantially homologous to such a probe and to each other.

The term "portion," as used herein, with regard to a protein (as in "a portion of a given protein") refers to fragments of that protein. The fragments may range in size from four amino acid residues to the entire amino acid sequence minus one amino acid. Thus, a protein "comprising at least a portion of the amino acid sequence of SEQ ID NO:3" encompasses the full-length human Chk1, and fragments thereof.

"Transformation," as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. It may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the host cell being transformed and may include, but is not limited to, viral infection, electroporation, lipofection, and particle bombardment. Such "transformed" cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome.

The term "transfection" as used herein refers to the introduction of foreign DNA into eukaryotic cells. Transfection may be accomplished by a variety of means known to the art including calcium phosphate-DNA co-precipitation, DEAE-dextran-mediated transfection, polybrene-mediated transfection, electroporation, microinjection, liposome fusion, lipofection, protoplast fusion, retroviral infection, and biolistics. Thus, the term "stable transfection" or "stably transfected" refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant" refers to a cell which has stably integrated foreign DNA into the genomic DNA. The term also encompasses cells which transiently express the inserted DNA or RNA for limited periods of time. Thus, the term "transient transfection" or "transiently transfected" refers to the introduction of foreign DNA into a cell where the foreign DNA fails to integrate into the genome of the transfected cell. The foreign DNA persists in the nucleus of the transfected cell for several days. During this time the foreign DNA is subject to the regulatory controls that govern the expression of endogenous genes in the chromosomes. The term "transient transfectant" refers to cells which have taken up foreign DNA but have failed to integrate this DNA.

The term "antigenic determinant," as used herein, refers to that portion of a molecule (i.e., an antigen) that makes contact with a particular antibody (i.e., an epitope). When a protein or fragment of a protein as used to immunize a host animal (e.g., an "immunocompetent" animal with "immunocompetent cells"), numerous regions of the protein may induce the production of antibodies which bind specifically to a given region or three-dimensional structure on the protein; these regions or structures are referred to as antigenic determinants. An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "specific binding" or "specifically binding," as used herein, in reference to the interaction of an antibody and a protein or peptide, mean that the interaction is dependent upon the presence of a particular structure (i.e., the antigenic determinant or epitope) on the protein; in other words, the antibody is recognizing and binding to a specific protein structure rather than to proteins in general. For example, if an antibody is specific for epitope "A", the presence of a protein containing epitope A (or free, unlabeled A)

in a reaction containing labeled "A" and the antibody will reduce the amount of labeled A bound to the antibody.

The term "sample," as used herein, is used in its broadest sense. The term encompasses biological sample(s) suspected of containing nucleic acid encoding human or murine Chk1 or fragments thereof, and may comprise a cell, chromosomes isolated from a cell (e.g., a spread of metaphase chromosomes), genomic DNA (in solution or bound to a solid support such as for Southern analysis), RNA (in solution or bound to a solid support such as for northern analysis), cDNA (in solution or bound to a solid support), an extract from cells or a tissue, and the like.

The term "correlates with expression of a polynucleotide," as used herein, indicates that the detection of the presence of ribonucleic acid that is similar to SEQ ID NO:1 or 2, by Northern analysis is indicative of the presence of mRNA encoding human or murine Chk1, in a sample and thereby correlates with expression of the transcript from the polynucleotide encoding the protein.

"Alterations" in the polynucleotide of SEQ ID NO:1 or 2, as used herein, comprise any alteration in the sequence of polynucleotides encoding human or murine Chk1, respectively, including deletions, insertions, and point mutations that may be detected using hybridization assays. Included within this definition is the detection of alterations to the genomic DNA sequence which encodes human Chk1 (e.g., by alterations in the pattern of restriction fragment length polymorphisms capable of hybridizing to SEQ ID NO:1), the inability of a selected fragment of SEQ ID NO:1 to hybridize to a sample of genomic DNA (e.g., using allele-specific oligonucleotide probes), and improper or unexpected hybridization, such as hybridization to a locus other than the normal chromosomal locus for the polynucleotide sequence encoding human Chk1 (e.g., using fluorescent in situ hybridization [FISH] to metaphase chromosomes spreads).

As used herein, the term "antibody" (or "immunoglobulin"), refers to intact molecules as well as fragments thereof, such as Fa, F(ab')₂, and Fv, which are capable of binding the epitopic determinant. Antibodies that bind human or murine Chk1 polypeptides can be prepared using intact polypeptides or fragments containing small peptides of interest at the immunizing antigen. The polypeptide or peptide used to immunize an animal can be derived from the transition of RNA or synthesized chemically, and can be conjugated to a carrier protein, if desired. Commonly and carriers that are chemically coupled to peptides include bovine serum albumin and thyroglobulin. The coupled peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit).

The term "humanized antibody," as used herein, refers to antibody molecules in which amino acids have been replaced in the non-antigen binding regions in order to more closely resemble a human antibody, while still retaining the original binding ability.

As used herein, the term "poly A⁺ RNA" refers to RNA molecules having a stretch of adenine nucleotides at the 3' end. This polyadenine stretch is also referred to as a "poly-A tail". Eukaryotic mRNA molecules contain poly-A tails and are referred to as poly A⁺ RNA.

As used herein, the term "fusion protein" refers to a chimeric protein containing the protein of interest (i.e., mouse or human [chk1] *Chk1* and fragments thereof) joined to an exogenous protein fragment (the fusion partner which consists of a non-[chk1] *Chk1* protein). The fusion partner may enhance solubility of the [chk1] *Chk1* protein as expressed in

a host cell, may provide an affinity tag to allow purification of the recombinant fusion protein from the host cell or culture supernatant, or both. If desired, the fusion protein may be removed from the protein of interest (i.e., [chk1] *Chk1* protein or fragments thereof) by a variety of enzymatic or chemical means known to the art.

As used herein, the term "affinity tag" refers to such structures as a "poly-histidine tract" or "poly-histidine tag," or any other structure or compound which facilitates the purification of a recombinant fusion protein from a host cell, host cell culture supernatant, or both. As used herein, the term "flag tag" refers to short polypeptide marker sequence useful for recombinant protein identification and purification.

As used herein, the terms "poly-histidine tract" and "poly-histidine tag," when used in reference to a fusion protein refers to the presence of two to ten histidine residues at either the amino- or carboxy-terminus of a protein of interest. A poly-histidine tract of six to ten residues is preferred. The poly-histidine tract is also defined functionally as being a number of consecutive histidine residues added to the protein of interest which allows the affinity purification of the resulting fusion protein on a nickel-chelate or IDA column.

As used herein, the term "chimeric protein" refers to two or more coding sequences obtained from different genes, that have been cloned together and that, after translation, act as a single polypeptide sequence. Chimeric proteins are also referred to as "hybrid proteins" As used herein, the term "chimeric protein" refers to coding sequences that are obtained from different species of organisms, as well as coding sequences that are obtained from the same species of organisms.

As used herein, the term "protein of interest" refers to the protein whose expression as desired within the fusion protein. In a fusion protein, the protein of interest will be joined or fused with another protein or protein domain, the fusion partner, to allow for enhanced stability of the protein of interest and/or ease of purification of the fusion protein.

As used herein, the term "abnormal Chk1 protein" refers to Chk1 that lacks function (i.e., does not function as Chk1 protein in normal cells), or is not recognized by sequences encoding full-length functional Chk1 protein.

As used herein, the term "cell culture" refers to any in vitro culture of cells. Included within this term are continuous cell lines (e.g., with an immortal phenotype), primary cell cultures, finite cell lines (e.g., non-transformed cells), and any other cell population maintained in vitro.

As used herein, the term "selectable marker" refers to the use of a gene which encodes an enzymatic activity that confers the ability to grow in medium lacking what would otherwise be an essential nutrient (e.g., the HIS3 gene in yeast cells); in addition, a selectable marker may confer resistance to an antibiotic or drug upon the cell in which the selectable marker is expressed. Selectable markers may be "dominant", a dominant selectable marker encodes an enzymatic activity which can be detected in any eukaryotic cell line. Examples of dominant selectable markers include the bacterial aminoglycoside 3' phosphotransferase gene (also referred to as the neo gene) which confers resistance to the drug G418 in mammalian cells, the bacterial hygromycin G phosphotransferase (*hyg*) gene which confers resistance to the antibiotic hygromycin and the bacterial xanthine-guanine phosphoribosyl transferase gene (also referred to as the *gpt* gene) which confers the ability to grow in the presence of mycophenolic acid. Other selectable markers are not dominant in that their use must be in conjunction with a cell line that lacks the relevant enzyme activity. Examples of non-

dominant selectable markers include the thymidine kinase (*tk*) gene which is used in conjunction with *tk*⁻ cell lines, the CAD gene which is used in conjunction with CAD-deficient cells and the mammalian hypoxanthine-guanine phosphoribosyl transferase (*hprt*) gene which is used in conjunction with *hprt*⁻ cell lines. A review of the use of selectable markers in mammalian cell lines is provided in Sambrook, J. et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York (1989) pp. 16.9-16.15.

As used herein, the term "vector" is used in reference to nucleic acid molecules that transfer DNA segment(s) from one cell to another. The term "vehicle" is sometimes used interchangeably with "vector."

The term "expression vector" as used herein refers to a recombinant DNA molecule containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular host organism. Nucleic acid sequences necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, often along with other sequences. Eukaryotic cells are known to utilize promoters, enhancers, and termination and polyadenylation signals.

The term "in operable combination," "in operable order," and "operably linked" as used herein refer to the linkage of nucleic acid sequences in such a manner that a nucleic acid molecule capable of directing the transcription of a given gene and/or the synthesis of a desired protein molecule is produced. The term also refers to the linkage of amino acid sequences in such a manner so that a functional protein is produced.

As used herein, the term "amplifiable nucleic acid" is used in reference to nucleic acids which may be amplified by any amplification method. It is contemplated that "amplifiable nucleic acid" will usually comprise "sample template."

As used herein, the term "sample template" refers to nucleic acid originating from a sample which is analyzed for the presence of "target" (defined below). In contrast, "background template" is used in reference to nucleic acid other than sample template which may be or may not be present in a sample. Background template is most often inadvertent. It may be the result of carryover, or it may be due to the presence of nucleic acid contaminants sought to be purified away from the sample. For example, nucleic acids from organisms other than those to be detected may be present as background in a test sample.

As used herein, the term "primer" refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product which is complementary to a nucleic acid strand is induced, (i.e., in the presence of nucleotides and an inducing agent such as DNA polymerase and at a suitable temperature and pH). The primer is preferably single stranded for maximum efficiency in amplification, but may alternatively be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. Preferably, the primer is an oligodeoxyribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the inducing agent. The exact lengths of the primers will depend on many factors, including temperature, source of primer and the use of the method.

As used herein, the term "probe" refers to an oligonucleotide (i.e., a sequence of nucleotides), whether occurring

naturally as in a purified restriction digest or produced synthetically, recombinantly or by PCR amplification, which is capable of hybridizing to another oligonucleotide of interest. A probe may be single-stranded or double-stranded. Probes are useful in the detection, identification and isolation of particular gene sequences. It is contemplated that any probe used in the present invention will be labelled with any "reporter molecule," so that is detectable in any detection system, including, but not limited to enzyme (e.g., ELISA, as well as enzyme-based histochemical assays), fluorescent, radioactive, and luminescent systems. It is not intended that the present invention be limited to any particular detection system or label.

As used herein, the term "target," when used in reference to the polymerase chain reaction, refers to the region of nucleic acid bounded by the primers used for polymerase chain reaction. Thus, the "target" is sought to be sorted out from other nucleic acid sequences. A "segment" is defined as a region of nucleic acid within the target sequence.

As used herein, the term "polymerase chain reaction" ("PCR") refers to the method of K. B. Mullis U.S. Pat. Nos. 4,683,195 4,683,202, and 4,965,188, hereby incorporated by reference, which describe a method for increasing the concentration of a segment of a target sequence in a mixture of genomic DNA without cloning or purification. This process for amplifying the target sequence consists of introducing a large excess of two oligonucleotide primers to the DNA mixture containing the desired target sequence, followed by a precise sequence of thermal cycling in the presence of a DNA polymerase. The two primers are complementary to their respective strands of the double stranded target sequence. To effect amplification, the mixture is denatured and the primers then annealed to their complementary sequences within the target molecule. Following annealing, the primers are extended with a polymerase so as to form a new pair of complementary strands. The steps of denaturation, primer annealing and polymerase extension can be repeated many times (i.e., denaturation, annealing and extension constitute one "cycle"; there can be numerous "cycles") to obtain a high concentration of an amplified segment of the desired target sequence. The length of the amplified segment of the desired target sequence is determined by the relative positions of the primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to as the "polymerase chain reaction" (hereinafter "PCR"). Because the desired amplified segments of the target sequence become the predominant sequences (in terms of concentration) in the mixture, they are said to be "PCR amplified".

With PCR, it is possible to amplify a single copy of a specific target sequence in genomic DNA to a level detectable by several different methodologies (e.g., hybridization with a labeled probe; uncoupling of biotinylated primers followed by avidin-enzyme conjugate detection; incorporation of ³²P-labeled deoxynucleotide triphosphates, such as dCTP or dATP, into the amplified segment). In addition to genomic DNA, any oligonucleotide sequence can be amplified with the appropriate set of primer molecules. In particular, the amplified segments created by the PCR process itself are, themselves, efficient templates for subsequent PCR amplifications.

"Amplification" is a special case of nucleic acid replication involving template specificity. It is to be contrasted with non-specific template replication (i.e., replication that is template-dependent but not dependent on a specific template). Template specificity is here distinguished from

fidelity of replication (i.e., synthesis of the proper polynucleotide sequence) and nucleotide (ribo- or deoxyribo-) specificity. Template specificity is frequently described in terms of "target" specificity. Target sequences are "targets" in the sense that they are sought to be sorted out from other nucleic acid. Amplification techniques have been designed primarily for this sorting out.

Template specificity achieved in most amplification techniques by the choice of enzyme. Amplification enzymes are enzymes that, under conditions they are used, will process only specific sequences of nucleic acid in a heterogenous mixture of nucleic acid. For example, in the case of Q β replicase, MDV-1 RNA is the specific template for the replicase (D. L. Kacian et al., Proc. Natl. Acad. Sci. USA 69:3038 [1972]). Other nucleic acid will not be replicated by this amplification enzyme. Similarly, in the case of T7 RNA polymerase, this amplification enzyme has a stringent specificity for its own promoters (M. Chamberlin et al., Nature 228:227 [1970]). In the case of T4 DNA ligase, the enzyme will not ligate the two oligonucleotides where there is a mismatch between the oligonucleotide substrate and the template at the ligation junction (D. Y. Wu and R. B. Wallace, Genomics 4:560 [1989]). Finally, Taq and Pfu polymerases, by virtue of their ability to function at high temperature, are found to display high specificity for the sequences bounded and thus defined by the primers; the high temperature results in thermodynamic conditions that favor primer hybridization with the target sequences and not hybridization with non-target sequences (H. A. Erlich (ed.), PCR Technology, Stockton Press [1989]).

As used herein, the terms "PCR product," "PCR fragment," and "amplification product" refer to the resultant mixture of compounds after two or more cycles of the PCR steps of denaturation, annealing and extension are complete. These terms encompass the case where there has been amplification of one or more segments of one or more target sequences.

As used herein, the term "amplification reagents" refers to those reagents (deoxyribonucleotide triphosphates, buffer, etc.), needed for amplification except for primers, nucleic acid template and the amplification enzyme. Typically, amplification reagents along with other reaction components are placed and contained in a reaction vessel (test tube, microwell, etc.).

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which are double-stranded DNA at or near a specific nucleotide sequence.

As used herein, the term "recombinant DNA molecule" as used herein refers to a DNA molecule which is comprised of segments of DNA joined together by means of molecular biological techniques.

DNA molecules are said to have "5' ends" and "3' ends" because mononucleotides are reacted to make oligonucleotides in a manner such that the 5' phosphate of one mononucleotide pentose ring is attached to the 3' oxygen of its neighbor in one direction via a phosphodiester linkage. Therefore, an end of an oligonucleotides referred to as the "5' end" if its 5' phosphate is not linked to the 3' oxygen of a mononucleotide pentose ring and as the "3' end" if its 3' oxygen is not linked to a 5' phosphate of a subsequent mononucleotide pentose ring. As used herein, a nucleic acid sequence, even if internal to a larger oligonucleotide, also may be said to have 5' and 3' ends. In either a linear or circular DNA molecule, discrete elements are referred to as being "upstream" or 5' of the "downstream" or 3' elements.

This terminology reflects the fact that transcription proceeds in a 5' to 3' fashion along the DNA strand. The promoter and enhancer elements which direct transcription of a linked gene are generally located 5' or upstream of the coding region. However, enhancer elements can exert their effect even when located 3' of the promoter element and the coding region, Transcription termination and polyadenylation signals are located 3' or downstream of the coding region.

As used herein, the term "an oligonucleotide having a nucleotide sequence encoding a gene" means a nucleic acid sequence comprising the coding region of a gene or in other words the nucleic acid sequence which encodes a gene product. The coding region may be present in either a cDNA, genomic DNA or RNA form. When present in a DNA form, the oligonucleotide may be single-stranded (i.e., the sense strand) or double-stranded. Suitable control elements such as enhancers/promoters, splice junctions, polyadenylation signals, etc. may be placed in close proximity to the coding region of the gene if needed to permit proper initiation of transcription and/or correct processing of the primary RNA transcript. Alternatively, the coding region utilized in the expression vectors of the present invention may contain endogenous enhancers/promoters, splice junctions, intervening sequences, polyadenylation signals, etc. or a combination of both endogenous and exogenous control elements.

As used herein, the term "regulatory element" refers to a genetic element which controls some aspect of the expression of nucleic acid sequences. For example, a promoter is a regulatory element which facilitates the initiation of transcription of an operably linked coding region. Other regulatory elements are splicing signals, polyadenylation signals, termination signals, etc. (defined *infra*).

Transcriptional control signals in eukaryotes comprise "promoter" and "enhancer" elements. Promoters and enhancers consist of short arrays of DNA sequences that interact specifically with cellular proteins involved in transcription (T. Maniatis et al., *Science* 236:1237 [1987]). Promoter and enhancer elements have been isolated from a variety of eukaryotic sources including genes in yeast, insect and mammalian cells and viruses (analogous control elements, i.e., promoters, are also found in prokaryote). The selection of a particular promoter and enhancer depends on what cell type is to be used to express the protein of interest. Some eukaryotic promoters and enhancers have a broad host range while others are functional in a limited subset of cell types (for review see, S. D. Voss et al., *Trends Biochem. Sci.*, 11:287 [1986]; and T. Maniatis et al., *supra*). For example, the SV40 early gene enhancer is very active in a wide variety of cell types from many mammalian species and has been widely used for the expression of proteins in mammalian cells (R. Dijkema et al., *EMBO J.* 4:761 [1985]). Two other examples of promoter/enhancer elements active in a broad range of mammalian cell types are those from the human elongation factor 1 α gene (T. Uetsuki et al., *J. Biol. Chem.*, 264:5791 [1989]; D. W. Kim et al., *Gene* 91:217 [1990]; and S. Mizushima and S. Nagata, *Nuc. Acids. Res.*, 18:5322 [1990]) and the long terminal repeats of the Rous sarcoma virus (C. M. Gorman et al., *Proc. Natl. Acad. Sci. USA* 79:6777 [1982]) and the human cytomegalovirus (M. Boshart et al., *Cell* 41:521 [1985]).

As used herein, the term "promoter/enhancer" denotes a segment of DNA which contains sequences capable of providing both promoter and enhancer functions (i.e., the functions provided by a promoter element and an enhancer element, see above for a discussion of these functions). For example, the long terminal repeats of retroviruses contain both promoter and enhancer functions. The enhancer/

promoter may be "endogenous" or "exogenous" or "heterologous." An "endogenous" enhancer/promoter is one which is naturally linked with a given gene in the genome. An "exogenous" or "heterologous" enhancer/promoter is one which is placed in juxtaposition to a gene by means of genetic manipulation (i.e., molecular biological techniques) such that transcription of that gene is directed by the linked enhancer/promoter.

The presence of "splicing signals" on an expression vector often results in higher levels of expression of the recombinant transcript. Splicing signals mediate the removal of introns from the primary RNA transcript and consist of a splice donor and acceptor site (J. Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York [1989], pp. 16.7-16.8). A commonly used splice donor and acceptor site is the splice junction from the 16S RNA of SV40.

Efficient expression of recombinant DNA sequences in eukaryotic cells request expression of signals directing the efficient termination and polyadenylation of the resulting transcript. Transcription termination signals are generally found downstream of the polyadenylation signal and are a few hundred nucleotides in length. The term "poly A site" or "poly A sequence" as used herein denotes a DNA sequence which directs both the termination and polyadenylation of the nascent RNA transcript. Efficient polyadenylation of the recombinant transcript is desirable as transcripts lacking a poly A tail are unstable and are rapidly degraded. The poly A signal utilized in an expression vector may be "heterologous" or "endogenous." An endogenous poly A signal is one that is found naturally at the 3' end of the coding region of a given gene in the genome. A heterologous poly A signal is one which is one which is isolated from one gene and placed 3' of another gene. A commonly used heterologous poly A signal is the SV40 poly A signal. The SV40 poly A signal is contained on a 237 bp BamHI/BclI restriction fragment and directs both termination and polyadenylation (J. Sambrook, *supra*, at 16.6-16.7).

Eucaryotic expression vectors may also contain "viral replicons" or "viral origins of replication." Viral replicons are viral DNA sequences which allow for the extrachromosomal replication of a vector in a host cell expressing the appropriate replication factors. Vectors which contain either the SV40 or polyoma virus origin of replication replicate to high copy number (up to 10^4 copies/cell) in cells that express the appropriate viral T antigen. Vectors which contain the replicons from bovine papillomavirus or Epstein-Barr virus replicate extrachromosomally at low copy number (~100 copies/cell).

As used herein, the terms "nucleic acid molecule encoding," "DNA sequence encoding," and "DNA encoding" refer to the order or sequence of deoxyribonucleotides along a strand of deoxyribonucleic acid. The order of these deoxyribonucleotides determines the order of amino acids along the polypeptide (protein) chain. The DNA sequence thus codes for the amino acid sequence.

The term "calcium phosphate co-precipitation" refers to a technique for the introduction of nucleic acids into a cell. The uptake of nucleic acids by cells is enhanced when the nucleic acid is presented as a calcium phosphate-nucleic acid coprecipitate. The original technique of Graham and van der Eb (Graham and van der Eb, *Virol.*, 52:456 [1973]), has been modified by several groups to optimize conditions for particular types of cells. The art is well aware of these numerous modifications.

The term "Southern blot" refers to the analysis of DNA on agarose or acrylamide gels to fractionate the DNA according

to size followed by transfer of the DNA from the gel to a solid support, such as nitrocellulose or a nylon membrane. The immobilized DNA is then probed with a labeled probe to detect DNA species complementary to the probe used. The DNA may be cleaved with restriction enzymes prior to electrophoresis. Following electrophoresis, the DNA may be partially depurinated and denatured prior to or during transfer to the solid support. Southern blots are a standard tool of molecular biologies (J. Sambrook et al. *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, New York, pp. 9.31–9.58 [1989]).

The term “Northern blot” as used herein refers to the analysis of RNA by electrophoresis of RNA on agarose gels to fractionate the RNA according to size followed by transfer of the RNA from the gel to a solid support, such as nitrocellulose or a nylon membrane. The immobilized RNA is then probed with a labeled probe to detect RNA species complementary to the probe used. Northern blots are a standard tool of molecular biologists (J. Sambrook, J. et al., *supra*, pp. 7.39–7.52 [1989]).

The term “YIp plasmid” refers to yeast integrating plasmids which contain selectable yeast genes but lack sequences which allow for the autonomous replication of the plasmid in a yeast cell. Transformation of the host yeast cell occurs by integration of the YIp plasmid into the yeast genome. This integration occurs by recombination between yeast sequences present on the YIp plasmid and homologous sequences present in the genome.

The term “prototrophic” or “prototrophy” refers to an organism that can survive growth under conditions where one or more essential nutrients are lacking in the growth medium. For example, if a yeast cell is transformed to histidine prototrophy, this means that the yeast cell now contains gene sequences encoding the enzyme necessary for the production of the amino acid histidine; therefore, the transformed yeast cell does not require the presence of histidine in the growth medium.

The term “isolated” when used in relation to a nucleic acid, as in “an isolated oligonucleotide” refers to a nucleic acid sequence that is identified and separated from at least one contaminant nucleic acid with which it is ordinarily associated in its natural source. Isolated nucleic acid is such present in a form or setting that is different from that in which it is found in nature. In contrast, non-isolated nucleic acids as nucleic acids such as DNA and RNA found in the state they exist in nature. For example, a given DNA sequence (e.g., a gene) is found on the host cell chromosome in proximity to neighboring genes; RNA sequences, such as a specific mRNA sequence encoding a specific protein, are found in the cell as a mixture with numerous other mRNAs which encode a multitude of proteins. However, isolated nucleic acid encoding a mammalian Chk1 protein includes, by way of example, such nucleic acid in cells ordinarily expressing a Chk1 protein where the nucleic acid is in a chromosomal location different from that of natural cells, or is otherwise flanked by a different nucleic acid sequence than that found in nature. The isolated nucleic acid or oligonucleotide may be present in single-stranded or double-stranded form. When an isolated nucleic acid or oligonucleotide is to be utilized to express a protein, the oligonucleotide will contain at a minimum the sense or coding strand (i.e., the oligonucleotide may single-stranded), but may contain both the sense and anti-sense strands (i.e., the oligonucleotide may be double-stranded).

As used herein, a “portion of a chromosome” refers to a discrete section of the chromosome. Chromosomes are

divided into sites or sections by cytogenetists as follows: the short (relative to the centromere) arm of a chromosome as termed the “p” arm; the long arm is termed the “q” arm. Each arm is then divided into 2 regions termed region 1 and region 2 (region 1 is closest to the centromere). Each region is further divided into bands. The bands may be further divided into sub-bands. For example, the 11p15.5 portion of human chromosome 11 is the portion located on chromosome 11 (11) on the short arm (p) in the first region (1) in the 5th band (5) in sub-band 5 (0.5). A portion of a chromosome may be “altered;” for instance the entire portion may be absent due to a deletion or may be rearranged (e.g., inversions, translocations, expanded or contracted due to changes in repeat regions). In the case of a deletion, an attempt to hybridize (i.e., specifically bind) a probe homologous to a particular portion of a chromosome could result in a negative result (i.e., the probe could not bind to the sample containing genetic material suspected of containing the missing portion of the chromosome). Thus, hybridization of a probe homologous to a particular portion of a chromosome may be used to detect alterations in a portion of a chromosome.

The term “sequences associated with a chromosome” means preparations of chromosomes (e.g., spreads of metaphase chromosomes), nucleic acid extracted from a sample containing chromosomal DNA (e.g., preparations of genomic DNA); the RNA which is produced by transcription of genes located on a chromosome (e.g., hnRNA and mRNA) and cDNA copies of the RNA transcribed from the DNA located on a chromosome. Sequences associated with a chromosome may be detected by numerous techniques including probing of Southern and Northern blots and in situ hybridization to RNA, DNA or metaphase chromosomes with probes containing sequences homologous to the nucleic acids in the above listed preparations.

As used herein the term “coding region” when used in reference to structural gene refers to the nucleotide sequences which encode the amino acids found in the nascent polypeptide as a result of translation of a mRNA molecule. The coding region is bounded, in eukaryotes, on the 5' side by the nucleotide triplet “ATG” which encodes the initiator methionine and on the 3' side by one of the three triplets which specify stop codons (i.e., TAA, TAG, TGA).

As used herein, the term “structural gene” refers to a DNA sequence coding for RNA or a protein. In contrast, “regulatory genes” are structural genes which encode products which control the expression of other genes (e.g., transcription factors).

As used herein, the term “gene” means the deoxyribonucleotide sequences comprising the coding region of a structural gene and the including sequences located adjacent to the coding region on both the 5' and 3' ends for a distance of about 1 kb on either end such that the gene corresponds to the length of the full-length mRNA. The sequences which are located 5' of the coding region and which are present on the mRNA are referred to as 5' non-translated sequences. The sequences which are located 3' or downstream of the coding region and which are present on the mRNA are referred to as 3' non-translated sequences; these sequences. The term “gene” encompasses both cDNA and genomic forms of a gene. A genomic form or clone of a gene contains the coding region interrupted with non-coding sequences termed “introns” or “intervening regions” or “intervening sequences.” Introns are segments of a gene which are transcribed into nuclear RNA (hnRNA); introns may contain regulatory elements such as enhancers. Introns are removed or “spliced out” from the nuclear or primary transcript;

intron therefore are absent in the messenger RNA (mRNA) transcript. The mRNA functions during translation to specify the sequence or order of amino acids in a nascent polypeptide.

In addition in containing introns, genomic forms of a gene may also include sequences located on both the 5' and 3' end of the sequences which are present on the RNA transcript. These sequences are referred to as "flanking" sequences or regions (those flanking sequences are located 5' or 3' to the non-translated sequences present on the mRNA transcript). The 5' flanking region may contain regulatory sequences such as promoters and enhancers which control or influence the transcription of the gene. The 3' flanking region may contain sequences which direct the termination of transcription, post-transcriptional cleavage and polyadenylation.

As used herein, the term "purified" or "to purify" refers to the removal of contaminants from a sample. For example, anti-chk1 antibodies are purified by removal of contaminating non-immunoglobulin proteins; they are also purified by the removal of immunoglobulin that does not bind [chk1] *Chk1*. The removal of non-immunoglobulin proteins and/or the removal of immunoglobulins that do not bind chk1 results in an increase in the percent of [chk1] *Chk1*-reactive immunoglobulins in the sample, in another example, recombinant [chk1] *Chk1* polypeptides are expressed in bacterial host cells and the polypeptides are purified by the removal of host cell proteins; the percent of recombinant [chk1] *Chk1* polypeptides is thereby increased in the sample.

The term "recombinant DNA molecule" as used herein refers to a DNA molecule which is comprised of segments of DNA joined together by means of molecular biological techniques.

The term "recombinant protein" or "recombinant polypeptide" as used herein refers to a protein molecule which is expressed from a recombinant DNA molecule.

The term "native protein" as used herein to indicate that a protein does not contain amino acid residues encoded by vector sequences; that is the native protein contains only those amino acids found in the protein as it occurs in nature. A native protein may be produced by recombinant means or may be isolated from a naturally occurring source.

As used herein, the term "[chk1] *Chk1* protein" or "[chk1] *Chk1* protein sequence" refers to a protein which is encoded by a chk1 gene sequence or to a protein.

As used herein, the term "[non-chk1] *non-Chk1* protein" or "[non-chk1] *non-Chk1* protein sequence" refers to that portion of a fusion protein which comprises a protein or protein sequence which is not derived from a [chk1] *Chk1* protein.

EXPERIMENTAL

The following examples are provided in order to demonstrate and further illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

In the experimental disclosure which follows, the following abbreviations apply: h (human); sc (*Saccharomyces cerevisiae*); sp (*Schizosaccharomyces pombe*); ce (*Caenorhabditis elegans*); dm (*Drosophila melanogaster*); ° C. (degrees Centigrade); rpm (revolutions per minute); BSA (bovine serum albumin); CFA (complete Freund's adjuvant); IFA (incomplete Freund's adjuvant); IgG (immunoglobulin G); IM (intramuscular); IP (intraperitoneal); IV (intravenous or intravascular); SC

(subcutaneous); H₂O (water); HCl (hydrochloric acid); aa (amino acid); bp (base pair); kb (kilobase pair); kD (kilodaltons); gm (grams); µg (micrograms); mg (milligrams); ng (nanograms); µl (microliters); ml (milliliters); mm (millimeters); nm (nanometers); µm (micrometer); M (molar); mM (millimolar); MW (molecular weight); sec (seconds); min(s) (minute/minutes); hr(s) (hour/hours), MgCl₂ (magnesium chloride); NaCl (sodium chloride); OD₂₈₀ (optical density at 280 nm); OD₆₀₀ (optical density at 600 nm); PAGE (polyacrylamide gel electrophoresis); PBS (phosphate buffered saline [150 mM NaCl, 10 mM sodium phosphate buffer, pH 7.2]); PEG (polyethylene glycol); PMSF (phenylmethylsulfonyl fluoride); SDS (Sodium dodecyl sulfate); Tris (tris (hydroxymethyl)aminomethane); w/v (weight to volume); v/v (volume to volume); [Amersham] *AMERSHAM*® (Amersham [Life Science, Inc. Arlington Heights, Ill.] *Biosciences, Piscataway, N.J.*); [ICN] *VALEANT*® ([ICN Pharmaceuticals, Inc., Costa Mesa] *Valeant Pharmaceuticals International, Aliso Viejo, Calif.*); [Amicon] *AMICON*® (Amicon, Inc., Beverly, Mass.); ATCC® ([American Type Culture Collection, Rockville, Md.] *AMERICAN TYPE CULTURE COLLECTION*®, *Manassas, Va.*); Becton Dickinson ([Becton Dickinson Labware, Lincoln Park] *BD, Franklin Lakes, N.J.*); [BioRad] *BIO-RAD*® ([BioRad, Richmond] *Bio-Rad Laboratories, Hercules, Calif.*); [Clontech] *CLONTECH*® (CLONTECH Laboratories, [Palo Alto] *Mountain View, Calif.*); [Difco] *DIFCO*® (Difco Laboratories, Incorporated, Detroit, Mich.); GIBCO® [BRL or Gibco BRL] ([Life Technologies, Inc., Gaithersburg, Md.] *Invitrogen Corporation, Carlsbad, Calif.*); [Invitrogen] *INVITROGEN*® (Invitrogen [Corp.] *Corporation, [San Diego] Carlsbad, Calif.*); [Kodak] *KODAK*® (Eastman Kodak Co., [New Haven, Conn.] *Rochester, N.Y.*); New England Biolabs (New England Biolabs, Inc., [Beverly] *Ipswich, Mass.*); [Novagen] *NOVAGEN*® (Novagen, Inc., Madison, Wis.); [Pharmacia] *PHARMACIA*® (Pharmacia [Inc., Piscataway,] & *Upjohn Company, North Peapack, N.J.*); [Sigma] *SIGMA-ALDRICH*® ([Sigma Chemical] *SIGMA-ALDRICH BIOTECHNOLOGY L.P. Co., St. Louis, Mo.*); [Sorvall] *SORVALL*® ([Sorvall Instruments, a subsidiary of DuPont Co., Biotechnology Systems, Wilmington, Del.] *Ivan Sorvall, Inc. Corporation, Norwalk, Conn.*); [Stratagene] *STRATAGENE*® (Stratagene [Cloning Systems], La Jolla, Calif.); [Whatman] *WHATMAN*® ([Whatman LabSales, Hillsboro, Ore.] *Whatman International Limited, Kent, UK*); Bethyl Laboratories (Bethyl Laboratories Inc., Montgomery, Tex.); and Zeiss (Carl Zeiss, Inc., Thornwood, N.Y.).

Unless otherwise indicated, all restriction enzymes were obtained from New England BioLabs and were used according to the manufacturer's instructions; all oligonucleotide primers, adapter and linkers were synthesized using standard methodologies on [an ABI] a DNA synthesizer. (*Applied Biosystems, Foster City, Calif.*). All chemicals were obtained from [Sigma] *SIGMA-ALDRICH*® unless otherwise indicated.

EXAMPLE 1

Human and Murine Chk1

In this Example, human and murine checkpoint genes were identified by searching for homologs of yeast checkpoint genes. In the experiments to identify the human homolog, degenerate PCR primers to conserved motifs in the kinase domains of spChk1 were used to screen a human B cell library by PCR, as known in the art.

A DNA fragment containing the ORF for the [hChk1] *hCHK1* gene was obtained using the polymerase chain reac-

tion (PCR) as follows. The DNA was obtained from human peripheral blood lymphocytes. Gel-purified degenerate primers used in the reaction were 5'-GGNGGNGAGT/CT/CTNATGGAT/CTT-3' (SEQ ID NO:5) and 5'-TTGGACAGGCCAAAGTC-3' (SEQ ID NO:6). The reaction conditions comprised the following steps: denaturation at 95° C. for 5 minutes, 80° C. for 1 minutes, during which Taq was added, and third cycles of 95° C. for 30 seconds; 52° C. for 30 seconds; and 72° C. for 2 minutes.

Four of 35 clones showed similarity to [spChk1] *spchk1*, and one clone was used to probe 2×10⁵ plaques from a λACT human B cell cDNA library. This probe was also used in subsequent experiments (it is referred to as the "PCR probe"). A partial sequence of this probe was determined and is shown below:

```
(5'gggggggagctgttgaccgaatagagc-
cagacatagcctgctgaaccagatgctcagagattcttccatca actcatgg-
gaggggtggtttatctgcatggtattg-
gaataactcacaggatattaaccagaaaatcttctgttgggaag
aaaggataacctcaaatctcagactttggc-3') (SEQ ID NO:11)
```

The library was constructed using methods known in the art ([See] *see*, T. Durfee et al., *Genes Develop.*, 7:555–569 [1993]), and deposited with the ATCC® (ATCC® Accession No. ATC 87003). The library was screened by hybridizing DNA present on filters with radiolabeled PCR probe (described above) in Hybridization Solution I (48% formamide, 5×SSC, 20 mM Tris-Cl, pH 7.6, 1×Denhardt's solution, 10% dextran sulfate, and 0.1% SDS) at 42° C. overnight. The filters were washed three times in low stringency wash (2×SSC/0.1% SDS) for 10 minutes at room temperature, and twice in high stringency wash (0.2×SSC/0.1% SDS) at 65° C., and then exposed for autoradiography.

The plaques showing hybridization to probe were isolated and used to infect a bacterial strain expressing the Cre enzyme. The recombinant plasmids containing the cDNA inserts were purified and screened by Southern analyses using the PCR-generated probe described and used above. Probing of this library resulted in the identification of two [hChk1] *hCHK1* cDNAs. Neither of these clones were complete, as the longest cDNA was lacking a few base pairs near the 5' end. The full-length clone was constructed by ligating the 5' end of the short clone onto the longer cDNA clone to produce the sequence shown in FIG. 1 ([SEQ ID NO:1] *SEQ ID NO:3*).

Once the human homolog was identical, human [Chk1] *CHK1* cDNA was used to isolate the murine [chk1] *Chk1* ([mChk1] *mChk1*). A mouse T cell cDNA library (ATCC® Accession No. ATC 87291) was screened using a NotI-ClaI fragment from the [hChk1] *hCHK1* cDNA as a probe as described above for the human library screen, with the exception being that the high stringency wash was conducted at 42° C. In addition, a genomic clone was isolated from a mouse ES cell library that contains [chk1] *Chk1*. This clone was found to contain the [chk1] *Chk1* exons (i.e., the sequence provided in [SEQ ID NO:2] *SEQ ID NO:4*).

The sequence of the longest human cDNA (1891 base pairs) predicted a translation product of 476 amino acids, with an approximate size of 54 kD. No in-frame stop codon was found upstream of the first methionine, which is located with the Kozak consensus sequence ([See] *see*, Kozak, *Cell* 44 283 [1986]), and is likely to be the initiation codon, as its encoded proteins is the same size as that observed in cells (as discussed below). FIG. 1 shows the sequence of the cDNA encoding human [chk1] *CHK1* ([SEQ ID NO:1] *SEQ ID NO:3*)(Genbank Accession No. AF016582), as well as the predicted amino acid sequence for the human Chk1 protein ([SEQ ID NO:3] *SEQ ID NO:1*). FIG. 2 shows the cDNA sequence of murine [chk1] *Chk1* ([SEQ ID NO:2] *SEQ ID*

NO:4)(Genbank Accession No. AF016583), as well as the predicted amino acid sequence for the murine Chk1 protein ([SEQ ID NO:4] *SEQ ID NO:2*).

The human [Chk1] *CHK1* gene is likely to be related to *Caenorabditis elegans* gene and the *Drosophila melanogaster* gene *grp*. However, during this experimental work, it was determined that the database DNA sequence for *ceChk1* has a likely frame shift in the COOH-terminus. FIG. 3 shows the domain structure of the predicted human Chk1 (hChk1) protein. In this Figure, the black boxes indicate regions of highest conservation. FIG. 4 shows the alignment of Chk1 homologs. In this Figure, amino acid identities are shown as black boxes, and conservative changes are shown as shaded boxes. In this Figure, "Hs" indicates *Homo sapiens*, "Sp" indicates *S. pombe*, "Ce" indicates *C. elegans*, and "Dm" *D. melanogaster*. In this Figure, the human sequence is SEQ ID NO:7, the *D. melanogaster* sequence is SEQ ID NO:8, the *C. elegans* sequence is SEQ ID NO:9, and the *S. pombe* sequence is SEQ ID NO:10.

The predicted hChk1 protein was found to be 29% identical and 44% similar to spChk1, 40% identical and 56% similar to the ceChk1, and 44% identical and 56% similar to dmChk1. Sequence analysis revealed several COOH-terminal domains that appear to be highly conserved in the Chk1 family of kinases.

EXAMPLE 2

Mapping of Chk1 and Its Expression

In this Example, Northern analysis of hChk1 and murine hChk1 was used to identify tissues that express Chk1. In addition, the chromosomal location of Chk1 was mapped to 11q24, by fluorescence in situ hybridization (FISH), as known in the art.

In the in situ hybridization experiments, tissues from adult mice, and murine embryos from day 15.5 post column (p.c.) were examined. Embryos or tissues were collected and fixed in 4% paraformaldehyde, embedded in paraffin and sectioned on a microtome (Zeiss) at 5μ. Specimens were hybridized with α-³⁵S-UTP labeled [I] riboprobes essentially as described (O. H. Sudin et al., *Develop.*, 108:47; and B. Lutz et al., *Develop.*, 120:25 [1994]). Briefly, pBluescript-[Chk1] *Chk1* was linearized using either BstEI and sense and antisense transcripts were generated using either T7 or T3 polymerase, respectively. Specimens were photographed by double exposure using darkfield illumination with a red filter and Hoechst epifluorescence optics.

In addition to the in situ hybridization with murine tissues, FISH hybridization was used to map, the genomic fragment containing [hchk1] *hCHK1*. For this analysis, metaphase chromosomes prepared from human cells (peripheral blood lymphocytes) were tested with fluorescently-labeled [I] human [chk1] *CHK1* DNA, as known in the art ([See] *see*, e.g., J. W. Ijdo et al., *Genomics* 14:1019–1025 [1992]). Briefly, the longest cDNA obtained from the human B cell library (See, Example 1) was subcloned into [pBlueScript] *pBLUESCRIPT*® (Stratagene, *La Jolla, Calif.*), and used as a probe to screen a human genomic library in the BAC vector by hybridization as described by the manufacturer (Genome Systems). One clone designated "BACH-190 (C16)" (Genome Systems control number 12883) was analyzed by PCR. The PCR conditions were the same as those described in Example 1 above, with the exceptions being that a 42° C. annealing temperature was used, and two primer pairs (primers 186 and 484; and 415 and 177) were used. The sequences of the primers are shown below:

Primer 177: 5'-cta gag gag cag aat cg-3' (SEQ ID NO:12)

Primer 186: 5'-gca gtt tgc agg aca gga taa tct tct cta gga ag-3' (SEQ ID NO:13)

Primer 415: 5'-ttg ctc cag aac ttc tg-3' (SEQ ID NO:14)

Primer 484: 5'-tat tgg ttg act tcc ggc-3' (SEQ ID NO:15)

By automated sequence analysis, it was determined that this clone contained the [chk1] *CHK1* sequence. This clone was then used in FISH analysis as known in the art, in order to determine the chromosomal location of the [chk1] *CHK1* gene.

The results of this analysis placed the gene at a position that is adjacent to the gene encoding ATM on chromosome 11 (i.e., at 11q23). Loss of heterozygosity at this region has been associated with a number of cancers, including breast, lung, and ovarian cancers (I. Vorechovsky et al., *Cancer Res.*, 56:2726 [1996]; and H. Gabra et al., *Cancer Res.*, 56:950 [1996]).

In the Northern analyses, mRNAs from human and mouse tissues were hybridized with 25 ng of labeled human or mouse cDNAs, as appropriate, overnight in 50 mM PIPES, 100 mM NaCl, 50 mM Na₂HPO₄, 1 mM EDTA, and 5% SDS, at 65° C. The blots were washed in PIPES at room temperature, followed by a high stringency wash in 0.1×SSC with 0.5% SDS, at 65° C., for 40 minutes.

The results of the Northern blot analysis (as shown in FIGS. 5A and 5B), revealed the ubiquitous expression of [hChk1] *hCHK1*, with large amounts present in human thymus, testis, small intestine, and colon. In adult mice, [mChk1] *mChk1* was detected in all tissues examined, and large amounts were found in the testis, spleen and lung. In addition, mouse embryos from embryonic day 15.5 also revealed ubiquitous expression, with large amounts detected in the brain, liver, kidney, pancreas, intestines, thymus, and lung. These results were of particular interest, as testis, spleen, and thymus have also been found to express large amounts of ATM (G. Chen and E. Y. H. P. Lee, *J. Biol. Chem.*, 271:33693 [1996]; and N. D. Lakin et al., *Oncogene* 13:2707 [1996]).

EXAMPLE 3

Antibodies Against Chk1

In this Example, affinity-purified antibodies to hChk1 protein ("anti-FL") and the 15 amino acids present on the carboxy terminus of the hChk1 protein ("anti-PEP") were produced. In these experiments, hChk1 protein was first produced in baculovirus as described below.

Recombinant baculovirus encoding glutathione S-transferase (GST) fusion proteins to [hChk1] *hChk1* (GST-[hChk1] *hCHK1*) or a to a mutation of hChk1 in which Asp at position 130 was mutated to Ala (GST-[hChk1] *hCHK1* (D130A)) were produced. Recombinant baculovirus encoding GST-hChk1 and GST-hChk1(D130A) (pYS71) were made by introducing an NdeI at the first ATG of the [hChk1] *hCHK1* open reading frame (ORF) using PCR, and subcloning the [hChk1] *hCHK1* cDNA as an Nde I-EcoRI fragment into pGEX2Tcs ([Invitrogen] *INVITROGEN*) to generate pYS45. The XbaI-EcoRI fragment from pYS45 containing GST-[hChk1] *hCHK1* was then subcloned into pVL1393 ([Invitrogen] *INVITROGEN*), which was cut with XbaI-EcoRI to generate pYS63.

The GST-hChk1(D130A) mutant was generated by the PCR and the XhoI-XmnI fragment containing the mutation was used to replace the wild-type fragment to generate pYS64. The hGST-Chk1(D130A) fragment from pYS64 was then subcloned into the baculovirus transfer vector using the Univector plasmid fusion strategy, as described in co-pending U.S. Patent Application Ser. No. 08/864,224, now issued as U.S. Pat. No. 5,851,808, hereby incorporated by reference.

Viruses were generated by standard methods (e.g., [Baculogold] *BACULOGOLD*, Pharmingen Corporation, San Diego, Calif.). Recombinant GST-hChk1 protein was isolated from infected Hi5 insect cells ([Invitrogen] *INVITROGEN*) on glutathione (GSH) agarose ([Pharmacia] *PHARMACIA*).

The GST-hChk1 protein was then used to produce affinity-purified antibodies. In addition, antibodies directed against the carboxy-terminal 15 amino acids were produced using synthetically produced sequence. Recombinant GST-hChk1 was affinity purified from the cell lysate by chromatography on Glutathione [Sepharose] *SEPHAROSE* 4B™ ([Pharmacia] *GE Healthcare Bio-Sciences AB LLC*, Uppsala, Sweden) according to the manufacturer's instructions.

Polyclonal antibodies against the purified GST-hChk1 or the carboxy terminal amino acids were generated in New Zealand white rabbits (Bethyl Laboratories), using standard techniques ([See] see e.g., E. Harlow and D. Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, New York [1988]). Briefly, rabbits were given an initial immunization comprising 100 µg of affinity purified GST-hChk1 or the 15 amino acid sequence in complete Freund's adjuvant (CFA). The antigen was delivered by SC injection. The animals received boosts comprising 50 µg of affinity purified GST-hChk1 or 15 amino acid sequence, in incomplete Freund's adjuvant (IFA), as appropriate, at the following intervals day 14, day 28, day 42, day 56 and day 70. Sera were collected by bleeding the rabbits from the ear vein and the sera were prepared using standard techniques (E. Harlow and D. Lane, supra, at pp. 117 and 119). The anti-[hChk1] *hChk1* antibodies were referred to as "anti-FL," while the antibodies directed against the carboxy-terminal 15 amino acids were referred to as "anti-PEP."

Anti-PEP antibodies were purified using an affinity column that was prepared by coupling a peptide representing the carboxy-terminal 15 amino acids, at its N terminus to activated CH-[Sepharose] *SEPHAROSE* 4B™ ([Pharmacia] *GE Healthcare*) according to the manufacturer's instructions. The anti-FL antibodies were purified using an affinity column that was prepared by coupling the GST-Chk1 fusion protein from baculovirus to [Affi-Gel] *AFFI-GEL*® 10 ([Biorad] *BIO-RAD*) according to the manufacturer's directions. The antibody concentrations were roughly determined by Bradford analyses. The antibodies were subsequently tested in titration experiments and in Western blots, to determine their titer and specificity.

Affinity purified antibodies to these hChk1 protein made in baculovirus ("anti-FL") or to its COOH-terminal 15 amino acids ("anti-PEP"), recognized a 54-kD protein (FIG. 6) that comigrates with hChk1 expressed in baculovirus. The anti-PEP but not anti-FL signal is competed by addition of excess peptide indicating that the two sera are recognizing different hChk1 epitopes, further confirming identity of the 54-kD band as endogenous hChk1. A 70-kD protein was also specifically recognized by anti-PEP.

Antibodies directed against mChk1 were also produced and purified, using the same methods as described above for the anti-hChk1 antibodies.

mChk1 expressed from the cytomegalovirus promoter, CMV, in baby hamster kidney cells (BHK) resulted in detection of a 54-kD nuclear protein only in transfected cells using antibodies directed against the C-terminal peptide of mChk1. These (and all other transfections) were carried out as follows. Tissue culture flasks (T25) at 70–80% confluence were incubated with 3–9 µg DNA and 15–18 µl lipofectamine (Gibco BRL), in 3 ml of [OptiMEMI] *OPTI-*

MEM® ([Gibco BRL] GIBCO®), for 5–7 hours at 37° C. The cells were washed three times with Dulbecco's PBS without calcium or magnesium, and fed with DMEM with high glucose ([Gibco BRL] GIBCO®) and 10% FBS ([Gibco BRL] GIBCO®). The cells were harvested for Western blots or FACS analyses 48 hours post transfection. The results indicated exogenous mChk1 comigrates with endogenous mChk1 from mouse lung tissue.

EXAMPLE 4

Effect of DNA Damage

To determine whether hChk1 is modified in response to DNA damage like spChk1, hChk1 protein in extracts from cells treated with ionizing radiation was examined.

In the first set of experiments, HeLa cells were synchronized with 2 mM thymidine, and treated without (–) or with (+) 10 Gy of ionizing radiation one hour after release from the block. Cells were collected in G2-M, and extracts were fractionated by 10% SDS-PAGE, and immunoblotted with anti-PEP.

In addition to the HeLa cells, Jurkat cells were treated (+IR) or not treated (–IR) with 10 Gy of ionizing radiation and incubated for two hours. Extracts from these cells were resolved in the first dimension by using isoelectric focusing (IEF), with pH 3 to 10 ampholytes, and in the second dimension on a 10% SDS-PAGE, followed by immunoblotting with anti-PEP.

hChk1 from extracts from damaged cells showed a minor but reproducible reduction in mobility compared to Chk1^{His} from untreated cells (FIG. 7). This modification was confirmed by 2-dimensional gel analysis which clearly demonstrated the generation of a more negatively charged Chk1 species 2 hours after γ -irradiation (FIG. 7). These results indicate that hChk1 may participate in transduction of the DNA damage signal like spChk1.

Indirect immunofluorescence was also conducted. In these experiments, human fibroblasts were fixed, stained with 4'6'-diamidino-2-phenylindole (DAP) to detect DNA, and were probed with affinity-purified anti-PEP, biotinylated antibody to rabbit IgG, and Texas Red streptavidin to reveal the subcellular location of the hChk1 protein. This indirect immunofluorescence revealed that hChk1 is localized to the nucleus in a punctate staining pattern, similar to that observed for ATM.

mChk1 was also tested as described above for hChk1, with the exception that it was expressed in BHK cells. These results also confirmed the nuclear localization of mChk1.

Finally, in order to test for the ability of Chk1^{His} to regulate the cell cycle, [hChk1] *hCHK1* or [hChk1] *hCHK1* (D130A) were transfected under the control of the cytomegalovirus (CMV) promoter, or the CMV vector alone into HeLa cells treated with and without 6 Gy of ionizing radiation. These transfections were accomplished as described in Example 3, above. No perturbation of the cell cycle by either kinase relative to vector alone was detected, suggesting that overproduction alone was insufficient to deregulate the system.

EXAMPLE 5

Phosphorylation of Cdk Tyrosine Phosphorylation Regulators

In this Example, the effects of phosphorylation of key regulators of Cdk tyrosine phosphorylation by [chk1] *Chk1* was investigated.

Tyrosine phosphorylation of Cdc2 has been implicated in cell cycle arrest in response to DNA damage and replication blocks in both *S. pombe* (T Enoch and P. Nurse, Cell 60 665 [1990]), and humans (P. Jin et al., J. Cell Biol., 134.963

[1996]). In *S. pombe*, Cdc2 mutants that cannot be phosphorylated on tyrosine display an inability to arrest the cell cycle in response to blockade of DNA replication. Although it was originally thought that the DNA damage checkpoint did not operate through tyrosine phosphorylation, tyrosine phosphorylation is apparently required for *S. pombe* cells to arrest in response to DNA damage. While it is now clear that tyrosine phosphorylation is required for proper checkpoint control, the experiments implicating tyrosine phosphorylation in this pathway do not distinguish between a regulatory role in which tyrosine phosphorylation rates are manipulated by the checkpoint pathways, or a passive role in which tyrosine phosphorylation is required to allow cell cycle arrest, but is not the actual target of the checkpoint pathway (S. J. Elledge, Science 274: 1664 [1996]; and D. J. Lew and S. Kombluth, Curr. Opin. Cell. Biol., 8:795 [1996]).

Next, the ability of hChk1 to phosphorylate key regulators of Cdk tyrosine phosphorylation, the Cdc25 dual specificity phosphatases, hCdc25A, hCdc25B, and hCdc25C was analyzed. These regulators were chosen for several reasons. First, overproduction of hCdk4 mutants in which the inhibitory tyrosine is changed to phenylalanine abrogates G1 arrest in response to UV light (Y. Terada et al., Nature 376:358 [1995]). Secondly, the UV-sensitivity of [chk1] *Chk1*[–] mutants in *S. pombe* is suppressed by inactivating *cdc25* with a Ts mutation (N. C. Walworth et al., Nature 363:368 [1993]). Finally, in *S. pombe* *wee1mik1* mutants, DNA damage still causes a partial cell cycle delay that could be due to regulation of spCdc25 activity.

GST-hChk1 and GST-hChk1(D130A) were introduced into baculovirus, purified from baculovirus-infected insect cells as described in Example 3 above, and incubated with either GST, His₆-Cdc25C, GST-hCdc25A, hGST-Cdc25B, GST-hCdc25C, or GST-Cdc25C(200–256), and (γ ³²P)ATP.

The kinase reactions contained hGST-Chk1 bound to GSH agarose and either His₆-Cdc25C, GST-Cdc25A, GST-Cdc25B, GST-Cdc25C or GST-Cdc25C(200 to 256) (i.e., amino acids 200 to 256 of Cdc25). Kinase reactions contained 1 to 3 μ g of GST-hChk1 or GST-hChk1(D130A) protein on beads and soluble substrate in 20 mM Hepes (pH 7.4), 10 mM MgCl₂, 10 mM MnCl₂, 2 μ M ATP and 15 μ Ci (γ ³²P)ATP for 30 minutes at 30° C. The proteins were resolved by SDS-PAGE (10%), and visualized by autoradiography for kinase assays (FIG. 8A), or by Coomassie staining (FIG. 8B). Less GST-Cdc25B was loaded than the other substrates (approximately 1/5 of the other substrates was loaded).

GST-Chk1 phosphorylated all three Cdc25 proteins but not GST alone (FIG. 8). Although Gst-Cdc25C^{His} co-migrated with Gst-Chk1^{His} which autophosphorylates, increased phosphorylation was observed at that position relative to that in the presence of kinase alone and phosphorylation of a Gst-Cdc25C^{His} breakdown product was visible.

Protein kinases often form complexes with their substrates. To examine this for hChk1, and the Cdc25 proteins, GST-Cdc25 proteins present on glutathione beads were incubated together with baculovirus extracts expressing His₆-tagged hChk1, and precipitated GST-hCdc25A, GST-hCdc25B, and GST-hCdc25C each specifically bound hChk1 while GST alone did not (FIG. 10). Furthermore, two other GST fusion proteins, GST-Dun1 and GST-Skp1, all failed to bind hChk1. These results indicate that Cdc25 can form complexes with hChk1.

To determine the site on Cdc25C that is phosphorylated by hChk1, the kinase reactions were carried out in a buffer consisting of 50 mM Tris (pH 7.4), 10 mM MgCl₂, 10 μ M ATP, 1 mM DTT and 10 μ Ci (γ ³²P)ATP. The proteins were

separated by SDS-PAGE, transferred to nitrocellulose membranes, and visualized by autoradiography. The nitrocellulose membrane containing His-Cdc25C was excised, blocked with 0.5% polyvinylpyrrolidone (PVP-40) in 100 mM acetic acid for 30 minutes at 37° C., washed six times with water, and digested with TPCK trypsin (Worthington Biochemical Corporation, Lakewood, N.J.) at a final concentration of 30 mg/ml, in 0.1 M NH₄CO₃ (pH 8.0). Further digestion on selected HPLC fractions was performed with 2 units of proline specific endopeptidase ([ICN] VALEANT®) in 0.1M sodium phosphate, 5 mM EDTA (pH 7.4), at 37° C. for 16 hours. Samples were acidified in 1% trifluoroacetic acid (TFA) and loaded onto a [Vydac] VYDAC® C18 column (25 cm×0.46 cm inner diameter, registered to Alltech Associates, Inc., Columbia, Md.). Reverse phase HPLC was performed at 37° C. Reactions were loaded in 0.1% TFA (Buffer A) and eluted with a gradient from 0 to 60% Buffer B (90% acetonitrile, 0.095% TFA). Fractions were collected at 0.5 minutes intervals up to 90 minutes, and counted for radioactivity. Selected fractions were immobilized on Sequenon-AA membrane discs (Millipore Corporation, Billerica, Mass.) for NH₂-terminal sequencing. Manual Edman degradation was done as known in the art ([See] see, J. E. Rodwell et al, J. Biol. Chem., 266:7549 [1991]; and S. Sullivan, and T. W. Wong, Anal. Biochem., 197: 65 [1991]) with a coupling and cleavage temperature of 55° C.

To establish the significance of the Cdc25 phosphorylation, the site of Chk1^{HS} phosphorylation on Cdc25C was mapped. Ser 216 is the main site of phosphorylation of Cdc25C^{HS} in vivo. hChk1 phosphorylated a 56 amino acid region of the hCdc25C protein fused to GST, but not GST alone (FIG. 8). This 56 amino acid motif contains 4 possible sites of phosphorylation. peptide analysis of proteolytic fragments of full length His₆-hCdc25 C phosphorylated with GST-hChk1 revealed a single phosphorylated tryptic peptide by high pressure liquid chromatography. Edman degradation of this peptide indicated release of radioactivity in the third cycle (FIG. 12B). FIG. 12A shows the radioactivity measured from column fractions obtained during reverse phase HPLC. Further degradation of this tryptic fragment with proline endopeptidase resulted in a peptide that released radioactivity in the first cycle. Serine 216 is the only site on Cdc25C^{HS} consistent with this phosphorylation pattern (FIG. 11), as amino acids inclusive of and surrounding Serine²¹⁶ contain amino-terminal trypsin and proline endopeptidase cleavage sites.

In addition, GST-hChk1 purified from baculovirus was incubated with either GST-hCdc25C(200–256) or GST-hCdc25C(200–256)(S216A), and (γ-³²P)ATP, using the same methods as described above. The results are shown in FIG. 9A). In addition, hChk1-His₆ purified from baculovirus was incubated with either GST-hCdc25C (lane 5, FIG. 9B), or GST-hCdc25c(S216A) and (γ-³²P)ATP. Proteins were resolved and visualized as described above. As shown in FIG. 9, there was clear phosphorylation of GST-hCdc25C. A catalytically inactive mutant (GST-hChk1(D130A)(k-) failed to phosphorylate itself or any of the Cdc25 proteins (See, FIG. 9).

To confirm this, the Cdc25C S216A mutation in Gst-Cdc25C and Cdc25C(200–256) were constructed. Both were found to be poor substrates for hChk1 confirming S216 as the site phosphorylation (FIG. 11). S216 has also been reported to be phosphorylated by spChk1, demonstrating phylogenetic conservation of this regulatory relationship.

EXAMPLE 6

Production of Monoclonal Antibodies

The antibodies of the present invention may be monoclonal or polyclonal. Thus, it is within the scope of this

invention to include other (e.g., second antibodies) (monoclonal or polyclonal) directed against or similar to the first antibodies discussed above. It is contemplated that these antibodies will find use in detection assays. Both the first and second antibodies may be used in the detection assays or a first antibody may be used with a commercially available anti-immunoglobulin antibody. An antibody as contemplated herein includes any antibody specific to any region of human or murine Chk1.

The production and use of monoclonal antibodies in an immunoassay is an alternative method to that described in Example 3. Monoclonals provide some advantages because of the ability to produce them in large quantities and the homogeneity of the product. The preparation of hybridoma cell lines for monoclonal antibody production derived by fusing an immortal cell line and lymphocytes sensitized against the immunogenic preparation can be done by techniques which are well known to those who are skilled in the art. (See e.g., Douillard and Hoffman, Basic Facts about Hybridomas, in Compendium of Immunology Vol. II, ed. by Schwartz [1981]; Kohler and Milstein, Nature 256: 495–499 [1975]; Eur. J Immunol., 6: 511–519, [1976]).

Unlike preparation of polyclonal sera, the choice of animal is dependent on the availability of appropriate immortal lines capable of fusing with lymphocytes. Mouse and rat have been the animals of choice in hybridoma technology and are preferably used. Humans can also be utilized as sources for sensitized lymphocytes if appropriate immortalized human (or nonhuman) cell lines are available. For the purpose of the present invention, the animal of choice may be injected with an antigenic amount, for example, from about 0.1 mg to about 20 mg of the enzyme or protein or antigenic parts thereof. Usually the injecting material is emulsified in Freund's complete adjuvant. Boosting injections may also be required. The detection of antibody production can be carried out by testing the antisera with appropriately labeled antigen. Lymphocytes can be obtained by removing the spleen of lymph nodes of sensitized animals in a sterile fashion and carrying out fusion. Alternatively, lymphocytes can be stimulated or immunized in vitro, as described, for example, in Reading, J. Immunol. Meth., 53: 261–291 [1982]D].

A number of cell lines suitable for fusion have been developed and the choice of any particular line for hybridization protocols is directed by any one of a number of criteria such as speed, uniformity of growth characteristics, deficiency of its metabolism for a component of the growth medium, and potential for good fusion frequency.

Intraspecies hybrids, particularly between like strains, work better than interspecies fusions. Several cell lines are available, including mutants selected for the loss of ability to secrete myeloma immunoglobulin.

Cell fusion can be induced either by virus, such as Epstein-Barr or Sendai virus, or polyethylene glycol. Polyethylene (PEG) is the most efficacious agent for the fusion of mammalian somatic cells. PEG itself may be toxic for cells and various concentrations should be tested for effects on viability before attempting fusion. The molecular weight range of PEG may be varied from 1000 to 6000. It gives best results when diluted to from about 20% to about 70% (w/w) in saline or serum-free medium. Exposure to PEG at 37° C. for about 30 seconds is preferred in the present case, utilizing murine cells. Extremes of temperature (i.e., about 45° C.) are avoided, and preincubation of each component of the fusion system at 37° C. prior to fusion can be useful. The ratio between lymphocytes and malignant cells is optimized to avoid cell fusion among spleen cells and a range of from about 1:1 to about 1:10 is commonly used.

The successfully fused cells can be separated from the myeloma line by any technique known by the art. The most common and preferred method is to choose a malignant line which is Hypoxanthine Guanine Phosphoribosyl Transferase (HGPRT) deficient, which will not grow in an aminopterin-containing medium used to allow only growth of hybrids and which is generally composed of hypoxanthine, $[1 \times 10^{-4} M]$ $1 \times 10^{-4} M$, aminopterin $1 \times 10^{-5} M$, and thymidine $3 \times 10^{-5} M$, commonly known as the HAT medium. The fusion mixture can be grown in the HAT-containing culture medium immediately after the fusion 24 hours later. The feeding schedules usually entail maintenance in HAT medium for two weeks and then feeding with either regular culture medium or hypoxanthine, thymidine-containing medium.

The growing colonies are then tested for the presence of antibodies that recognize the antigenic preparation. Detection of hybridoma antibodies can be performed using an assay where the antigen is bound to a solid support and allowed to react to hybridoma supernatants containing putative antibodies. The presence of antibodies may be detected by "sandwich" techniques using a variety of indicators. Most of the common methods are sufficiently sensitive for use in the range of antibody concentrations secreted during hybrid growth.

Cloning of hybrids can be carried out after 21–23 days of cell growth in selected medium. Cloning can be performed by cell limiting dilution in fluid phase or by directly selecting single cells growing in semi-solid agarose. For limiting dilution, cell suspension are diluted serially to yield a statistical probability of having only one cell per well. For the agarose technique, hybrids are seeded in a semi-solid upper layer, over a lower layer containing feeder cells. The colonies from the upper layer may be picked up and eventually transferred to wells.

Antibody-secreting hybrids can be grown in various tissue culture flasks, yielding supernatants with variable concentrations of antibodies. In order to obtain higher concentrations, hybrids may be transferred into animals to obtain inflammatory ascites. Antibody-containing ascites can be harvested 8–12 days after intraperitoneal injection. The ascites contain a higher concentration of antibodies but include both monoclonals and immunoglobulins from the inflammatory ascites. Antibody purification may then be achieved by, for example, affinity chromatography.

Antibodies produced by these methods can then be used in immunoassay methods to detect human or murine Chk1. Such methods include, but are not limited to ELISA (enzyme-linked immunosorbent assay), IFA (immunofluorescence assay), or RIA (radioimmunoassay).

From the above it should be clear that the present invention provides gene sequences encoding mammalian checkpoint genes and proteins useful as probes for a tumors and other malignancies, as well as growth and/or development deficiencies.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 15

<210> SEQ ID NO 1

<211> LENGTH: 476

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Met Ala Val Pro Phe Val Glu Asp Trp Asp Leu Val Gln Thr Leu Gly
 1             5             10             15

Glu Gly Ala Tyr Gly Glu Val Gln Leu Ala Val Asn Arg Val Thr Glu
 20             25             30

Glu Ala Val Ala Val Lys Ile Val Asp Met Lys Arg Ala Val Asp Cys
 35             40             45

Pro Glu Asn Ile Lys Lys Glu Ile Cys Ile Asn Lys Met Leu Asn His
 50             55             60

Glu Asn Val Val Lys Phe Tyr Gly His Arg Arg Glu Gly Asn Ile Gln
 65             70             75             80

Tyr Leu Phe Leu Glu Tyr Cys Ser Gly Gly Glu Leu Phe Asp Arg Ile
 85             90             95

Glu Pro Asp Ile Gly Met Pro Glu Pro Asp Ala Gln Arg Phe Phe His
 100            105            110

Gln Leu Met Ala Gly Val Val Tyr Leu His Gly Ile Gly Ile Thr His
 115            120            125

```


-continued

Arg Asp Ile Lys Pro Glu Asn Leu Leu Leu Asp Glu Arg Asp Asn Leu
 130 135 140
 Lys Ile Ser Asp Phe Gly Leu Ala Thr Val Phe Arg Tyr Asn Asn Arg
 145 150 155 160
 Glu Arg Leu Leu Asn Lys Met Cys Gly Thr Leu Pro Tyr Val Ala Pro
 165 170 175
 Glu Leu Leu Lys Arg Arg Glu Phe His Ala Glu Pro Val Asp Val Trp
 180 185 190
 Ser Cys Gly Ile Val Leu Thr Ala Met Leu Ala Gly Glu Leu Pro Trp
 195 200 205
 Asp Gln Pro Ser Asp Ser Cys Gln Glu Tyr Ser Asp Trp Lys Glu Lys
 210 215 220
 Lys Thr Tyr Leu Asn Pro Trp Lys Lys Ile Asp Ser Ala Pro Leu Ala
 225 230 235 240
 Leu Leu His Lys Ile Leu Val Glu Asn Pro Ser Ala Arg Ile Thr Ile
 245 250 255
 Pro Asp Ile Lys Lys Asp Arg Trp Tyr Asn Lys Pro Leu Lys Lys Gly
 260 265 270
 Ala Lys Arg Pro Arg Val Thr Ser Gly Gly Val Ser Glu Ser Pro Ser
 275 280 285
 Gly Phe Ser Lys His Ile Gln Ser Asn Leu Asp Phe Ser Pro Val Asn
 290 295 300
 Ser Ala Ser Ser Glu Glu Asn Val Lys Tyr Ser Ser Ser Gln Pro Glu
 305 310 315 320
 Pro Arg Thr Gly Leu Ser Leu Trp Asp Thr Ser Pro Ser Tyr Ile Asp
 325 330 335
 Lys Leu Val Gln Gly Ile Ser Phe Ser Gln Pro Thr Cys Pro Asp His
 340 345 350
 Met Leu Leu Asn Ser Gln Leu Leu Gly Thr Pro Gly Ser Ser Gln Asn
 355 360 365
 Pro Trp Gln Arg Leu Val Lys Arg Met Thr Arg Phe Phe Thr Lys Leu
 370 375 380
 Asp Ala Asp Lys Ser Tyr Gln Cys Leu Lys Glu Thr Cys Glu Lys Leu
 385 390 395 400
 Gly Tyr Gln Trp Lys Lys Ser Cys Met Asn Gln Val Thr Ile Ser Thr
 405 410 415
 Thr Asp Arg Arg Asn Asn Lys Leu Ile Phe Lys Val Asn Leu Leu Glu
 420 425 430
 Met Asp Asp Lys Ile Leu Val Asp Phe Arg Leu Ser Lys Gly Asp Gly
 435 440 445
 Leu Glu Phe Lys Arg His Phe Leu Lys Ile Lys Gly Lys Leu Ile Asp
 450 455 460
 Ile Val Ser Ser Gln Lys Val Trp Leu Pro Ala Thr
 465 470 475

<210> SEQ ID NO 2

<211> LENGTH: 476

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 2

Met Ala Val Pro Phe Val Glu Asp Trp Asp Leu Val Gln Thr Leu Gly
 1 5 10 15
 Glu Gly Ala Tyr Gly Glu Val Gln Leu Ala Val Asn Arg Ile Thr Glu
 20 25 30

-continued

Gln Ala Val Ala Val Lys Ile Val Asp Met Lys Arg Ala Ile Asp Cys
 35 40 45

Pro Gln Asn Ile Lys Lys Glu Ile Cys Ile Asn Lys Met Leu Ser His
 50 55 60

Glu Asn Val Val Lys Phe Tyr Gly His Arg Arg Glu Gly His Ile Gln
 65 70 75 80

Tyr Leu Phe Leu Glu Tyr Cys Ser Gly Gly Glu Leu Phe Asp Arg Ile
 85 90 95

Glu Pro Asp Ile Gly Met Pro Glu Gln Asp Ala Gln Arg Phe Phe His
 100 105 110

Gln Leu Met Ala Gly Val Val Tyr Leu His Gly Ile Gly Ile Thr His
 115 120 125

Arg Asp Ile Lys Pro Glu Asn Leu Leu Leu Asp Glu Arg Asp Asn Leu
 130 135 140

Lys Ile Ser Asp Phe Gly Leu Ala Thr Val Phe Arg His Asn Asn Arg
 145 150 155 160

Glu Arg Leu Leu Asn Lys Met Cys Gly Thr Leu Pro Tyr Val Ala Pro
 165 170 175

Glu Leu Leu Lys Arg Lys Glu Phe His Ala Glu Pro Val Asp Val Trp
 180 185 190

Ser Cys Gly Ile Val Leu Thr Ala Met Leu Ala Gly Glu Leu Pro Trp
 195 200 205

Asp Gln Pro Ser Asp Ser Cys Gln Glu Tyr Ser Asp Trp Lys Glu Lys
 210 215 220

Lys Thr Tyr Leu Asn Pro Trp Lys Lys Ile Asp Ser Ala Pro Leu Ala
 225 230 235 240

Leu Leu His Lys Ile Leu Val Glu Thr Pro Ser Ala Arg Ile Thr Ile
 245 250 255

Pro Asp Ile Lys Lys Asp Arg Trp Tyr Asn Lys Pro Leu Asn Arg Gly
 260 265 270

Ala Lys Arg Pro Arg Ala Thr Ser Gly Gly Met Ser Glu Ser Ser Ser
 275 280 285

Gly Phe Ser Lys His Ile His Ser Asn Leu Asp Phe Ser Pro Val Asn
 290 295 300

Asn Gly Ser Ser Glu Glu Thr Val Lys Phe Ser Ser Ser Gln Pro Glu
 305 310 315 320

Pro Arg Thr Gly Leu Ser Leu Trp Asp Thr Gly Pro Ser Asn Val Asp
 325 330 335

Lys Leu Val Gln Gly Ile Ser Phe Ser Gln Pro Thr Cys Pro Glu His
 340 345 350

Met Leu Val Asn Ser Gln Leu Leu Gly Thr Pro Gly Phe Ser Gln Asn
 355 360 365

Pro Trp Gln Arg Leu Val Lys Arg Met Thr Arg Phe Phe Thr Lys Leu
 370 375 380

Asp Ala Asp Lys Ser Tyr Gln Cys Leu Lys Glu Thr Phe Glu Lys Leu
 385 390 395 400

Gly Tyr Gln Trp Lys Lys Ser Cys Met Asn Gln Val Thr Val Ser Thr
 405 410 415

Thr Asp Arg Arg Asn Asn Lys Leu Ile Phe Lys Ile Asn Leu Val Glu
 420 425 430

Met Asp Glu Lys Ile Leu Val Asp Phe Arg Leu Ser Lys Gly Asp Gly
 435 440 445

-continued

Leu Glu Phe Lys Arg His Phe Leu Lys Ile Lys Gly Lys Leu Ser Asp
 450 455 460

Val Val Ser Ser Gln Lys Val Trp Phe Pro Val Thr
 465 470 475

<210> SEQ ID NO 3
 <211> LENGTH: 1821
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

ggccggacag tccgccgagg tgctcgggtg agtcatggca gtgccctttg tggaagactg 60
 ggacttggag caaaccttg gagaaggtgc ctatggagaa gttcaacttg ctgtgaatag 120
 agtaactgaa gaagcagtcg cagtgaagat tgtagatatg aagcgtgccg tagactgtcc 180
 agaaaatatt aagaaagaga tctgtatcaa taaaatgcta aatcatgaaa atgtagtaaa 240
 attctatggt cacaggagag aaggcaatat ccaatattta tttctggagt actgtagtgg 300
 aggagagctt tttgacagaa tagagccaga cataggcatg cctgaaccag atgctcagag 360
 attcttccat caactcatgg caggggtggg ttatctgcat ggtattggaa taactcacag 420
 ggatattaata ccagaaaatc ttctgttggg tgaaagggat aacctcaaaa tctcagactt 480
 tggcttggca acagtatttc ggtataataa tcgtgagcgt ttgttgaaca agatgtgtgg 540
 tactttacca tatgttctc cagaacttct gaagagaaga gaatttcatg cagaaccagt 600
 tgatgtttgg tcctgtggaa tagtacttac tgcaatgctc gctggagaat tgccatggga 660
 ccaaccaggt gacagctgtc aggagtattc tgactggaaa gaaaaaaaa catacctcaa 720
 cccttggaata aaaatcgatt ctgctcctct agctctgctg cataaaatct tagttgagaa 780
 tccatcagca agaattacca ttccagacat caaaaaagat agatggtaca acaaaccct 840
 caagaaaggg gcaaaaaggc cccgagtcac ttcaggtggg gtgtcagagt ctcccagtg 900
 attttctaag cacattcaat ccaatttggg cttctctcca gtaaacagtg cttctagtga 960
 agaaaatgtg aagtactcca gttctcagcc agaaccctgc acaggtcttt ccttatggga 1020
 taccagcccc tcatacattg ataaattggg acaagggatc agcttttccc agcccacatg 1080
 tcctgatcat atgcttttga atagtcagtt acttggcacc ccaggatcct cacagaaccc 1140
 ctggcagcgg ttggtcaaaa gaatgacacg attctttacc aaattggatg cagacaaatc 1200
 ttatcaatgc ctgaaagaga cttgtgagaa gttgggctat caatggaaga aaagttgtat 1260
 gaatcaggtt actatatcaa caactgatag gagaacaat aaactcattt tcaaagtgaa 1320
 tttgttagaa atggatgata aaatattggg tgacttccgg ctttctaagg gtgatggatt 1380
 ggagttcaag agacacttcc tgaagattaa agggaagctg attgatattg tgagcagcca 1440
 gaaggtttgg ctctctgcca catgatcgga ccatcggctc tggggaatcc tgggtaatat 1500
 agtctgctta tgttgacatt attcttctca gagaagatta tcctgtcctg caaactgcaa 1560
 atagtagttc ctgaagtgtt cacttccctg tttatccaaa catcttcaa tttatttgt 1620
 ttgttcggca tacaataat acctatatct taattgtaag caaaactttg gggaaaggat 1680
 gaatagaatt catttgatta tttcttcatg tgtgttagt atctgaattt gaaactcatc 1740
 tggtggaaac caagtttcag gggacatgag ttttccagct tttatacaca cgtatctcat 1800
 ttttatcaaa acattttgtt t 1821

<210> SEQ ID NO 4
 <211> LENGTH: 1962

-continued

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 4

```

gcttgctgct gtgcttggag tcatggcagt gccttttgtg gaagactggg atttggtgca    60
aactttggga gaaggtgcct atggagaagt tcaacttgcct gtgaatagaa taactgaaca    120
agctgttgca gtgaaaattg tagacatgaa gcgggccata gactgtccac aaaatattaa    180
gaaagagatc tgcacataa aaatgttaag ccacgagaat gtagtgaaat tctatggcca    240
caggagggaa ggccatatcc agtatctggt tctggagtac tgtagtggag gagaactttt    300
tgatagaatt gagccagaca tagggatgcc tgaacaagat gctcagaggt tcttccacca    360
actcatggca ggggtggttt atcttcatgg aattggaata actcacaggg atattaacc    420
agaaaacctc ctcttggatg aaagggataa cctcaaaatc tctgactttg gcttggcaac    480
ggtatttcgg cataataatc gtgaacgctt actgaacaag atgtgtggga ctttacctta    540
tgttgctccg gagcttctaa agagaaaaga atttcatgca gaaccagttg atgtttggtc    600
ctgtggaata gtacttactg caatgttggc tggagaattg ccgtgggacc agcccagtga    660
tagctgtcag gaatattctg attggaaaga aaaaaaaaaacc tatctcaatc cttggaaaaa    720
aattgattct gctcctctgg ctttgcttca taaaattcta gttgagactc catcagcaag    780
gatcaccatc ccagacatta agaaagatag atggtacaac aaaccactta acagaggagc    840
aaagaggcca cgcgccacat caggtggtat gtcagagtct tctagtggat tctctaagca    900
cattcattcc aatttggact tttctccagt aaataatggt tccagtgaag aaaccgtgaa    960
gttctctagt tcccagccag agccgagaac agggctttcc ttgtgggaca ctggtccttc   1020
gaacgtggac aaactggttc agggcatcag tttttcccag cctacgtgtc ctgagcatat   1080
gcttgtaaac agtcagttac tcggtacccc tggattttca cagaaccctt ggcagcgctt   1140
ggtcaaaagg atgacacgat tctttactaa attggatgcg gacaaatctt accaatgcct   1200
gaaagagacc ttcgagaagt tgggctatca gtggaagaag agttgtatga atcaggttac   1260
tgtatcaaca actgatagaa gaaacaataa gttgattttc aaaataaatt tggtagaaat   1320
ggatgagaag atactggttg acttccgact ttctaagggt gatggattag agttcaagag   1380
acacttctg aagattaaag ggaagctcag cgatgttgtg agcagccaga aggtttggtt   1440
tcctgttaca tgaggaagct gtcagctctg cacattcctg gtgaatagag tgctgctatg   1500
tgacattttt ctctctagag aagattatct attctgcaa ctgcaaaca tagttgttga   1560
agagttctct tcccattacc caaacatctt ccgatttgta gtgtttggca tacaataact   1620
aatgtatttt aattgtatgt aatgctttgg ggaaaggatg gatcaaattc attaggtatt   1680
tgtccagctg tctttaaatt gtctggattt gaaaccaagt tatgggatac ttgagtttgc   1740
cagcttttat acccatgtag tagtatcact tttgaaaaat caaaagcttg tttcatccca   1800
agcaaaatat tttcttctct gcctatttaa ttgtaaggat gaataaacac agaccatata   1860
cagttgattg gttcatgaat gaggccagcc aaaaaaatgt gtatgttaat gtatgtactg   1920
tattttcagt ttgggtatat gtgctgcaca agggcttgac ca                          1962

```

<210> SEQ ID NO 5

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

<220> FEATURE:

-continued

<221> NAME/KEY: misc_feature
 <222> LOCATION: (3)(6)
 <223> OTHER INFORMATION: n = "t" or "c"
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (6)
 <223> OTHER INFORMATION: n = "t" or "c"
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (13)
 <223> OTHER INFORMATION: n = "t" or "c"

<400> SEQUENCE: 5

ggngngaggt ttnatggatt t

21

<210> SEQ ID NO 6
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

<400> SEQUENCE: 6

ttggacaggc caaagtc

17

<210> SEQ ID NO 7
 <211> LENGTH: 476
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Met Ala Val Pro Phe Val Glu Asp Asn Asp Leu Val Gln Thr Leu Gly
 1 5 10 15

Glu Gly Ala Val Gly Glu Val Gln Leu Ala Val Asn Arg Val Thr Glu
 20 25 30

Glu Ala Val Ala Val Lys Ile Val Asp Met Arg Arg Ala Val Asp Cys
 35 40 45

Pro Glu Asn Ile Lys Lys Glu Ile Cys Ile Asn Lys Met Leu Asn Asn
 50 55 60

Glu Asn Val Val Lys Phe Tyr Gly His Arg Arg Arg Glu Gly Asn Ile
 65 70 75 80

Gln Tyr Leu Phe Leu Glu Tyr Cys Ser Gly Gly Glu Leu Phe Asp Arg
 85 90 95

Ile Glu Pro Asp Ile Gly Met Pro Glu Pro Asp Ala Gln Arg Phe Phe
 100 105 110

His Gln Leu Met Ala Gly Val Val Tyr Leu His Gly Ile Gly Ile Thr
 115 120 125

His Arg Asp Ile Lys Pro Glu Asn Leu Leu Leu Asp Glu His Asp Asn
 130 135 140

Leu Lys Ile Ser Asp Phe Gly Leu Ala Thr Val Phe Arg Tyr Asn Asn
 145 150 155 160

Arg Glu Arg Leu Leu Asn Lys Met Cys Gly Thr Leu Pro Tyr Val Ala
 165 170 175

Pro Glu Leu Leu Lys Arg Arg Glu Phe His Ala Glx Pro Val Asp Val
 180 185 190

Trp Ser Cys Gly Ile Val Leu Thr Ala Met Leu Ala Gly Glu Leu Pro
 195 200 205

Trp Asp Gln Pro Ser Asp Ser Cys Gln Glu Tyr Ser Asp Trp Lys Glu
 210 215 220

Lys Lys Thr Tyr Leu Asn Pro Trp Lys Lys Ile Asp Ser Ala Pro Leu

-continued

225	230	235	240
Ala Leu Leu His Lys Ile Leu Val Glu Asn Pro Ser Ala Arg Ile Thr 245 250 255			
Ile Pro Asp Ile Lys Lys Asp Arg Trp Tyr Asn Lys Pro Leu Lys Lys 260 265 270			
Gly Ala Lys Arg Pro Arg Val Thr Ser Gly Gly Val Ser Glu Ser Pro 275 280 285			
Ser Gly Phe Ser Lys His Ile Gln Ser Asn Leu Asp Phe Ser Pro Val 290 295 300			
Asn Ser Ala Ser Ser Glu Glu Asn Val Lys Tyr Ser Ser Ser Gln Pro 305 310 315 320			
Glu Pro Arg Thr Gly Leu Ser Leu Trp Asp Thr Ser Pro Ser Tyr Ile 325 330 335			
Asp Lys Leu Val Gln Gly Ile Ser Phe Ser Gln Pro Thr Cys Pro Asp 340 345 350			
His Met Leu Leu Asn Ser Gln Leu Leu Gly Thr Pro Gly Ser Ser Gln 355 360 365			
Asn Pro Trp Gln Arg Leu Val Lys Arg Met Thr Arg Phe Phe Thr Lys 370 375 380			
Leu Asp Ala Asp Lys Ser Tyr Gln Cys Leu Lys Glu Thr Glu Lys Leu 385 390 395 400			
Gly Tyr Gln Trp Lys Lys Ser Cys Met Met Gln Val Thr Ile Ser Thr 405 410 415			
Thr Asp Arg Arg Asn Asn Lys Leu Ile Phe Lys Val Asn Leu Leu Glu 420 425 430			
Met Asp Asp Lys Ile Leu Val Asp Phe Arg Leu Ser Lys Gly Asp Gly 435 440 445			
Leu Glu Phe Lys Arg His Phe Leu Lys Ile Lys Gly Lys Leu Ile Asp 450 455 460			
Ile Val Ser Ser Gln Lys Val Trp Leu Pro Ala Thr 465 470 475			

<210> SEQ ID NO 8

<211> LENGTH: 513

<212> TYPE: PRT

<213> ORGANISM: Drosophila melanogaster

<400> SEQUENCE: 8

Met Ala Ala Thr Leu Thr Glu Ala Gly Thr Gly Pro Ala Ala Thr Arg 1 5 10 15
Glu Phe Val Glu Gly Trp Thr Leu Ala Gln Thr Leu Gly Glu Gly Ala 20 25 30
Tyr Gly Glu Val Lys Leu Leu Ile Asn Arg Gln Thr Gly Gly Gly Cys 35 40 45
Gly Met Lys Met Val Asp Leu Lys Lys His Pro Asp Ala Ala Asn Ser 50 55 60
Val Arg Lys Glu Val Cys Ile Gln Lys Met Leu Gln Asp Lys His Ile 65 70 75 80
Leu Arg Phe Phe Gly Lys Arg Ser Gln Gly Ser Val Glu Tyr Ile Phe 85 90 95
Leu Glu Tyr Ala Ala Gly Gly Glu Leu Phe Asp Arg Ile Glu Pro Asp 100 105 110
Val Gly Met Pro Gln His Glu Ala Gln Arg Tyr Phe Thr Gln Leu Leu 115 120 125

-continued

Ser Gly Leu Asn Tyr Leu His Gln Arg Gly Ile Ala His Arg Asp Leu
 130 135 140

Lys Pro Glu Asn Leu Leu Leu Asp Glu His Asp Asn Val Lys Ile Ser
 145 150 155 160

Asp Phe Gly Met Ala Thr Met Phe Arg Cys Lys Gly Lys Glu Arg Leu
 165 170 175

Leu Asp Lys Arg Cys Gly Thr Leu Pro Tyr Val Ala Pro Glu Val Leu
 180 185 190

Gln Lys Ala Tyr Gln Pro Gln Pro Ala Asp Leu Trp Ser Cys Gly Val
 195 200 205

Ile Leu Val Thr Met Leu Ala Gly Glu Leu Pro Trp Asp Gln Pro Ser
 210 215 220

Thr Asn Cys Thr Glu Phe Thr Asn Trp Arg Asp Asn Asp His Trp Gln
 225 230 235 240

Leu Gln Thr Pro Trp Ser Lys Leu Asp Thr Leu Ala Ile Ser Leu Leu
 245 250 255

Arg Lys Leu Leu Leu Ala Thr Ser Pro Gly Thr Arg Leu Thr Leu Glu
 260 265 270

Lys Thr Leu Asp His Lys Trp Cys Asn Met Gln Phe Ala Asp Asn Glu
 275 280 285

Arg Ser Tyr Asp Leu Val Asp Ser Ala Ala Ala Leu Glu Ile Cys Ser
 290 295 300

Pro Lys Ala Lys Arg Gln Arg Leu Gln Ser Ser Ala His Leu Ser Asn
 305 310 315 320

Gly Leu Asp Asp Ser Ile Ser Arg Asn Tyr Cys Ser Gln Pro Met Pro
 325 330 335

Thr Met Arg Thr Asp Asp Asp Phe Asn Val Arg Leu Gly Ser Gly Arg
 340 345 350

Ile Gln Gly Gly Trp Arg Arg Pro Gln Thr Leu Ala Gln Glu Ala Arg
 355 360 365

Leu Ser Tyr Ser Phe Ser Gln Pro Ala Leu Leu Asp Asp Leu Leu Leu
 370 375 380

Ala Thr Gln Met Asn Gln Thr Gln Asn Ala Ser Gln Asn Tyr Phe Gln
 385 390 395 400

Arg Leu Val Arg Arg Met Thr Arg Phe Phe Val Thr Thr Arg Trp Asp
 405 410 415

Asp Thr Ile Lys Arg Leu Val Gly Thr Ile Glu Arg Leu Gly Gly Tyr
 420 425 430

Thr Cys Lys Phe Gly Asp Asp Gly Val Val Thr Val Ser Thr Val Asp
 435 440 445

Arg Asn Lys Leu Arg Leu Val Phe Lys Ala His Ile Ile Glu Met Asp
 450 455 460

Gly Lys Ile Leu Val Asp Cys Arg Leu Ser Lys Gly Cys Gly Leu Glu
 465 470 475 480

Phe Lys Arg Arg Phe Ile Lys Ile Lys Asn Ala Leu Glu Asp Ile Val
 485 490 495

Leu Lys Gly Pro Thr Thr Trp Pro Ile Ala Ile Ala Thr Asn Ser Val
 500 505 510

Pro

<210> SEQ ID NO 9

<211> LENGTH: 483

<212> TYPE: PRT

<213> ORGANISM: Caenorhabditis elegans

-continued

<400> SEQUENCE: 9

Met Ser Ala Ala Ser Thr Thr Ser Thr Pro Ala Ala Ala Ala Val Ala
 1 5 10 15
 Pro Gln Gln Pro Glu Ser Leu Tyr Arg Val Val Gln Thr Leu Gly Glu
 20 25 30
 Gly Ala Phe Gly Glu Val Leu Leu Ile Val Asn Thr Lys Asn Pro Glu
 35 40 45
 Val Ala Ala Ala Met Lys Lys Ile Asn Ile Ala Asn Lys Ser Lys Asp
 50 55 60
 Phe Ile Asp Asn Ile Arg Lys Glu Tyr Leu Leu Gln Lys Arg Val Ser
 65 70 75 80
 Ala Val Gly His Asp Asn Val Ile Arg Met Ile Gly Met Arg Asn Asp
 85 90 95
 Pro Gln Phe Tyr Tyr Leu Phe Leu Glu Tyr Ala Asp Gly Gly Glu Leu
 100 105 110
 Phe Asp Lys Ile Glu Pro Asp Cys Gly Met Ser Pro Val Phe Ala Gln
 115 120 125
 Phe Tyr Phe Lys Gln Leu Ile Cys Gly Leu Lys Phe Ile His Asp Asn
 130 135 140
 Asp Val Val His Arg Asp Ile Lys Pro Glu Asn Leu Leu Leu Thr Gly
 145 150 155 160
 Thr His Val Leu Lys Ile Ser Asp Phe Gly Met Ala Thr Leu Tyr Arg
 165 170 175
 Asn Lys Gly Glu Glu Arg Leu Leu Asp Leu Ser Cys Gly Thr Ile Pro
 180 185 190
 Tyr Ala Ala Pro Glu Leu Cys Ala Gly Lys Lys Tyr Arg Gly Pro Pro
 195 200 205
 Val Asp Val Trp Ser Ser Gly Ile Val Leu Ile Ala Met Leu Thr Gly
 210 215 220
 Glu Leu Pro Trp Asp Arg Ala Ser Asp Ala Ser Gln Ser Tyr Met Gly
 225 230 235 240
 Trp Ile Ser Asn Thr Ser Leu Asp Glu Arg Pro Trp Lys Lys Ile Asp
 245 250 255
 Val Arg Ala Leu Cys Met Leu Arg Lys Ile Val Thr Asp Lys Thr Asp
 260 265 270
 Lys Arg Ala Thr Ile Glu Gln Ile Gln Ala Asp Pro Trp Tyr Gln His
 275 280 285
 Asn Phe Gly Gln Val Glu Thr Pro Asn Gly Arg Pro Leu Lys Arg Ala
 290 295 300
 Arg Asn Asn Asp Glu Asn Ile Thr Cys Thr Gln Gln Ala Glu Cys Ser
 305 310 315 320
 Ala Lys Arg Arg His Leu Glu Thr Pro Asn Glu Lys Ser Thr Leu Ala
 325 330 335
 Glu Arg Gln Asn Ala Ser Phe Ser Gln Pro Thr Lys Thr Glu Asp Leu
 340 345 350
 Leu Leu Thr Gln His Ile Asp Met Ser Gln Thr Asn Ser Asn Leu Leu
 355 360 365
 Gln Arg Met Val Cys Arg Met Thr Arg Phe Cys Val Val Thr Asp Ile
 370 375 380
 Arg Ser Thr Tyr Gln Lys Val Ala Arg Ala Ser Glu His Ala Gly Phe
 385 390 395 400
 Gly Leu Arg Glu Thr Asp Asp Tyr Arg Leu Leu Val Thr Trp Arg Glu

-continued

	405		410		415
Val	Ser Met Met	Val Ser Leu Tyr	Thr Met Gly Asp	Ile Pro Asp Lys	
	420		425	430	
Pro	Arg Val Met	Val Asp Phe Arg	Ser Leu Ala Val	Thr Glu Ser Ser	
	435		440	445	
Leu	Arg Arg Cys	Ser Trp Thr	Leu Glu Thr Val	Cys Met Ser Gly Tyr	
	450		455	460	
Val	Pro Thr Glu	Thr Thr Gly Ser	Pro Ile Leu Asp	Met Cys Gln Glu	
465		470	475	480	
Ile	Arg Arg				

<210> SEQ ID NO 10

<211> LENGTH: 496

<212> TYPE: PRT

<213> ORGANISM: Schizosaccharomyces pombe

<400> SEQUENCE: 10

Met	Ala	Gln	Lys	Leu	Asp	Asn	Phe	Pro	Tyr	His	Ile	Gly	Arg	Glu	Ile
1				5					10					15	
Gly	Thr	Gly	Ala	Phe	Ala	Ser	Val	Arg	Leu	Cys	Tyr	Asp	Asp	Asn	Ala
			20					25					30		
Lys	Ile	Tyr	Ala	Val	Lys	Phe	Val	Asn	Lys	Lys	His	Ala	Thr	Ser	Cys
		35					40					45			
Met	Asn	Ala	Gly	Val	Trp	Ala	Arg	Arg	Met	Ala	Ser	Glu	Ile	Gln	Leu
	50					55					60				
His	Lys	Leu	Cys	Asn	Gly	His	Lys	Asn	Ile	Ile	His	Phe	Tyr	Asn	Thr
65					70					75					80
Ala	Glu	Asn	Pro	Gln	Trp	Arg	Trp	Val	Val	Leu	Glu	Phe	Ala	Gln	Gly
				85					90					95	
Gly	Asp	Leu	Phe	Asp	Lys	Ile	Glu	Pro	Asp	Val	Gly	Ile	Asp	Glu	Asp
		100						105					110		
Val	Ala	Gln	Phe	Tyr	Phe	Ala	Gln	Leu	Met	Glu	Gly	Ile	Ser	Phe	Met
		115					120					125			
His	Ser	Lys	Gly	Val	Ala	His	Arg	Asp	Leu	Lys	Pro	Glu	Asn	Ile	Leu
	130					135					140				
Leu	Asp	Tyr	Asn	Gly	Asn	Leu	Lys	Ile	Ser	Asp	Phe	Gly	Phe	Ala	Ser
145					150					155					160
Leu	Phe	Ser	Tyr	Lys	Gly	Lys	Ser	Arg	Leu	Leu	Asn	Ser	Pro	Val	Gly
				165					170					175	
Ser	Pro	Pro	Tyr	Ala	Ala	Pro	Glu	Ile	Thr	Gln	Gln	Tyr	Asp	Gly	Ser
			180					185					190		
Lys	Val	Asp	Val	Trp	Ser	Cys	Gly	Ile	Ile	Leu	Phe	Ala	Leu	Leu	Leu
	195					200						205			
Gly	Asn	Thr	Pro	Trp	Asp	Glu	Ala	Ile	Ser	Asn	Thr	Gly	Asp	Tyr	Leu
	210					215						220			
Leu	Tyr	Lys	Lys	Gln	Cys	Glu	Arg	Pro	Ser	Tyr	His	Pro	Trp	Asn	Leu
225					230					235					240
Leu	Ser	Pro	Gly	Ala	Tyr	Ser	Ile	Ile	Thr	Gly	Met	Leu	Arg	Ser	Asp
			245						250					255	
Pro	Phe	Lys	Arg	Tyr	Ser	Val	Lys	His	Val	Val	Gln	His	Pro	Trp	Leu
			260					265						270	
Thr	Ser	Ser	Thr	Pro	Phe	Arg	Thr	Lys	Asn	Gly	Asn	Cys	Ala	Asp	Pro
	275						280					285			
Val	Ala	Leu	Ala	Ser	Arg	Leu	Met	Leu	Lys	Leu	Arg	Ile	Asp	Leu	Asp

-continued

290	295	300			
Lys Pro Arg Leu Ala Ser Ser Arg Ala Ser Gln Asn Asp Ser Gly Phe					
305	310	315			320
Ser Met Thr Gln Pro Ala Phe Lys Lys Asn Asp Gln Lys Glu Leu Asp					
	325	330			335
Arg Val Glu Val Tyr Gly Ala Leu Ser Gln Pro Val Gln Leu Asn Lys					
	340	345			350
Asn Ile Asp Val Thr Glu Ile Leu Glu Lys Asp Pro Ser Leu Ser Gln					
	355	360			365
Phe Cys Glu Asn Glu Gly Phe Ile Lys Arg Leu Ala Lys Lys Ala Lys					
	370	375			380
Asn Phe Tyr Glu Ile Cys Pro Pro Glu Arg Leu Thr Arg Phe Tyr Ser					
385	390	395			400
Arg Ala Ser Arg Glu Thr Ile Ile Asp His Leu Tyr Asp Ser Leu Arg					
	405	410			415
Leu Leu Ala Ile Ser Val Thr Met Lys Tyr Val Arg Asn Gln Thr Ile					
	420	425			430
Leu Tyr Val Asn Leu His Asp Lys Arg Lys Cys Leu Leu Gln Gly Val					
	435	440			445
Ile Glu Leu Thr Asn Leu Gly His Asn Leu Glu Leu Ile Asn Phe Ile					
	450	455			460
Lys Arg Asn Gly Asp Pro Leu Glu Trp Arg Lys Phe Phe Lys Asn Val					
465	470	475			480
Val Ser Ser Ile Gly Lys Pro Ile Val Leu Thr Asp Val Ser Gln Asn					
	485	490			495

<210> SEQ ID NO 11
 <211> LENGTH: 186
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

<400> SEQUENCE: 11

```

gggggggagc tgtttgaccg aatagagcca gacataggca tgcctgaacc agatgctcag      60
agattcttcc atcaactcat gggaggggtg gtttatctgc atggtattgg aataactcac      120
aggatatta aaccagaaaa tcttctgttg gaagaaaggg ataacctcaa aatctcagac      180
tttggc                                           186
  
```

<210> SEQ ID NO 12
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

<400> SEQUENCE: 12

```

ctagaggagc agaatcg                                           17
  
```

<210> SEQ ID NO 13
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

<400> SEQUENCE: 13

```

gcagtttgca ggacaggata atcttctcta ggaag                                           35
  
```


-continued

<210> SEQ ID NO 14
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

<400> SEQUENCE: 14

ttgctccaga acttctg

17

<210> SEQ ID NO 15
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

<400> SEQUENCE: 15

tattggttga cttceggc

18

What is claimed is:

1. A purified *human* protein, wherein the protein is a polypeptide having checkpoint kinase activity and encoded by [the nucleotide sequence of] SEQ ID NO: 3.

2. A purified *human* protein [comprising the amino acid sequence set forth in] having checkpoint kinase activity, wherein the protein comprises SEQ ID NO: 1.

3. A fusion protein comprising [a portion of] at least 15 sequential amino acids of [the] a carboxy-terminus [of the CHK1 protein of claim 2,] sequence of SEQ ID NO:1 and a non-[CHK1] *Chk1* protein [sequence].

25 4. The fusion protein of claim 3, wherein said non-*Chk1* protein sequence [comprises] is combined with an affinity tag for purification of the fusion protein.

5. The fusion protein of claim 4, wherein said affinity tag [comprises] is a histidine tag.

30 6. A purified [Chk1] protein [encoded by the nucleotide sequence] of claim 5.

7. The fusion protein of either claim 3 or claim 6, wherein the non-*Chk1* protein is glutathione *S*-transferase.

* * * * *