

US00RE41491E

(19) **United States**
 (12) **Reissued Patent**
 DeLuca et al.

(10) **Patent Number:** **US RE41,491 E**
 (45) **Date of Reissued Patent:** ***Aug. 10, 2010**

(54) **(20S)-1 α -HYDROXY-2-METHYLENE-19-NOR-BISHOMOPREGNACALCIFEROL AND ITS USES**

(75) Inventors: **Hector F. DeLuca**, Deerfield, WI (US);
Lori A. Plum, Arena, WI (US);
Margaret Clagett-Dame, Deerfield, WI (US)

(73) Assignee: **Wisconsin Alumni Research Foundation**, Madison, WI (US)

(*) Notice: This patent is subject to a terminal disclaimer.

(21) Appl. No.: **12/116,066**

(22) Filed: **May 6, 2008**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **6,627,622**
 Issued: **Sep. 30, 2003**
 Appl. No.: **10/078,204**
 Filed: **Feb. 18, 2002**

U.S. Applications:

(60) Provisional application No. 60/341,138, filed on Dec. 13, 2001.

(51) **Int. Cl.**
A61K 31/59 (2006.01)

(52) **U.S. Cl.** **514/167; 552/653**

(58) **Field of Classification Search** **514/167; 552/653**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,666,634 A 5/1987 Miyamoto

4,800,198 A 1/1989 DeLuca et al.
 5,089,641 A 2/1992 DeLuca et al.
 5,536,713 A 7/1996 Deluca et al.
 5,578,587 A 11/1996 DeLuca et al.
 5,587,497 A 12/1996 DeLuca et al.
 5,843,928 A 12/1998 Deluca et al.
 5,936,133 A 8/1999 Deluca et al.
 5,945,410 A 8/1999 DeLuca et al.
 6,579,861 B2 6/2003 DeLuca et al.
 6,835,723 B2 12/2004 DeLuca et al.

FOREIGN PATENT DOCUMENTS

WO WO 8606255 11/1986
 WO WO 96/01811 1/1996
 WO WO 9841501 9/1998
 WO WO 0174766 10/2001
 WO WO 0220021 3/2002

OTHER PUBLICATIONS

Brown et al, "New Active Analogues of Vitamin D with Low Calcemic Activity", *Kidney International*, vol. 38, Suppl. 29, 1990, pp. S-22-S-27.

(Continued)

Primary Examiner—Sang-ming Hui


(74) *Attorney, Agent, or Firm*—Andrus Scales, Starke & Sawall, LLP

(57) **ABSTRACT**


This invention discloses (20S)-1 α -hydroxy-2-methylene-19-nor-bishomopregnacalciferol and pharmaceutical uses therefor. This compound exhibits pronounced activity in arresting the proliferation of undifferentiated cells and inducing their differentiation to the monocyte thus evidencing use as an anti-cancer agent and for the treatment of skin diseases such as psoriasis as well as skin conditions such as wrinkles, slack skin, dry skin and insufficient sebum secretion. This compound also has little, if any, calcemic activity and therefore may be used to treat immune disorders in humans as well as renal osteodystrophy.

1 Claim, 6 Drawing Sheets

1 α , 25-DIHYDROXY-19-NORVITAMIN D3



1 α , 25-DIHYDROXYVITAMIN D3
 A.K.A. NATIVE HORMONE



OTHER PUBLICATIONS

DeLuca, Hector F. et al, Data Accession No. 108:180270 XP002238138, "Analogues of the Hormonal Form of Vitamin D and Their Possible Use in Leukemia", Progress in Clinical and Biological Research (1988), 259 (Nutr., Growth, Cancer), 41-55.

Hareau et al, "Asymmetric Synthesis of 1 α ,25-Dihydroxyvitamin D₃-A-Ring Precursor Starting with 5-Tert-Butyldimethylsiloxy-2-Cyclohexanone," *Tetrahedron Letters*, 41, 2000, pp. 2385-2388.

Hekutaa, Efu Deruuka et al, Data Accession No. 120:153700 XP002238139, "Cell Differentiation-Inducing Agents Containing Seosterols", JP 05 238937 A, WARF, Japan (Sep. 17, 1993).

Ostrem, Voula K. et al, Data Accession No. 107:191147 XP002238137, "Induction of Monocytic Differentiation of HL-60 Cells by 1,25-Dihydroxyvitamin D Analogs", *Journal of Biological Chem.* (1987), 262(29), 14164-71.


Posner et al, "2-Fluoroalkyl A-Ring Analogs of 1,25-Dihydroxyvitamin D₃ Stereocontrolled Total Synthesis Via Intramolecular and Intermolecular Diels-Alder Cycloadditions, Preliminary Biological Testing", *Journal of Organic Chemistry*, 1995, vol. 60, No. 14, pp. 4617-4628.

Sicinski et al, "New 1 α , 25-Hydroxyvitamin D₃ Compounds of High Biological Activity: Synthesis and Biological Evaluation of 2-Hydroxymethyl, 25-Methyl, and 2-Methylene Analogues," *Journal of Medical Chemistry*, 1998, vol. 41, No. 23, pp. 4662-4674.


Sicinski et al, "New 1 α ,25-Dihydroxy-19-Norvitamin D₃ Compounds of High Biological Activity: Synthesis and Biological Evaluation of 2-Hydroxymethyl, 2-Methyl, and 2-Methylene Analogues" *J. Med. Chem.*, 41 (1998) pp. 4662-4674.

Posner et al. "2-Fluoroalkyl A-Ring Analogs of 1,25-Dihydroxyvitamin D₃. Stereocontrolled Total Synthesis via Intramolecular and Intermolecular Diels-Alder Cycloadditions. Preliminary Biological Testing", *J. Org. Chem.* 60, 4617 (1995), pp. 4617-4626.


1 α , 25-DIHYDROXY-19-NORVITAMIN D3




1 α ,25-DIHYDROXYVITAMIN D3
A.K.A. NATIVE HORMONE



1 α -HYDROXY-2-METHYLENE-
19-NOR-PREGNACALCIFEROL
A.K.A 2-Mpregna



(20S)-1 α -HYDROXY-2-METHYLENE-
19-NOR-BISHOMOPREGNACALCIFEROL
A.K.A 2-MbisP



1 α -HYDROXY-2-METHYLENE-19-
NOR-HOMOPREGNACALCIFEROL
A.K.A 2-MP




FIG. 2





FIG. 3






FIG. 4A




FIG. 4B




FIG. 4C




FIG. 5A




FIG. 5B





FIG. 5C

FIG. 6



3

reaction, and rejection of organ transplants; and additionally for the treatment of inflammatory diseases, such as rheumatoid arthritis, asthma, and inflammatory bowel diseases such as celiac disease and Crohn's disease, as well as the improvement of bone fracture healing and improved bone grafts. Acne, alopecia and hypertension are other conditions which may be treated with the compound of the invention.

The above compound is also characterized by relatively high cell differentiation activity. Thus, this compound also provides a therapeutic agent for the treatment of psoriasis, or as an anti-cancer agent, especially against leukemia, colon cancer, breast cancer and prostate cancer. In addition, due to its relatively high cell differentiation activity, this compound provides a therapeutic agent for the treatment of various skin conditions including wrinkles, lack of adequate dermal hydration, i.e. dry skin, lack of adequate skin firmness, i.e. slack skin, and insufficient sebum secretion. Use of this compound thus not only results in moisturizing of skin but also improves the barrier function of skin.

The compound may be present in a composition to treat the above-noted diseases and disorders in an amount from about 0.01 $\mu\text{g/gm}$ to about 100 $\mu\text{g/gm}$ of the composition, and may be administered topically, transdermally, orally or parenterally in dosages of from about 0.01 $\mu\text{g/day}$ to about 100 $\mu\text{g/day}$.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A–1E illustrate the structures of the compounds described and tested herein, namely, 1 α ,25-dihydroxy-19-nor-vitamin D₃, hereinafter 1,25(OH)₂19-nor-D₃ (FIG. 1A); the native hormone 1 α ,25-dihydroxyvitamin D₃, hereinafter 1,25(OH)₂D₃ (FIG. 1B); 1 α -hydroxy-2-methylene-19-nor-pregnacalciferol, hereinafter 2-Mpregna (FIG. 1C); (20S)-1 α -hydroxy-2-methylene-19-nor-bishomopregnacalciferol, hereinafter 2-MbisP (FIG. 1D); and 1 α -hydroxy-2-methylene-19-nor-homopregnacalciferol, hereinafter 2-MP (FIG. 1E);

FIG. 2 is a graph illustrating the relative activity of 1,25(OH)₂19-nor-D₃, 2-Mpregna, 2-MbisP, 2-MP and 1,25(OH)₂D₃ to compete for binding with [³H]-1,25-(OH)₂-D₃ to the full-length recombinant rat vitamin D receptor;

FIG. 3 is a graph illustrating the percent HL-60 cell differentiation as a function of the concentration of 1,25(OH)₂19-nor-D₃, 2-Mpregna, 2-MbisP, 2-MP and 1,25(OH)₂D₃;

FIGS. 4A–4C are bar graphs illustrating the bone calcium mobilization activity of 1,25(OH)₂D₃ and 1,25(OH)₂19-nor-D₃ as compared to 2-MbisP (FIG. 4A), 2-Mpregna (FIG. 4B), and 2-MP (FIG. 4C);

FIGS. 5A–5C are bar graphs illustrating the intestinal calcium transport activity of 1,25(OH)₂D₃ and 1,25(OH)₂19-nor-D₃ as compared to 2-MbisP (FIG. 5A), 2-Mpregna (FIG. 5B), and 2-MP (FIG. 5C); and

FIG. 6 is a bar graph illustrating blood serum calcium levels in female rats after treatment with chronic doses of 1,25(OH)₂D₃ and 1,25(OH)₂19-nor-D₃ as compared to 2-MbisP (FIG. 6A), 2-Mpregna (FIG. 6B), and 2-MP (FIG. 6C).




DETAILED DESCRIPTION OF THE INVENTION

(20S)-1 α -hydroxy-2-methylene-19-nor-bishomopregnacalciferol (referred to herein as 2-MbisP) was synthesized and tested. Structurally, this 19-nor analog is characterized by the general formula I previously illustrated herein.

The preparation of (20S)-1 α -hydroxy-2-methylene-19-nor-bishomopregnacalciferol having the basic structure I can

4

be accomplished by a common general method, i.e. the condensation of a bicyclic Windaus-Grundmann type ketone II with the allylic phosphine oxide III to the corresponding 2-methylene-19-nor-vitamin D analog IV followed by deprotection at C-1 and C-3 in the latter compound:



In the structures II, III, and IV groups Y₁ and Y₂ are hydroxy-protecting groups, it being also understood that any functionalities that might be sensitive, or that interfere with the condensation reaction, be suitably protected as is well-known in the art. The process shown above represents an application of the convergent synthesis concept, which has been applied effectively for the preparation of vitamin D compounds [e.g. Lythgoe et al., J. Chem. Soc. Perkin Trans. I, 590 (1978); Lythgoe, Chem. Soc. Rev. 9, 449 (1983); Toh et al., J. Org. Chem. 48, 1414 (1983); Baggolini et al., J. Org. Chem. 51, 3098 (1986); Sardina et al., J. Org. Chem. 51, 1264 (1986); J. Org. Chem. 51, 1269 (1986); DeLuca et al., U.S. Pat. No. 5,086,191; DeLuca et al., U.S. Pat. No. 5,536,713].

Hydrindanones of the general structure II are known, or can be prepared by known methods.

For the preparation of the required phosphine oxides of general structure III, a synthetic route has been developed starting from a methyl quinate derivative which is easily obtained from commercial (1R,3R,4S,5R)-(-)-quinic acid as described by Perlman et al., Tetrahedron Lett. 32, 7663 (1991) and DeLuca et al., U.S. Pat. No. 5,086,191.

The overall process of the synthesis of compound I is illustrated and described more completely in U.S. Pat. No. 5,843,928 entitled "2-Alkylidene-19-Nor-Vitamin D Compounds" the specification of which is specifically incorporated herein by reference.

5

Biological Activity of (20S)-1 α -Hydroxy-2-Methylene-19-Nor-Bishomopregnacalciferol

The introduction of a methylene group to the 2-position of 1 α -hydroxy-19-nor-pregnacalciferol had little or no effect on binding to the porcine intestinal vitamin D receptor, as compared to 1 α ,25-dihydroxyvitamin D₃. This compound bound equally well to the porcine receptor as compared to the standard 1,25-(OH)₂D₃ (FIG. 2). It might be expected from these results that this compound would have equivalent biological activity. Surprisingly, however, the 2-methylene substitution in 2-MbisP produced a highly selective analog with unique biological activity.

FIG. 4B shows that 2-MbisP has very little activity as compared to that of 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃), the natural hormone, in stimulating intestinal calcium transport.

FIG. 5B demonstrates that 2-MbisP has very little bone calcium mobilization activity, as compared to 1,25(OH)₂D₃.

FIGS. 4B and 5B thus illustrate that 2-MbisP may be characterized as having little, if any, calcemic activity.

FIG. 3 illustrates that 2-MbisP is almost as potent as 1,25(OH)₂D₃ on HL-60 differentiation, making it an excellent candidate for the treatment of psoriasis and cancer, especially against leukemia, colon cancer, breast cancer and prostate cancer. In addition, due to its relatively high cell differentiation activity, this compound provides a therapeutic agent for the treatment of various skin conditions including wrinkles, lack of adequate dermal hydration, i.e. dry skin, lack of adequate skin firmness, i.e. slack skin, and insufficient sebum secretion. Use of this compound thus not only results in moisturizing of skin but also improves the barrier function of skin.

FIG. 6 shows an analysis of serum calcium in rats after administration of chronic doses of 2-MbisP. These data provide further support for the data in FIGS. 4B and 5B that 2-MbisP has very little calcemic activity and thus a relatively low risk of producing hypercalcemia at recommended doses.

The battery of in vitro and in vivo assays described in Sicinski et al (J. Med. Chem. 41, 4662-4674, 1998) were used to compare the biological activities of the compounds, 2-Mpregna, 2-MbisP and 2-MP, with 1,25(OH)₂ 19-nor-D₃, and 1,25(OH)₂D₃, the native vitamin D hormone.

The differentiation of HL-60 promyelocytic into monocytes was determined as described by Ostrem et al (J. Biol. Chem. 262, 14164-14171, 1987).

Interpretation of Data

VDR binding and HL60 cell differentiation. 2-MbisP, 2-Mpregna and 2-MP are nearly equivalent (K_i=0.3, 0.6 and 0.3 nM for 2-Mpregna, 2-MbisP and 2-MP, respectively) in their ability to compete with [³H]-1,25(OH)₂D₃ for binding to the full-length recombinant rat vitamin D receptor (FIG. 2). Furthermore, the competition binding activity of these three compounds is similar to that of 1,25(OH)₂ 19-nor-D₃ (K_i=0.2 nM), as well as the native hormone, 1,25(OH)₂D₃ (K_i=0.1 nM). There is little difference between any of these five compounds in their ability (efficacy or potency) to promote HL60 differentiation with the possible exception of 2-Mpregna (EC₅₀=17 nM), which is slightly less potent than 2-MbisP (EC₅₀=7 nM), 2-MP (EC₅₀=6 nM), 1,25(OH)₂ 19-nor-D₃ (EC₅₀=4 nM) and 1,25(OH)₂D₃ (EC₅₀=5 nM) (See FIG. 3). This result suggests that 2-MbisP will be very effective in psoriasis because it has direct cellular activity in causing cell differentiation and in suppressing cell growth. It also indicates that it will have significant activity as an anti-

6

cancer agent, especially against leukemia, colon cancer, breast cancer and prostate cancer, as well as against skin conditions such as dry skin (lack of dermal hydration), undue skin slackness (insufficient skin firmness), insufficient sebum secretion and wrinkles.

Calcium mobilization from bone and intestinal calcium absorption in vitamin D-deficient animals. Using vitamin D-deficient rats on a low calcium diet (0.02%), the activities of these compounds in intestine and bone were tested. As expected, the native hormone (1,25(OH)₂D₃) increased serum calcium levels at all dosages (FIG. 4). FIG. 4 also shows that 2-MbisP, 2-Mpregna and 2-MP have little, if any, activity in mobilizing calcium from bone. Administration of 2-MbisP, 2-Mpregna, 2-MP or 1,25(OH)₂ 19-nor-D₃ at 260 pmol/day for 7 days did not result in mobilization of bone calcium, and increasing the amount of 2-MbisP, 2-Mpregna and 2-MP to 1300 pmol/day (5-fold) was without effect. Similar findings were obtained with 1,25(OH)₂ 19-nor-D₃.

Intestinal calcium transport was evaluated in the same groups of animals using the everted gut sac method (FIG. 5). These results show that 2-MbisP and 2-Mpregna do not promote intestinal calcium transport when administered at either 260 or 1300 pmol/day, whereas 1,25(OH)₂D₃ promotes a significant increase at the 260 pmol/day dose. In contrast to 2-MbisP and 2-Mpregna, 2-MP showed calcium transport activity equivalent to 1,25(OH)₂D₃ at this and the 5-fold higher concentration but higher doses actually decreased this activity. As shown in FIG. 4, 1,25(OH)₂ 19-nor-D₃, like the 2-MbisP and 2-Mpregna derivatives, is devoid of intestinal calcium transport activity.

Serum calcium response in vitamin D-sufficient animals on a normal calcium diet. The desirability of 2-MbisP, 2-Mpregna and 2-MP is exemplified by their inability to produce hypercalcemia in normal animals compared to 1,25(OH)₂ 19-nor-D₃, and 1,25(OH)₂D₃. In the experiment shown in FIG. 6, normal, female rats were given 2500 pmol/day for 7 days of 1,25(OH)₂D₃, or 5000 pmol/day of 2-MbisP, 2-Mpregna, 2-MP or 1,25(OH)₂ 19-nor-D₃. The doses were administered to two separate groups of animals by either the oral or intraperitoneal route, and serum calcium levels were assessed 4 hours after the last dose. Animals receiving 1,25(OH)₂D₃ by either route exhibited hypercalcemia, some severe enough to require euthanasia. Likewise, 1,25(OH)₂ 19-nor-D₃ produced frank hypercalcemia. In contrast, no increase in serum calcium was seen in any of the animals receiving the compounds, 2-MP, 2-MbisP or 2-Mpregna.

These results illustrate that 2-MbisP is an excellent candidate for numerous human therapies and that it may be useful in a number of circumstances such as autoimmune diseases, cancer, and psoriasis. 2-MbisP is an excellent candidate for treating psoriasis because: (1) it has significant VDR binding and cellular differentiation activity; (2) it is devoid of hypercalcemic liability unlike 1,25(OH)₂ 19-nor-D₃ and 1,25(OH)₂D₃; and (3) it is easily synthesized. Since 2-MbisP has significant binding activity to the vitamin D receptor, but has little ability to raise blood serum calcium, it may also be useful for the treatment of renal osteodystrophy.

For treatment purposes, the compound of this invention defined by formula I may be formulated for pharmaceutical applications as a solution in innocuous solvents, or as an emulsion, suspension or dispersion in suitable solvents or carriers, or as pills, tablets or capsules, together with solid carriers, according to conventional methods known in the art. Any such formulations may also contain other pharmaceutically-acceptable and non-toxic excipients such

7

as stabilizers, anti-oxidants, binders, coloring agents or emulsifying or taste-modifying agents.

The compound may be administered orally, topically, parenterally or transdermally. The compound is advantageously administered by injection or by intravenous infusion or suitable sterile solutions, or in the form of liquid or solid doses via the alimentary canal, or in the form of creams, ointments, patches, or similar vehicles suitable for transdermal applications. Doses of from 0.01 μg to 100 μg per day of the compounds are appropriate for treatment purposes, such doses being adjusted according to the disease to be treated, its severity and the response of the subject as is well understood in the art. Since the compound exhibits specificity of action, each may be suitably administered alone, or together with graded doses of another active vitamin D compound—e.g. 1α -hydroxyvitamin D_2 or D_3 , or $1\alpha,25$ -dihydroxyvitamin D_3 —in situations where different degrees of bone mineral mobilization and calcium transport stimulation is found to be advantageous.

Compositions for use in the above-mentioned treatments comprise an effective amount of the (20S)- 1α -hydroxy-2-methylene-19-nor-bishomopregnacalciferol as defined by the above formula I as the active ingredient, and a suitable carrier. An effective amount of such compound for use in accordance with this invention is from about 0.01 μg to about 100 μg per gm of composition, and may be administered topically, transdermally, orally or parenterally in dosages of from about 0.01 $\mu\text{g}/\text{day}$ to about 100 $\mu\text{g}/\text{day}$.

The compound may be formulated as creams, lotions, ointments, topical patches, pills, capsules or tablets, or in liquid form as solutions, emulsions, dispersions, or suspensions in pharmaceutically innocuous and acceptable solvent or oils, and such preparations may contain in addition other pharmaceutically innocuous or beneficial components, such as stabilizers, antioxidants, emulsifiers, coloring agents, binders or taste-modifying agents.

The compound is advantageously administered in amounts sufficient to effect the differentiation of promyelocytes to normal macrophages. Dosages as described above are suitable, it being understood that the amounts given are to be adjusted in accordance with the severity of the disease, and the condition and response of the subject as is well understood in the art.

The formulations of the present invention comprise an active ingredient in association with a pharmaceutically acceptable carrier therefore and optionally other therapeutic ingredients. The carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulations and not deleterious to the recipient thereof.

Formulations of the present invention suitable for oral administration may be in the form of discrete units as capsules, sachets, tablets or lozenges, each containing a pre-

8

determined amount of the active ingredient; in the form of a powder or granules; in the form of a solution or a suspension in an aqueous liquid or non-aqueous liquid; or in the form of an oil-in-water emulsion or a water-in-oil emulsion.

Formulations for rectal administration may be in the form of a suppository incorporating the active ingredient and carrier such as cocoa butter, or in the form of an enema.

Formulations suitable for parenteral administration conveniently comprise a sterile oily or aqueous preparation of the active ingredient which is preferably isotonic with the blood of the recipient.


Formulations suitable for topical administration include liquid or semi-liquid preparations such as liniments, lotions, applicants, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions or suspensions such as drops; or as sprays.

For asthma treatment, inhalation of powder, self-propelling or spray formulations, dispensed with a spray can, a nebulizer or an atomizer can be used. The formulations, when dispensed, preferably have a particle size in the range of 10 to 100 μ .

The formulations may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy. By the term "dosage unit" is meant a unitary, i.e. a single dose which is capable of being administered to a patient as a physically and chemically stable unit dose comprising either the active ingredient as such or a mixture of it with solid or liquid pharmaceutical diluents or carriers.

We claim:

1. (20S)- 1α -hydroxy-2-methylene-19-nor-bishomopregnacalciferol having the formula:



* * * * *