



US00RE41800E

(19) **United States**
 (12) **Reissued Patent**
Ueda et al.

(10) **Patent Number:** **US RE41,800 E**
 (45) **Date of Reissued Patent:** **Oct. 5, 2010**

(54) **METHOD FOR PRODUCING L-GLUTAMIC ACID**
 (75) Inventors: **Hiroshi Ueda**, Kawasaki (JP); **Takayuki Koda**, Kawasaki (JP); **Masakazu Sato**, Kawasaki (JP)
 (73) Assignee: **Ajinomoto Co., Inc.**, Tokyo (JP)
 (21) Appl. No.: **12/485,550**
 (22) Filed: **Jun. 16, 2009**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **7,319,025**
 Issued: **Jan. 15, 2008**
 Appl. No.: **10/338,915**
 Filed: **Jan. 9, 2003**

U.S. Applications:

(63) Continuation of application No. 10/077,999, filed on Feb. 20, 2002, now abandoned.

(51) **Int. Cl.**
G12P 13/14 (2006.01)

(52) **U.S. Cl.** **435/110**

(58) **Field of Classification Search** None
 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,032,474	A	5/1962	Foster
3,220,929	A	11/1965	Kinoshita et al.
3,331,828	A	7/1967	Inamine et al.
5,908,768	A	6/1999	Ono et al.
6,197,559	B1	3/2001	Moriya et al.
6,331,419	B1	12/2001	Moriya et al.
6,596,517	B2	7/2003	Izui et al.
6,653,110	B2	11/2003	Sato et al.
6,682,912	B2	1/2004	Ito et al.
6,995,250	B1	2/2006	Hirano et al.
7,015,010	B1 *	3/2006	Izui et al.
7,037,690	B2	5/2006	Hara et al.
7,090,998	B2	8/2006	Ishikawa et al.
7,125,977	B2	10/2006	Hirano et al.
7,183,403	B2	2/2007	Hirano et al.
7,198,943	B2	4/2007	Koda et al.
7,205,132	B2	4/2007	Hirano et al.
7,208,296	B2	4/2007	Izui et al.
7,247,459	B1	7/2007	Izui et al.
7,262,035	B2	8/2007	Nakamura et al.
7,294,491	B2	11/2007	Ueda et al.
7,319,025	B2	1/2008	Ueda et al.
7,344,874	B2	3/2008	Hara et al.
7,354,744	B2	4/2008	Takahashi et al.
7,432,085	B2	10/2008	Hara et al.
7,501,282	B2	3/2009	Hara et al.
2001/0019836	A1	9/2001	Moriya et al.
2002/0137150	A1	9/2002	Ohtaki et al.
2002/0182688	A1	12/2002	Izui et al.
2002/0192772	A1	12/2002	Sato et al.
2003/0119153	A1	6/2003	Moriya et al.
2003/0148474	A1	8/2003	Gusyatiner et al.
2003/0172698	A1	9/2003	Koda et al.
2004/0180404	A1	9/2004	Ishikawa et al.
2005/0196846	A1	9/2005	Hara et al.

2005/0227334	A1	10/2005	Izui et al.
2006/0019355	A1	1/2006	Ueda et al.
2006/0040365	A1	2/2006	Kozlov et al.
2006/0084151	A1 *	4/2006	Ueda et al.
2006/0110813	A1	5/2006	Takahashi et al.
2006/0115878	A1	6/2006	Hara et al.
2006/0141588	A1	6/2006	Nakamura et al.
2006/0160191	A1	7/2006	Kataoka et al.
2006/0246552	A1	11/2006	Hara et al.
2007/0038419	A1	2/2007	Usuda et al.
2007/0092953	A1	4/2007	Nakamura et al.
2007/0134773	A1	6/2007	Izui et al.
2007/0172932	A1	7/2007	Hirano et al.
2009/0162907	A1	6/2009	Izui et al.
2009/0215131	A1	8/2009	Hara et al.
2009/0226981	A1	9/2009	Hara et al.
2009/0246835	A1	10/2009	Iwatani et al.
2009/0258402	A1	10/2009	Hirano et al.
2009/0263874	A1	10/2009	Moriya et al.

FOREIGN PATENT DOCUMENTS

AU	761990	4/2000
CN	1069109 C	3/1999
EP	0 636 695	2/1995
EP	0 670 370	9/1995
EP	0 952 221	10/1999
EP	1 078 989	2/2001
EP	1078989	2/2001
GB	921 075	3/1963
JP	44-30249	12/1969
JP	62-288	1/1987
WO	WO 97/08294	3/1997

OTHER PUBLICATIONS

Francoise Gavini, et al., "Transfer of *Enterobacter agglomerans* (Beijerinck 1888) Ewing and Fife 1972 to *Pantoea gen. nov.* as *Pantoea agglomerans comb. nov.* and Description of *Pantoea dispersa sp. nov.*," International Journal of Systematic Bacteriology, vol. 39, No. 3, Jul. 1989, pp. 337-345.

Mergaert, J., et al., "Transfer of *Erwinia ananas* (synonym, *Erwinia uredovora*) and *Erwinia stewartii* to the Genus *Pantoea emend.* as *Pantoea ananas* (Serrano 1928) comb. nov. and *Pantoea stewartii* (Smith 1898) comb. nov., Respective-ly, and Description of *Pantoea stewartii* subsp. *indologenes* subsp. *nov.*," International Journal of Systematic Bacteriology, vol. 43, No. 1, Jan. 1993, 162-173.

(Continued)

Primary Examiner—Herbert J. Lilling

(74) Attorney, Agent, or Firm—Oblon, Spivak, McClelland, Maier & Neustadt, L.L.P.

(57) **ABSTRACT**

A method for producing L-glutamic acid by fermentation, which comprises culturing a microorganism having L-glutamic acid-producing ability in a liquid medium of which pH is adjusted to a condition under which L-glutamic acid produced by the microorganism is allowed to be precipitated, to allow L-glutamic acid to be produced and accumulated with precipitation of L-glutamic acid accompanied, wherein an operation causing existence of L-glutamic acid crystals in the medium is performed when a concentration of L-glutamic acid in the medium is lower than the concentration at which spontaneous crystallization occurs.

17 Claims, 7 Drawing Sheets

OTHER PUBLICATIONS

Robacker, D. C., et al., "Purine Metabolizing Capability of *Enterobacter agglomerans* Affects Volatiles Production and Attractiveness to Mexican Fruit Fly," *Journal of Chemical Ecology*, vol. 28, No. 8, Aug. 2002, pp. 1549–1563.

Moutaouakkil, A., et al., "Purification and Partial Characterization of Azoreductase from *Enterobacter agglomerans*," *Archives of Biochemistry and Biophysics* 413, 2003, pp. 139–146.

Moutaouakkil, A., et al., "Decolorization of Azo Dyes with *Enterobacter agglomerans* Immobilized in Different Supports by Using Fluidized Bed Bioreactor," *Current Microbiology*, vol. 48, 2004, pp. 124–129.

Bergey's Manual of Systematic Bacteriology, 2nd Edition, vol. 2, published in 2005, Springer Science+Business Media, Inc., New York, NY, pp. 712–721.

Food and Fermentation Industries, 1995, No. 2, pp. 22–26 (w/English Abstract).

Abstract of Crit. Rev. Biotechnol., vol. 15, No. 1, 1 page, "Recent Advances in the Physiology and Genetics of Amino Acid-Producing Bacteria", 1995.

Separation TEchnology Discussion Group, pp. 159–153, "Separation Technology Series 4, Industrial Crystalization", 1992 (with summarized English translation).

Kwei-Chao Chao et al., *Departemne3t of Bacteriology, The University of TEXas, Austin*, pp. 715–725, "A Glutamic Acid Producing *Bacillus*", Nov. 10, 1958.

Database WPI, Section Ch, Week 197630, Derwent Publications Ltd., London, GB; Class B05, AN 1976–57114X, XP002214438 & JP 51 020488 B (Takeda Chem Ind Ltd), Jun. 25, 1976.

R.M. Borischewski, *Journal of Bacteriology*, vol. 93, No. 2, pp. 597–599, "Keto Acids as Growth-Limiting Factors in Autotrophic Growth of *Thiobacillus* Thiooxidants", Feb. 1967.

Database WPI, Section C, Week 198706, Derwent Publications Ltd., Lend, GB; AN 1987–040881, XP002214439 & JP 62 000288A (Sanraku Ocewan Co Ltd), Jan. 6, 1987.

Kramer R. "Genetic and Physiological Approaches for the Production of Amino Acids", *Journal of Biotechnology*<Elsevier Science Publishers, Amsterdam, NL. vol. 45, No. 1, Feb. 12, 1996, pp. 1–21 XP004036833.

* cited by examiner




FIG. 1

[88.0% / 935 aa]

1' MQNSAMKPMLDSSWLAGANQSYIEQLYEDFLTOPDSVDAVIRSMFQQLPGTGVKPEQFHS
 1' MQNSALKAMLDSSYLSGANQSYIEQLYEDFLTOPDSVDAMTRSTFQQLPGTGVKPDQFHS

61' ATREYFRRLAKDASRYTSSYTDPATNSKQVKYLQLINAFRFRGHQEANLDPLGLWKQDRV
 61' QTREYFRRLAKDASRYSSSTISDPOTHVKQVKYLQLINAYRFRGHQHANLDPLGLWQDKV

121' ADLDPAFHDLTDADFQESFNVGSFAIGKEMKLDLFDALKQTYCGSIGAEMHINNTEE
 121' ADLDPFHDLTEADFQETFNVGSFASGKEMKLGELLEALKQTYCGPIGAEMHITSTEE

181' KRNIQQRIESGASQTSFSGEEKKGFLELTAEGLEKYLGAKFPGAKRFSLEGGDALVPM
 181' KRNIQQRIESG--RATFNSEKKRFLSELTAEGLEKYLGAKFPGAKRFSLEGGDALIPM

241' LREMIRHAGKSGTREVYLGMAHRGRLNVLINVLGKKPQDLFDEFSGKHKEHLGTGDVKYH
 239' LKEMIRHAGNSGTREVYLGMAHRGRLNVLINVLGKKPQDLDEFAGKHKEHLGTGDVKYH

301' MGFSSDIETEGGLVHLALAFNPSHLEIVSPVWGSYRARRLDRLAEPVSNKVLPTIHGDA
 299' MGFSSDFQTDGGLVHLALAFNPSHLEIVSPVWIGSVRARRLDRLEPSSNKVLPTIHGDA

361' AVICQGVVQETLNMSQARGYEVGGTVRIVINNQGFTTSHPKDARSTPYCTDIGKMLAP
 359' AVTGQGVVQETLNMSKARGYEVGGTVRIVINNQGFTTSHPLDARSTPYCTDIGKMQAP

421' IFHVNADPEAVAFVTRLALDYRNTFKRDVFIDLVCYRRHGHNEADEPSATQPLMYQIK
 419' IFHVNADPEAVAFVTRLALDFRNTFKRDVFIDLVSYRRHGHNEADEPSATQPLMYQIK

481' KHPTPRKIYADRLEGEVASQEDATEMVNLYRDALDAGECVPEWRPMSLHSFTWSPYLN
 479' KHPTPRKIYADKLEQKVATLEDATEMVNLYRDALDAGDCVVAEWRPMMHHSFTWSPYLN

541' HEWDEPYPAQYDMKRLKELALRISQVPEQIEVQSRVAKIYNDRKLMAGEKAFDWGGAEN
 539' HEWDEEYPNKVEMKRLQELAKRISTVPEAVEMQSRVAKIYGDRQAMAAGEKLDWGGAEN

601' LAYATLVOEGIPVRLSGEDSGRGTFFHRHAVVHNQANGSTYTPLHHIHNSQGEFKVWOSV
 599' LAYATLVOEGIPVRLSGEDSGRGTFFHRHAVIHNQNSHGSTYTPLQHHNGQGAFRVWOSV

661' LSEEAVLAFEYGYATAEPRVLTINEAQFGDFANGAQVVIDQFISSEGEQKNGRMCGLVMLL
 659' LSEEAVLAFEYGYATAEPRTLTINEAQFGDFANGAQVVIDQFISSEGEQKNGRMCGLVMLL

721' PHGYEGQGPEHSSARLERYLQLCAEQNMQVCVPSTPAQVYHMLRRQALRGMRPLVVMSP
 719' PHGYEGQGPEHSSARLERYLQLCAEQNMQVCVPSTPAQVYHMLRRQALRGMRPLVVMSP

781' KSLLRHPLAISSLELANGSFQPAIGEIDLOPQGVCRVVLCSGKVVYDLLEQRRKDEXT
 779' KSLLRHPLAVSSLELANGTFLPAIGEIDLOPKGVKRVVMCSGKVVYDLLEQRRKNNQH

841' DVAIVRIEQLYPPFHQAVQEALKAYSHVQDFVWCQEELNQGAWYCSQHHFRDVVYVPGAT
 839' DVAIVRIEQLYPPFKAMQEVLLQFAHVKDFVWCQEELNQGAWYCSQHHFREVYVPGAS

901' LRYAGRPASASPAVGYMSVHQQQQDLVNDALNVN
 899' LRYAGRPASASPAVGYMSVHQQQQDLVNDALNVE

FIG. 2

[88.2% / 407 aa]

1' MSSVDILVPDLPEVADATVATYHKKPGDAVSFIDEVIVEIETDKVYLEVPASADGVLEAY

 1" MSSVDILVPDLPEVADATVATYHKKPGDAVVRIDEVLVEIETDKVYLEVPASADGILDAV

61' LEDEGATVTSRQILGRLKEGNSAGKESAKAE;NODTPAQRQTASLEEESSDALSPAIRR

 61" LEDEGTTVTSRQILGRLREGNSAGKETSAXSE;EKASTPAQRQQASLEEQNNDALSPAIRR

121' LIAEHNLDAAQIKGTGVGGRLTREDVEKHLANKPQAEKAAAPAGAATAQQPVANRSEKR

 121" LLAEHNLDASAIAIKGTGVGGRLTREDVEKHLAKAPAKE--SAPAAAAPAAQPALAARSEKR

181' VPMTRLRKRVAERLLEAKNSTAMLTTFNEINMKPIMDLRKQYGDAFEKRHGVRLLGFMSFY

 179" VPMTRLRKRVAERLLEAKNSTAMLTTFNEVNMKPIMDLRKQYGEAFEKRHGIRLLGFMSFY

241' IKAVVEALKRYPEVNASIDGEDVYHNYFDVSIYVSTPRGLVTPVLRDVALSMADIEKK

 239" VKAVVEALKRYPEVNASIDGDVVYHNYFDVSMYVSTPRGLVTPVLRDVTLLGMADIEKK

301' IKELAVKGRDGKLTVDLGGNFTITNGGVFGSLMSTPIINPPQSAILGMHAIKDRPMAV

 299" IKELAVKGRDGKLTVEDLGGNFTITNGGVFGSLMSTPIINPPQSAILGMHAIKDRPMAV

361' NGQVVILPMMYLALSYDHRLIDGRESVGYLVAVKEMLEDPARLLLDV

 359" NGQVEILPMMYLALSYDHRLIDGRESVGFVLIKELLEDPTRLLLDV

FIG. 3

[95.1% / 41 aa]

1' MNLHEYQAKQLFARYGMPAPTGYACTTPREAE EAASKIGAG

 1" MNLHEYQAKQLFARYGLPAPVGYACTTPREAE EAASKIGAGPMWVKCQVHAGGRGKAGGY

FIG. 4

[97.4% / 39 aa]

1' AFSVFRCHSIMNCVSVCPKGLNPTRAI GHIKSM L LQ RSA

 181" FLIDSRDTETDSRLOGLSDAFSVFRCHSIMNCVSVCPKGLNPTRAI GHIKSM L LQRNA

FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9

METHOD FOR PRODUCING L-GLUTAMIC ACID

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.

This application is a continuation of U.S. application Ser. No. 10/077,999, filed on Feb. 20, 2002, now abandoned.

BACKGROUND OF THE INVENTION

The present invention relates to a method for producing L-glutamic acid by fermentation. L-Glutamic acid is widely used as a raw material of seasonings and so forth.

L-Glutamic acid is produced mainly by fermentation utilizing so-called L-glutamic acid-producing coryneform bacteria belonging to the genus *Brevibacterium*, *Corynebacterium* or *Microbacterium* or mutant strains thereof (Amino Acid Fermentation, Gakkai Shuppan Center, pp. 195-215, 1986). As methods for producing L-glutamic acid by fermentation by using other bacterial strains, there are known a method using a microorganism belonging to the genus *Bacillus*, *Streptomyces*, *Penicillium* or the like (U.S. Pat. No. 3,220,929), a method using a microorganism belonging to the genus *Pseudomonas*, *Arthrobacter*, *Serratia*, *Candida* or the like (U.S. Pat. No. 3,563,857), a method using a microorganism belonging to the genus *Bacillus*, *Pseudomonas*, *Serratia*, *Aerobacter aerogenes* (currently referred to as *Enterobacter aerogenes*) or the like (Japanese Patent Publication (Kokoku) No. 32-9393), a method using a mutant strain of *Escherichia coli* (Japanese Patent Application Laid-open (Kokai) No. 5-244970) and so forth. In addition, the inventors of the present invention proposed a method for producing L-glutamic acid by using a microorganism belonging to the genus *Klebsiella*, *Erwinia* or *Pantoea* (Japanese Patent Application Laid-open No. 2000-106869).

Further, there have been disclosed various techniques for improving L-glutamic acid-producing ability by enhancing activities of L-glutamic acid biosynthetic enzymes through use of recombinant DNA techniques. For example, it was reported that introduction of a gene coding for citrate synthase derived from *Escherichia coli* or *Corynebacterium glutamicum* was effective for enhancement of L-glutamic acid-producing ability in *Corynebacterium* or *Brevibacterium* bacteria (Japanese Patent Publication (Kokoku) No. 7-121228). In addition, Japanese Patent Application Laid-open No. 61-268185 discloses a cell harboring recombinant DNA containing a glutamate dehydrogenase gene derived from *Corynebacterium* bacteria. Further, Japanese Patent Application Laid-open No. 63-214189 discloses a technique for increasing L-glutamic acid-producing ability by amplifying a glutamate dehydrogenase gene, an isocitrate dehydrogenase gene, an aconitate hydratase gene and a citrate synthase gene.

Although L-glutamic acid productivity has been considerably increased by the aforementioned breeding of microorganisms or improvement of production methods, development of methods for more efficiently producing L-glutamic acid at a lower cost is required to meet to further increase of the demand in future.

There is known a method wherein fermentation is performed as L-amino acid accumulated in culture is crystallized (Japanese Patent Application Laid-open No. 62-288). In this method, the L-amino acid concentration in the culture

is maintained below a certain level by precipitating the accumulated L-amino acid in the culture. Specifically, L-tryptophan, L-tyrosine or L-leucine is precipitated during fermentation by adjusting temperature and pH of the culture or adding a surfactant to a medium.

While a method of carrying out fermentation with precipitation of L-amino acid accompanied is known as described above, amino acids suitable for this method are those showing a relatively low water solubility, and no example of applying the method to highly water-soluble amino acids such as L-glutamic acid is known. In addition, the medium must have low pH to precipitate L-glutamic acid. However, L-glutamic acid-producing bacteria such as those mentioned above cannot grow under an acidic condition, and therefore L-glutamic acid fermentation is performed under neutral conditions (U.S. Pat. Nos. 3,220,929 and 3,032,474; K. C. Chao & J. W. Foster, *J. Bacteriol.*, 77, pp. 715-725 (1959)). Thus, production of L-glutamic acid by fermentation accompanied by precipitation is not known. Furthermore, it is known that growth of most acidophile bacteria is inhibited by organic acids such as acetic acid, lactic acid and succinic acid (Yasuro Oshima Ed., "Extreme Environment Microorganism Handbook", p. 231, Science Forum; R. M. Borichewski, *J. Bacteriol.*, 93, pp. 597-599 (1967) etc.). Therefore, it is considered that many microorganisms are susceptible to L-glutamic acid, which is also an organic acid, under acidic conditions, and there has been no report that search of microorganisms showing L-glutamic acid-producing ability under acidic conditions was attempted.

SUMMARY OF THE INVENTION

Under the circumstances as described above, an object of the present invention is to provide a method for producing L-glutamic acid by fermentation, with precipitation of L-glutamic acid accompanied.

The inventors of the present invention found that, in a method for producing L-glutamic acid by fermentation, which comprising culturing a microorganism having L-glutamic acid-producing ability, to allow L-glutamic acid to be produced and accumulated in a medium with precipitation of L-glutamic acid accompanied, a yield of L-glutamic acid when existence of crystals is artificially caused before spontaneous crystallization occurs, is better than that when spontaneous crystallization of L-glutamic acid produced by the microorganism is allowed to occur. Thus, they accomplished the present invention.

The present invention provides the followings.

- (1) A method for producing L-glutamic acid by fermentation, which comprises culturing a microorganism having L-glutamic acid-producing ability in a liquid medium of which pH is adjusted to a condition under which L-glutamic acid produced by the microorganism is allowed to be precipitated, to allow L-glutamic acid to be produced and accumulated with precipitation of L-glutamic acid accompanied, wherein an operation causing existence of L-glutamic acid crystals in the medium is performed when a concentration of L-glutamic acid in the medium is lower than the concentration at which spontaneous crystallization occurs.
- (2) The method according (1), wherein the operation causing existence of L-glutamic acid crystals in the medium is addition of L-glutamic acid crystals.
- (3) The method according to (2), wherein the crystals are α -crystals.
- (4) The method according to (3), wherein an amount of the crystals added to the medium is 0.2 g/L or more.

(5) The method according to any one of (1) to (4), wherein the microorganism belongs to the genus **[Enterobacter]** *Pantoea*.

(6) The method according to (5), wherein the microorganism is **[Enterobacter agglomerans]** *Pantoea ananatis*.

(7) The method according to any one of (1) to (6), wherein the microorganism can metabolize a carbon source in a liquid medium containing L-glutamic acid at a saturation concentration and the carbon source, at a specific pH, and has an ability to accumulate L-glutamic acid in an amount exceeding the saturation concentration of L-glutamic acid in the liquid medium at the pH.

(8) The method according to (7), wherein the specific pH is 5.0 or less.

According to the methods of the present invention, L-glutamic acid can be efficiently produced. Also, in a preferred embodiment, α -crystals which are preferable as a product, can be produced.

BRIEF EXPLANATION OF THE DRAWINGS

FIG. 1 is a restriction enzyme map of a DNA fragment derived from **[Enterobacter agglomerans]** *Pantoea ananatis* in pTWVEK101.

FIG. 2 shows comparison of an amino acid sequence deduced from a nucleotide sequence of an *sucA* gene derived from **[Enterobacter agglomerans]** *Pantoea ananatis* and that derived from *Escherichia coli* (upper: **[Enterobacter agglomerans]** *Pantoea ananatis*, column: *Escherichia coli*, the same shall apply to the followings).

FIG. 3 shows comparison of an amino acid sequence deduced from a nucleotide sequence of an *sucB* gene derived from **[Enterobacter agglomerans]** *Pantoea ananatis* and that derived from *Escherichia coli*.

FIG. 4 shows comparison of an amino acid sequence deduced from a nucleotide sequence of an *sucC* gene derived from **[Enterobacter agglomerans]** *Pantoea ananatis* and that derived from *Escherichia coli*.

FIG. 5 shows comparison of an amino acid sequence deduced from a nucleotide sequence of an *sdhB* gene derived from **[Enterobacter agglomerans]** *Pantoea ananatis* and that derived from *Escherichia coli*.

FIG. 6 shows construction of a plasmid pMWCPG containing a *gltA* gene, a *ppc* gene and a *gdhA* gene.

FIG. 7 shows construction of a plasmid RSF-Tet containing a replication origin of a broad-host-range plasmid RSF1010 and a tetracycline resistance gene.

FIG. 8 shows construction of a plasmid RSFCPG containing a replication origin of a broad-host-range plasmid RSF1010, a tetracycline resistance gene, a *gltA* gene, a *ppc* gene and a *gdhA* gene.

FIG. 9 shows the construction of plasmid pSTVCB containing a *gltA* gene.

DETAILED DESCRIPTION OF THE INVENTION

Hereafter, the present invention will be explained in detail.

The production method according to the present invention is a method for producing L-glutamic acid by fermentation, which comprises culturing a microorganism having L-glutamic acid-producing ability in a liquid medium of which pH is adjusted to a condition under which L-glutamic acid produced by the microorganism is allowed to be precipitated, to allow L-glutamic acid to be produced and accumulated with precipitation of L-glutamic acid accompanied, wherein an operation causing existence of

L-glutamic acid crystals in the medium is performed when a concentration of L-glutamic acid in the medium is lower than the concentration at which spontaneous crystallization occurs.

The terms "spontaneous crystallization" used herein means that due to accumulation of L-glutamic acid by the microorganism having L-glutamic acid-producing ability, a concentration of L-glutamic acid in the medium exceeds a saturation concentration and L-glutamic acid spontaneously precipitates in the medium.

The operation causing existence of L-glutamic acid crystals in the medium means an operation by which existence of the crystals is artificially caused. Examples of the operation include addition of the crystals to the medium, and forced precipitation by lowering, during culture, pH of a medium in which a certain amount of L-glutamic acid has been dissolved at the beginning of the culture.

The amount of crystals to be existed in the medium is usually 0.01 to 10 g/L. The time at which the crystals are to be existed is preferably when the accumulated amount of L-glutamic acid in the medium increases to around the saturation concentration (for example, 25 g/L or more at pH 4.5). The amount of crystals existing in the medium and the concentration of L-glutamic acid can be determined by methods known to one skilled in the art. The existing amount of the crystals of L-glutamic acid may be determined by allowing the medium to stand and isolating the crystals from the medium by decantation. The concentration of L-glutamic acid in the medium means a concentration of dissolved L-glutamic acid. When crystals precipitate in the medium, the concentration is a determined concentration of a clarified solution obtained by separating solids by centrifugation (or filtration).

The operation causing existence of L-glutamic acid crystals is preferably addition of L-glutamic acid crystals.

As for the L-glutamic acid crystals, there are α -form and β -form of crystals (H. Takahashi, T. Takenishi, N. Nagashima Bull. Chem. Soc. Japan, 35, 923 (1962); J. D. Bernal, Z. Krist., 78, 363 (1931); S. Hirokawa, Acta Cryst, 8, 637 (1955)). When it is intended to obtain α -form of crystals, the crystals to be added are preferably of α -form.

A preferred amount of crystals varies depending on conditions such as crystal form of crystals. If the crystals are of α -form, it is usually 0.2 g/L or more. If it is more than this concentration, crystals of α -form may be obtained with good reproducibility. Because of their shape, the crystals of α -form may be handled more easily compared with crystals of β -form.

The microorganism having L-glutamic acid-producing ability used in the first production method of the present invention and the second production method of the present invention is a microorganism that accumulates a significant amount of L-glutamic acid in a medium when it is cultured in the medium. Examples thereof include microorganisms belonging to the genus **[Enterobacter]** *Pantoea*. Preferred is **[Enterobacter agglomerans]** *Pantoea ananatis*.

Further, the microorganism having L-glutamic acid-producing ability used in the production method of the present invention is preferably a microorganism that can metabolize a carbon source in a liquid medium containing L-glutamic acid at a saturation concentration and the carbon source, at a specific pH, and has an ability to accumulate L-glutamic acid in an amount exceeding the saturation concentration of L-glutamic acid in the liquid medium at the aforementioned pH (henceforth also referred to as "L-glutamic acid-accumulating microorganism"). The

5

aforementioned specific pH is preferably a pH at which L-glutamic acid precipitates in the medium, and such a pH is usually 5.0 or less.

The "saturation concentration" means a concentration of L-glutamic acid dissolved in the liquid medium when the liquid medium is saturated with L-glutamic acid.

When an L-glutamic acid-accumulating microorganism is used, the pH suitable for the production of L-glutamic acid is preferably a pH at which L-glutamic acid precipitates in the medium. By performing the culture at this pH, L-glutamic is produced and accumulated in the medium with its precipitation accompanied.

The L-glutamic acid-accumulating microorganism can be obtained as follows. A sample containing microorganisms is inoculated into a liquid medium containing L-glutamic acid at a saturation concentration and a carbon source, at a specific pH, and a strain that metabolizes the carbon source is selected. Although the specific pH is not particularly limited, it is usually about 5.0 or less, preferably about 4.5 or less, further preferably about 4.3 or less. The L-glutamic acid-accumulating microorganism is used for production of L-glutamic acid by fermentation with precipitation of the L-glutamic acid accompanied. If the pH is too high, it becomes difficult to allow the microorganism to produce L-glutamic acid in an amount sufficient for precipitation. Therefore, pH is preferably in the aforementioned range.

If pH of an aqueous solution containing L-glutamic acid is lowered, the solubility of L-glutamic acid significantly falls around pKa of γ -carboxyl group (4.25, 25° C.). The solubility becomes the lowest at the isoelectric point (pH 3.2) and L-glutamic acid exceeding the amount corresponding to the saturation concentration is precipitated. While it depends on the medium composition, L-glutamic acid is dissolved in an amount of 10-20 g/L at pH 3.2, 30-40 g/L at pH 4.0 and 50-60 g/L at pH 4.7, at about 30° C. Usually pH does not need to be made 3.0 or lower, because the L-glutamic acid precipitating effect reaches its upper limit when pH goes below a certain value. However, pH may be 3.0 or less.

In addition, the expression that a microorganism "can metabolize a carbon source" means that it can proliferate or can consume a carbon source even though it cannot proliferate, that is, it indicates that it catabolizes a carbon source such as sugars or organic acids. Specifically, for example, if a microorganism proliferates when it is cultured in a liquid medium containing L-glutamic acid at a saturation concentration at pH 5.0 to 4.0, preferably pH 4.5 to 4.0, more preferably pH 4.3 to 4.0, particularly preferably pH 4.0, at an appropriate temperature, for example, 28° C., 37° C. or 50° C., for 2 to 4 days, this microorganism can metabolize the carbon source in the medium. Further, for example, if a microorganism consumes a carbon source even though the microorganism does not proliferate, when it is cultured in a synthetic liquid medium containing L-glutamic acid at a saturation concentration at pH 5.0 to 4.0, preferably pH 4.5 to 4.0, more preferably pH 4.3 to 4.0, particularly preferably pH 4.0, at an appropriate temperature, for example, 28° C., 37° C. or 50° C., for 2 to 4 days, the microorganism is a microorganism that can metabolize the carbon source in the medium.

The microorganism that can metabolize a carbon source include a microorganism that can grow in the aforementioned liquid medium.

Further, the expression that a microorganism "can grow" means that it can proliferate or can produce L-glutamic acid even though it cannot proliferate. Specifically, for example, if a microorganism proliferates when it is cultured in a liquid

6

medium containing L-glutamic acid at a saturation concentration at pH 5.0 to 4.0, preferably pH 4.5 to 4.0, more preferably pH 4.3 to 4.0, particularly preferably pH 4.0, at an appropriate temperature, for example, 28° C., 37° C. or 50° C., for 2 to 4 days, this microorganism can grow in the medium. Further, for example, if a microorganism increases an amount of L-glutamic acid in a synthetic liquid medium even though the microorganism does not proliferate, when the microorganism is cultured in the synthetic liquid medium containing L-glutamic acid at a saturation concentration at pH 5.0 to 4.0, preferably pH 4.5 to 4.0, more preferably pH 4.3 to 4.0, particularly preferably pH 4.0, at an appropriate temperature, for example, 28° C., 37° C. or 50° C., for 2 to 4 days, this microorganism is a microorganism that can grow in the medium.

The selection described above may be repeated two or more times under the same conditions or with changing pH or the concentration of L-glutamic acid. A selection for an early stage can be performed in a medium containing L-glutamic acid at a concentration lower than the saturation concentration, and thereafter a subsequent selection can be performed in a medium containing L-glutamic acid at a saturation concentration. Further, strains with favorable properties such as superior proliferation rate may be selected.

The L-glutamic acid-accumulating microorganism is a microorganism that has an ability to accumulate L-glutamic acid in an amount exceeding the amount corresponding to the saturation concentration of L-glutamic acid in a liquid medium, in addition to the properties described above. The pH of the aforementioned liquid medium is preferably the same as or close to that of the medium used for screening a microorganism having the aforementioned properties. Usually, a microorganism becomes susceptible to L-glutamic acid at a high concentration as pH becomes lower. Therefore, it is preferred that pH is not low in view of resistance to L-glutamic acid, but low pH is preferred in view of production of L-glutamic acid with its precipitation accompanied. To satisfy these conditions, pH can be in the range of 3 to 5, preferably 4 to 5, more preferably 4 to 4.7, further preferably 4 to 4.5, particularly preferably 4.0 to 4.3.

As the L-glutamic acid-accumulating microorganism of or breeding materials therefor, there can be mentioned, for example, microorganisms belonging to the genus *Enterobacter*, *Klebsiella*, *Serratia*, *Pantoea*, *Erwinia*, *Escherichia*, *Corynebacterium*, *Alicyclobacillus*, *Bacillus*, *Saccharomyces* or the like. Among these, microorganisms belonging to the genus *Enterobacter* are preferred. Hereafter, the microorganism of the present invention will be explained mainly for microorganisms belonging to the genus *Enterobacter*. However, the microorganism is not limited to those belonging to the genus *Enterobacter*, and those belonging to other genera can be similarly used.

As a microorganism belonging to the [*Enterobacter*] *Pantoea*, there can be specifically mentioned [*Enterobacter* agglomerans] *Pantoea ananatis*, preferably the [*Enterobacter* agglomerans] *Pantoea ananatis* AJ13355 strain. This strain was isolated from soil in Iwata-shi, Shizuoka, Japan as a strain that can proliferate in a medium containing L-glutamic acid and a carbon source at low pH.

The physiological properties of AJ13355 are shown below:

- (1) Gram staining: negative
- (2) Behavior against oxygen: facultative anaerobic
- (3) Catalase: positive
- (4) Oxidase: negative
- (5) Nitrate-reducing ability: negative

- (6) Voges-Proskauer test: positive
 (7) Methyl Red test: negative
 (8) Urease: negative
 (9) Indole production: positive
 (10) Motility: motile
 (11) H₂S production in TSI medium: weakly active
 (12) β-Galactosidase: positive
 (13) Saccharide-assimilating property:
 Arabinose: positive
 Sucrose: positive
 Lactose: positive
 Xylose: positive
 Sorbitol: positive
 Inositol: positive
 Trehalose: positive
 Maltose: positive
 Glucose: positive
 Adonitol: negative
 Raffinose: positive
 Salicin: negative
 Melibiose: positive
 (14) Glycerose-assimilating property: positive
 (15) organic acid-assimilating property:
 Citric acid: positive
 Tartaric acid: negative
 Gluconic acid: positive
 Acetic acid: positive
 Malonic acid: negative
 (16) Arginine dehydratase: negative
 (17) Ornithine decarboxylase: negative
 (18) Lysine decarboxylase: negative
 (19) Phenylalanine deaminase: negative
 (20) Pigment formation: yellow
 (21) Gelatin liquefaction ability: positive
 (22) Growth pH: growth possible at pH 4, good growth at pH 4.5 to 7
 (23) Growth temperature: good growth at 25° C., good growth at 30° C., good growth at 37° C., growth possible at 42° C., growth impossible at 45° C.

Based on these bacteriological properties, AJ13355 was determined as [Enterobacter agglomerans] *Pantoea ananatis*.

The [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 was deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry (now, International Patent Organism Depositary, National Institute of Advanced Industrial Science and Technology) on Feb. 19, 1998 and received an accession number of FERM P-16644. It was then transferred to an international deposition under the provisions of Budapest Treaty on Jan. 11, 1999 and received an accession number of FERM BP-6614.

The L-glutamic acid-accumulating microorganism may be a microorganism originally having L-glutamic acid-producing ability or one having L-glutamic acid-producing ability imparted or enhanced by breeding through use of mutagenesis treatment, recombinant DNA techniques or the like.

The L-glutamic acid-producing ability can be imparted or enhanced by, for example, increasing activity of an enzyme

that catalyzes a reaction for biosynthesis of L-glutamic acid. The L-glutamic acid-producing ability can also be enhanced by decreasing or eliminating activity of an enzyme that catalyzes a reaction which branches off from the biosynthetic pathway of L-glutamic acid and generates a compound other than L-glutamic acid.

As examples of the enzyme that catalyzes the reaction for biosynthesis of L-glutamic acid, there can be mentioned glutamate dehydrogenase (hereafter, also referred to as "GDH"), glutamine synthetase, glutamate synthase, isocitrate dehydrogenase, aconitate hydratase, citrate synthase (hereafter, also referred to as "CS"), phosphoenolpyruvate carboxylase (hereafter, also referred to as "PEPC"), pyruvate dehydrogenase, pyruvate kinase, enolase, phosphoglyceromutase, phosphoglycerate kinase, glyceraldehyde-3-phosphate dehydrogenase, triosephosphate isomerase, fructose bisphosphate aldolase, phosphofructokinase, glucose phosphase isomerase and so forth. Among these enzymes, one, two or three of CS, PEPC and GDH are preferred. Further, it is preferred that the activities of all the three enzymes, CS, PEPC and GDH, are enhanced in the L-glutamic acid-accumulating microorganism. In particular, CS of *Brevibacterium lactofermentum* is preferred, because it does not suffer from inhibition by α-ketoglutaric acid, L-glutamic acid and NADH.

In order to enhance the activity of CS, PEPC or GDH, for example, a gene coding for CS, PEPC or GDH can be cloned on an appropriate plasmid and a host microorganism can be transformed with the obtained plasmid. The copy number of the gene coding for CS, PEPC or GDH (hereafter, abbreviated as "gltA gene", "ppc gene" and "gdhA gene", respectively) in the transformed strain cell increases, resulting in the increase of the activity of CS, PEPC or GDH.

The cloned gltA, ppc and gdhA genes are introduced into the aforementioned starting parent strain solely or in combination of arbitrary two or three kinds of them. When two or three kinds of the genes are introduced, two or three kinds of the genes may be cloned on one kind of plasmid and introduced into the host, or separately cloned on two or three kinds of plasmids that can coexist and introduced into the host.

Two or more kinds of genes coding for an enzyme of the same kind, but derived from different microorganisms, may be introduced into the same host.

The plasmids described above are not particularly limited so long as they are autonomously replicable in a cell of a microorganism belonging to, for example, the genus *Enterobacter* or the like. However, there can be mentioned, for example, pUC19, pUC18, pBR322, pHSG299, pHSG298, pHSG399, pHSG398, RSF1010, pMW119, pMW118, pMW219, pMW218, pACYC177, pACYC184 and so forth. Besides these, vectors of phage DNA can also be used.

Transformation can be performed by, for example, the method of D. M. Morrison (Methods in Enzymology, 68, 326 (1979)), the method wherein permeability of recipient bacterium cells for DNA is increased by treating the cells with calcium chloride (Mandel M. and Higa A., J. Mol. Biol., 53, 159 (1970)), electroporation (Miller J. H., "A Short Course in Bacterial Genetics", Cold Spring Harbor Laboratory Press, U.S.A., 1992) or the like.

The activity of CS, PEPC or GDH can also be increased by allowing multiple copies of the gltA gene, the ppc gene or the gdhA gene to be present on chromosomal DNA of the aforementioned starting parent strain to be a host. In order to introduce multiple copies of the gltA gene, the ppc gene or the gdhA gene on chromosomal DNA of a microorganism

belonging to the genus *Enterobacter* or the like, a sequence of which multiple copies are present on the chromosomal DNA, such as repetitive DNA and inverted repeats present at terminus of a transposable element, can be used. Alternatively, multiple copies of the genes can be introduced onto chromosomal DNA by utilizing transfer of a transposon containing the *gltA* gene, the *ppc* gene or the *gdhA* gene. As a result, the copy number of *gltA* gene, the *ppc* gene or the *gdhA* gene in a transformed strain cell is increased, and thus the activity of CS, PEPC or GDH is increased.

As organisms used as a source of the *gltA* gene, the *ppc* gene or the *gdhA* gene of which copy number is to be increased, any organism can be used so long as it has activity of CS, PEPC or GDH. Inter alia, bacteria, which are prokaryotes, for example, those belonging to the genus *Enterobacter*, *Klebsiella*, *Erwinia*, *Pantoea*, *Serratia*, *Escherichia*, *Corynebacterium*, *Brevibacterium* or *Bacillus* are preferred. As specific examples, there can be mentioned *Escherichia coli*, *Brevibacterium lactofermentum* and so forth. The *gltA* gene, the *ppc* gene and the *gdhA* gene can be obtained from chromosomal DNA of the microorganisms described above.

The *gltA* gene, the *ppc* gene and the *gdhA* gene can be obtained by using a mutant strain which is deficient in the activity of CS, PEPC or GDH to isolate a DNA fragment that supplements its auxotrophy from chromosomal DNA of the aforementioned microorganism. Further, since the nucleotide sequences of these genes of *Escherichia* and *Corynebacterium* bacteria have already been elucidated (Biochemistry, 22, pp. 5243-5249, (1983); J. Biochem., 95, pp. 909-916, (1984); Gene, 27, pp. 193-199, (1984); Microbiology, 140, pp. 1817-1828, (1994); Mol. Gen. Genet., 218, pp. 330-339, (1989); Molecular Microbiology, 6, pp. 317-326, (1992)), they can also be obtained by PCR utilizing primers synthesized based on each nucleotide sequence and chromosomal DNA as a template.

The activity of CS, PEPC or GDH can also be increased by enhancing the expression of the *gltA* gene, the *ppc* gene or the *gdhA* gene, besides the aforementioned amplification of the genes. For example, the expression can be enhanced by replacing a promoter for the *gltA* gene, the *ppc* gene or the *gdhA* gene with another stronger promoter. For example, *lac* promoter, *trp* promoter, *trc* promoter, *tac* promoter, P_R promoter and P_L promoter of the lambda phage and so forth are known as strong promoters. The *gltA* gene, the *ppc* gene and the *gdhA* gene of which promoter is replaced are cloned on a plasmid and introduced into the host microorganism, or introduced onto the chromosomal DNA of the host microorganism by using repetitive DNA, inverted repeat, transposon or the like.

The activity of CS, PEPC or GDH can also be increased by replacing the promoter of the *gltA* gene, the *ppc* gene or the *gdhA* gene on the chromosome with another stronger promoter (see WO87/03006 and Japanese Patent Application Laid-open No. 61-268183), or inserting a strong promoter in the upstream of the coding sequence of each gene (see Gene, 29, pp. 231-241 (1984)). Specifically, homologous recombination can be performed between the *gltA* gene, the *ppc* gene or the *gdhA* gene of which promoter is replaced with a stronger one or DNA containing a part thereof and the corresponding gene on the chromosome.

Examples of the enzyme that catalyzes the reaction which branches off from the biosynthetic pathway of the L-glutamic acid and generates a compound other than L-glutamic acid include α -ketoglutarate dehydrogenase (hereafter, also referred to as " α KGDH"), isocitrate lyase,

phosphate acetyltransferase, acetate kinase, acetohydroxy acid synthase, acetolactate synthase, formate acetyltransferase, lactate dehydrogenase, glutamate decarboxylase, 1-pyrroline dehydrogenase and so forth. Among these enzymes, α KGDH is preferred.

In order to decrease or eliminate the activities of the aforementioned enzymes in a microorganism belonging to the genus *Enterobacter* or the like, mutations for decreasing or eliminating the intracellular activity of the enzymes can be introduced into genes of the aforementioned enzymes by a usual mutagenesis treatment method or a genetic engineering method.

Examples of the mutagenesis treatment method include, for example, methods utilizing irradiation with x-ray or ultraviolet ray, methods utilizing treatment with a mutagenesis agent such as N-methyl-N'-nitro-N-nitrosoguanidine, and so forth. The site of a gene where the mutation is introduced may be in a coding region coding for an enzyme protein or a region for regulating expression such as a promoter.

Examples of the genetic engineering methods include, for example, methods utilizing gene recombination, transduction, cell fusion and so forth. For example, a drug resistance gene is inserted into a cloned target gene to prepare a gene that has lost its function (defective gene). Subsequently, this defective gene is introduced into a cell of a host microorganism, and the target gene on the chromosome is replaced with the aforementioned defective gene by utilizing homologous recombination (gene disruption).

Decrease or deficiency of intracellular activity of the target enzyme and the degree of decrease of the activity can be confirmed by measuring the enzyme activity of a cell extract or a purified fraction thereof obtained from a candidate strain and comparing it with that of a wild strain. For example, the α KGDH activity can be measured by the method of Reed et al. (Reed L. J. and Mukherjee B. B., Methods in Enzymology, 13, pp. 55-61 (1969)).

Depending on the target enzyme, a target mutant strain can be selected based on a phenotype of the mutant strain. For example, a mutant strain wherein the α KGDH activity is eliminated or decreased cannot proliferate or shows a markedly reduced proliferation rate in a minimal medium containing glucose or a minimal medium containing acetic acid or L-glutamic acid as an exclusive carbon source under an aerobic culture condition. However, normal proliferation is enabled even under the same condition by adding succinic acid or lysine, methionine and diaminopimelic acid to a minimal medium containing glucose. By utilizing these phenomena as indicators, a mutant strain with decreased α KGDH activity or deficient in the activity can be selected.

A method for preparing an α KGDH gene-deficient strain of *Brevibacterium lactofermentum* by utilizing homologous recombination is described in detail in WO95/34672. Similar methods can be applied to other microorganisms.

Further, techniques such as cloning of genes and digestion and ligation of DNA, transformation and so forth are described in detail in Molecular Cloning, 2nd Edition, Cold Spring Harbor Press (1989) and so forth.

As a specific example of a mutant strain deficient in α KGDH activity or with decreased α KGDH activity obtained as described above, there can be mentioned [*Enterobacter agglomerans*] *Pantoea ananatis* AJ13356. [*Enterobacter agglomerans*] *Pantoea ananatis* AJ13356 was deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry (now, International Patent Organism Depository, National

Institute of Advanced Industrial Science and Technology) on Feb. 19, 1998 and received an accession number of FERM P-16645. It was then transferred to an international deposition under the provisions of Budapest Treaty on Jan. 11, 1999 and received an accession number of FERM BP-6615. The [Enterobacter agglomerans] *Pantoea ananatis* AJ13356 is deficient in α KGDH activity as a result of disruption of the α KGDH-E1 subunit gene (sucA).

When [Enterobacter agglomerans] *Pantoea ananatis*, which is an example of the microorganism used in the present invention, is cultured in a medium containing a saccharide, mucus is extracellularly secreted, occasionally resulting in low operation efficiency. Therefore, when [Enterobacter agglomerans] *Pantoea ananatis* having such a property of secreting mucus is used, it is preferable to use a mutant strain that secretes less mucus compared with a wild strain. Examples of mutagenesis treatment include, for example, methods utilizing irradiation with X-ray or ultraviolet ray, method utilizing treatment with a mutagenesis agent such as N-methyl-N'-nitro-N-nitrosoguanidine, and so forth. A mutant strain with decreased secretion of mucus can be selected by inoculating mutagenized bacterial cells in a medium containing a saccharide, for example, LB medium plate containing 5 g/L of glucose, culturing them with tilting the plate about 45 degrees and selecting a colony that does not show flowing down of mucus.

In the present invention, impartation or enhancement of L-glutamic acid-producing ability and impartation of other favorable properties such as mutation for less mucus secretion described above can be carried out in an arbitrary order.

By culturing the microorganism having L-glutamic acid-producing ability in a liquid medium that is adjusted to pH condition that allows precipitation of L-glutamic acid, L-glutamic acid can be produced and accumulated with its precipitation in the medium accompanied.

Preferably, by culturing the L-glutamic acid-accumulating microorganism in a liquid medium that is adjusted to pH condition that allows precipitation of L-glutamic acid, L-glutamic acid can be produced and accumulated with its precipitation in the medium accompanied. Furthermore, it is possible that the culture is started at a neutral pH, and pH becomes the condition that allows precipitation of L-glutamic acid when the culture is completed.

The "condition that allows precipitation of L-glutamic acid" referred to herein means a condition that allows precipitation of L-glutamic acid when the above-mentioned microorganism produces and accumulates L-glutamic acid. For example, it is usually 3 to 5 when the microorganism is an Enterobacter bacterium.

The microorganism may be cultured at pH suitable for growth thereof at the beginning and then cultured under the condition which allows precipitation of L-glutamic acid. For example, when the medium contains a sugar source which the microorganism can not assimilate under the condition which allows precipitation of L-glutamic acid, or an organic acid which inhibits the growth of the microorganism under the condition which allows precipitation of L-glutamic acid, the microorganism may be cultured under a condition under which the microorganism can assimilate the sugar source or growth of the microorganism is not inhibited by the organic acid to allow the microorganism to consume the sugar source or the organic acid, and then cultured under the condition allows precipitation of L-glutamic acid.

As the media used for culture, a usual nutrient medium containing a carbon source, a nitrogen source, mineral salts and organic trace nutrients such as amino acids and vitamins

as required can be used so long as pH is adjusted so as to satisfy the predetermined condition. Either a synthetic medium or a natural medium can be used. The carbon source and the nitrogen source used in the medium can be any ones so long as they can be used by the strain to be cultured.

As the carbon source, saccharides such as glucose, glycerol, fructose, sucrose, maltose, mannose, galactose, starch hydrolysate and molasses are used. In addition, organic acids such as acetic acid and citric acid may be used each alone or in combination with another carbon source.

As the nitrogen source, ammonia, ammonium salts such as ammonium sulfate, ammonium carbonate, ammonium chloride, ammonium phosphate and ammonium acetate, nitrates and so forth are used.

As the organic trace nutrients, amino acids, vitamins, fatty acids, nucleic acids, those containing these substances such as peptone, casamino acid, yeast extract and soybean protein decomposition products are used. When an auxotrophic mutant strain that requires an amino acid and so forth for metabolism or growth is used, the required nutrient must be supplemented.

As mineral salts, phosphates, magnesium salts, calcium salts, iron salts, manganese salts and so forth are used.

The culture is usually performed with aeration under the condition of a culture temperature at 20 to 42° C., and pH at 3 to 5, preferably 4 to 5, more preferably 4 to 4.7, particularly preferably 4 to 4.5. A considerable amount of L-glutamic acid is usually accumulated after culture of from about 10 hours to about 4 days. A portion of the accumulated L-glutamic acid which exceeds the saturation concentration precipitates in the medium.

After completion of the culture, L-glutamic acid precipitated in the culture can be collected by centrifugation, filtration or the like. L-Glutamic acid dissolved in the medium can be also collected by known methods. For example, the L-glutamic acid can be isolated by concentrating the culture broth to crystallize it or isolated by ion exchange chromatography or the like. It is also possible to crystallize L-glutamic acid dissolved in the medium and then collect the L-glutamic acid precipitated in the culture broth together with the crystallized L-glutamic acid.

When L-glutamic acid exceeding a saturation concentration precipitates, the concentration of L-glutamic acid dissolved in the medium is maintained at a constant level. Therefore, influence of L-glutamic acid at a high concentration on microorganisms can be reduced. Accordingly, it also becomes possible to breed a microorganism having further improved L-glutamic acid-producing ability. Further, since L-glutamic acid is precipitated as crystals, acidification of the culture broth by accumulation of L-glutamic acid is suppressed, and therefore the amount of alkali used for maintaining pH of the culture can significantly be reduced. Provided that the present invention is not intended to be bound by any theories, the reason why the yield of L-glutamic acid is improved by performing an operation causing existence of L-glutamic acid crystals in the medium when a concentration of L-glutamic acid in the medium is lower than the concentration at which spontaneous crystallization occurs, in the case that L-glutamic acid precipitates as described above, is considered as follows.

In the fermentation of L-glutamic acid with precipitation accompanied, production of L-glutamic acid tends to reach its limit when spontaneous crystallization occurs. It is presumed that this is because a concentration of L-glutamic acid is the highest on the surfaces of cells when the spontaneous crystallization occurs, and precipitation occurs on the

surfaces, thereby hindering movement of substances around a membrane to reduce activity of a bacterium. If existence of crystals is artificially caused before the spontaneous crystallization occurs, the crystals serve as seed crystals to cause precipitation on surfaces of the crystals, thereby preventing the bacterium from the activity reduction.

EXAMPLES

Hereafter, the present invention will be more specifically explained with reference to the following examples. In the examples, amino acids are L-amino acids unless otherwise indicated.

Reference Example 1

<1> Screening of Microorganism Having L-glutamic Acid Resistance in Acidic Environment

Screening of a microorganism having L-glutamic acid resistance in acidic environment was performed as follows. One (1) g each of about 500 samples obtained from nature including soil, fruits, plant bodies, river water and so forth was suspended in 5 mL of sterilized water, and 200 μ L thereof was coated on 20 mL of solid medium adjusted to pH 4.0 with HCl. The composition of the medium was as follows: 3 g/L of glucose, 1 g/L of ammonium sulfate, 0.2 g/L of magnesium sulfate heptahydrate, 0.5 g/L of potassium dihydrogenphosphate, 0.2 g/L of sodium chloride, 0.1 g/L of calcium chloride dihydrate, 0.01 g/L of ferrous sulfate heptahydrate, 0.01 g/L of manganese sulfate tetrahydrate, 0.72 mg/L of zinc sulfate dihydrate, 0.64 mg/L of copper sulfate pentahydrate, 0.72 mg/L of cobalt chloride hexahydrate, 0.4 mg/L of boric acid, 1.2 mg/L of sodium molybdate dihydrate, 50 μ g/L of biotin, 50 μ g/L of calcium pantothenate, 50 μ g/L of folic acid, 50 μ g/L of inositol, 50 μ g/L of niacin, 50 μ g/L of p-aminobenzoic acid, 50 μ g/L of pyridoxine hydrochloride, 50 μ g/L of riboflavin, 50 μ g/L of thiamine hydrochloride, 50 mg/L of cycloheximide and 20 g/L of agar.

The media plated with the above samples were incubated at 28° C., 37° C. or 50° C. for 2 to 4 days and 378 strains forming colonies were obtained.

Subsequently, each of the strains obtained as described above was inoculated in a test tube of 16.5 cm in length and 14 mm in diameter containing 3 mL of liquid medium (adjusted to pH 4.0 with HCl) containing a saturation concentration of L-glutamic acid and cultured at 28 C., 37° C. or 50° C. for 24 hours to 3 days with shaking. Then, the grown strains were selected. The composition of the aforementioned medium was follows: 40 g/L of glucose, 20 g/L of ammonium sulfate, 0.5 g/L of magnesium sulfate heptahydrate, 2 g/L of potassium dihydrogenphosphate, 0.5 g/L of sodium chloride, 0.25 g/L of calcium chloride dihydrate, 0.02 g/L of ferrous sulfate heptahydrate, 0.02 g/L of manganese sulfate tetrahydrate, 0.72 mg/L of zinc sulfate dihydrate, 0.64 mg/L of copper sulfate pentahydrate, 0.72 mg/L of cobalt chloride hexahydrate, 0.4 mg/L of boric acid, 1.2 mg/L of sodium molybdate dihydrate and 2 g/L of yeast extract.

Thus, 78 strains of microorganisms showing L-glutamic acid resistance in an acidic environment were successfully obtained.

<2> Selection of Strains Showing Superior Growth Rate From Microorganisms Having L-glutamic Acid Resistance in Acidic Environment

The various microorganisms having L-glutamic acid resistance in an acidic environment obtained as described above are each inoculated into a test tube of 16.5 cm in length and 14 mm in diameter containing 3 mL of medium

(adjusted to pH 4.0 with HCl) obtained by adding 20 g/L of glutamic acid and 2 g/L of glucose to M9 medium (Sambrook, J., Fritsh, E. F. and Maniatis, T., "Molecular Cloning", Cold Spring Harbor Laboratory Press, U.S.A., 1989), and the turbidity of the medium was measured in the time course to select strains showing a favorable growth rate. As a result, as a strain showing favorable growth, the AJ13355 strain was obtained from soil in Iwata-shi, Shizuoka, Japan. This strain was determined as [Enterobacter agglomerans] *Pantoea ananatis* based on its bacteriological properties described above.

<3> Acquisition of Strain with Less Mucus Secretion From [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 Strain

Since the [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 strain extracellularly secretes mucus when cultured in a medium containing a saccharide, operation efficiency is not favorable. Therefore, a strain with less mucus secretion was obtained by the ultraviolet irradiation method (Miller, J. H. et al., "A Short Course in Bacterial Genetics; Laboratory Manual", p. 150, 1992, Cold Spring Harbor Laboratory Press, U.S.A.).

The [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 strain was irradiated with ultraviolet ray for 2 minutes at a position 60 cm away from a 60-W ultraviolet lamp and cultured in LB medium overnight to fix mutation. The mutagenized strain was diluted and inoculated in LB medium containing 5 g/L of glucose and 20 g/L of agar so that about 100 colonies per plate would emerge and cultured at 30° C. overnight with tilting the plate about 45 degrees, and then 20 colonies showing not flowing down of mucus were selected.

As a strain satisfying conditions that no revertant emerged even after 5 times of subculture in LB medium containing 5 g/L of glucose and 20 g/L of agar, and that there should be observed growth equivalent to the parent strain in LB medium, LB medium containing 5 g/L of glucose and M9 medium (Sambrook, J. et al., Molecular Cloning, 2nd Edition, Cold Spring Harbor Press, U.S.A., 1989) supplemented with 20 g/L of L-glutamic acid and 2 g/L of glucose and adjusted to pH 4.5 with HCl, SC17 strain was selected from the strains selected above.

<4> Construction of Glutamic Acid-Producing Bacterium From [Enterobacter agglomerans] *Pantoea ananatis* SC17 Strain

(1) Preparation of α KGDH Deficient Strain From [Enterobacter agglomerans] *Pantoea ananatis* SC17 Strain

A strain that was deficient in α KGDH and had enhanced L-glutamic acid biosynthetic system was prepared from the [Enterobacter agglomerans] *Pantoea ananatis* SC17 strain.

(i) Cloning of α KGDH Gene (Hereafter, Referred to as "sucAB") of [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 Strain

The sucAB gene of the [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 strain was cloned by selecting a DNA fragment complementing the acetic acid-unassimilating property of the α KGDH-E1 subunit gene (hereafter, referred to as "sucA")-deficient strain of *Escherichia coli* from chromosomal DNA of the [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 strain.

The chromosomal DNA of the [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 strain was isolated by a method usually employed for extracting chromosomal DNA from *Escherichia coli* (Text for Bioengineering Experiments, Edited by the Society for Bioscience and Bioengineering, Japan, pp. 97-98, Baifukan, 1992). The pTWV228 (resistant to ampicillin) used as a vector was a commercial product of Takara Shuzo Co., Ltd.

The chromosomal DNA of the AJ13355 strain digested with EcoT221 and pTWV228 digested with PstI were ligated by using T4 ligase and used to transform the sucA-deficient Escherichia coli JRG465 strain (Herbert, J. et al., Mol. Gen. Genetics, 105, 182 (1969)). A strain growing in an acetate minimal medium was selected from the transformant strains obtained above, and a plasmid was extracted from it and designated as pTWVEK101. The Escherichia coli JRG465 strain harboring pTWVEK101 recovered auxotrophy for succinic acid or L-lysine and L-methionine besides the trait of acetic acid-unassimilating property. This suggests that pTWVEK101 contained the sucA gene of [Enterobacter agglomerans] *Pantoea ananatis*.

FIG. 1 shows a restriction enzyme map of a DNA fragment derived from [Enterobacter agglomerans] *Pantoea ananatis* in pTWVEK101. In the nucleotide sequence of the hatched portion in FIG. 1, nucleotide sequences considered to be two full length ORFs and two nucleotide sequences considered to be partial sequences of ORFs were found. As a result of homology search for these, it was revealed that the portions of which nucleotide sequences were determined contained a 3' end partial sequence of the succinate dehydrogenase iron-sulfur protein gene (sdhB), full length sucA and α KGDH-E2 subunit gene (sucB gene), and a 5' end partial sequence of the succinyl CoA synthetase β subunit gene (sucC gene). The results of comparison of the amino acid sequences deduced from these nucleotide sequences with those derived from Escherichia coli (Eur. J. Biochem., 141, pp. 351-359 (1984); Eur. J. Biochem., 141, pp. 361-374 (1984); Biochemistry, 24, pp. 6245-6252 (1985)) are shown in FIGS. 2 to 5. Thus, the amino acid sequences showed very high homology to each other. In addition, it was found that a cluster of sdhB-sucA-sucB-sucC was constituted on the chromosome of [Enterobacter agglomerans] *Pantoea ananatis* as in Escherichia coli (Eur. J. Biochem., 141, pp. 351-359 (1984); Eur. J. Biochem., 141, pp. 361-374 (1984); Biochemistry, 24, pp. 6245-6252 (1985)).

(ii) Acquisition of α KGDH-deficient Strain Derived from [Enterobacter agglomerans] *Pantoea ananatis* SC17 Strain

The homologous recombination was performed by using the sucAB gene of [Enterobacter agglomerans] *Pantoea ananatis* obtained as described above to obtain an α KGDH-deficient strain of [Enterobacter agglomerans] *Pantoea ananatis*.

After pTWVEK101 was digested with SphI to excise a fragment containing sucA, the fragment was blunt-ended with Klenow fragment (Takara Shuzo Co., Ltd.) and ligated with pBR322 digested with EcoRI and blunt-ended with Klenow fragment, by using T4 DNA ligase (Takara Shuzo Co., Ltd.). The obtained plasmid was digested at the restriction enzyme BglIII recognition site positioned approximately at the center of sucA by using the enzyme, blunt-ended with Klenow fragment, and then ligated again by using T4 DNA ligase. It was considered that the sucA gene became unfunctional because a frameshift mutation was introduced into sucA of the plasmid newly constructed through the above procedure.

The plasmid constructed as described above was digested with a restriction enzyme ApaLI, and subjected to agarose gel electrophoresis to recover a DNA fragment containing sucA into which the frameshift mutation was introduced and a tetracycline resistance gene derived from pBR322. The recovered DNA fragment was ligated again by using T4 DNA ligase to construct a plasmid for disrupting the α KGDH gene.

The plasmid for disrupting the α KGDH gene obtained as described above was used to transform the [Enterobacter agglomerans] *Pantoea ananatis* SC17 strain by electroporation (Miller, J. H., "A Short Course in Bacterial Genetics; Handbook", p. 279, Cold Spring Harbor Laboratory Press, U.S.A., 1992), and a strain wherein sucA on the chromo-

some was replaced with a mutant type one of the plasmid by homologous recombination was obtained by using the tetracycline resistance as a marker. The obtained strain was designated as SC17sucA strain.

In order to confirm that the SC17sucA strain was deficient in the α KGDH activity, the enzyme activity was measured by the method of Reed et al. (Reed, L. J. and Mukherjee, B. B., Methods in Enzymology, 13, pp. 55-61, (1969)) by using cells of the strain cultured in LB medium to the logarithmic growth phase. As a result, α KGDH activity of 0.073 (Δ ABS/min/mg protein) was detected from the SC17 strain, whereas no α KGDH activity was detected from the SC17sucA strain, and thus it was confirmed that the sucA was eliminated as intended.

(2) Enhancement of L-glutamic Acid Biosynthesis System of [Enterobacter agglomerans] *Pantoea ananatis* SC17sucA Strain

Subsequently, the citrate synthase gene, phosphoenolpyruvate carboxylase gene and glutamate dehydrogenase gene derived from Escherichia coli were introduced into the SC17sucA strain.

(i) Preparation of Plasmid Having gltA Gene, ppc Gene and gdhA Gene derived From Escherichia coli

The procedures of preparing a plasmid having the gltA gene, the ppc gene and the gdhA gene will be explained by referring to FIGS. 6 and 7.

A plasmid having the gdhA gene derived from Escherichia coli, pBRGDH (Japanese Patent Application Laid-open No. 7-203980), was digested with HindIII and SphI, the both ends were blunt-ended by the T4 DNA polymerase treatment, and then the DNA fragment having the gdhA gene was purified and recovered. Separately, a plasmid having the gltA gene and ppc gene derived from Escherichia coli, pMWCP (WO97/08294), was digested with XbaI, and then the both ends were blunt-ended by using T4 DNA polymerase. This was mixed with the above purified DNA fragment having the gdhA gene and ligated by using T4 ligase to obtain a plasmid pMWCPG, which corresponded to pMWCP further containing the gdhA gene (FIG. 6).

At the same time, the plasmid pVIC40 (Japanese Patent Application Laid-open No. 8-047397) having the replication origin of the broad-host-range plasmid RSF1010 was digested with NotI, treated with T4 DNA polymerase and digested with PstI. pBR322 was digested with EcoT141, treated with T4 DNA polymerase and digested with PstI. The both products were mixed and ligated by using T4 ligase to obtain a plasmid RSF-Tet having the replication origin of RSF1010 and the tetracycline resistance gene (FIG. 7).

Subsequently, pMWCPG was digested with EcoRI and PstI, and a DNA fragment having the gltA gene, the ppc gene and the gdhA gene was purified and recovered. RSF-Tet was similarly digested with EcoRI and PstI, and a DNA fragment having the replication origin of RSF1010 was purified and recovered. The both products were mixed and ligated by using T4 ligase to obtain a plasmid RSFCPPG, which corresponded to RSF-Tet containing the gltA gene, the ppc gene and the gdhA gene (FIG. 8). It was confirmed that the obtained plasmid RSFCPPG expressed the gltA gene, the ppc gene and the gdhA gene based on the supplementation of the auxotrophy of the gltA gene-, ppc gene- or gdhA gene-deficient strain derived from Escherichia coli and measurement of each enzyme activity.

(ii) Preparation of Plasmid Having gltA Gene derived From Brevibacterium lactofermentum

A plasmid having the gltA gene derived from Brevibacterium lactofermentum was constructed as follows. PCR was performed by using the primer DNAs which were prepared based on the nucleotide sequence of the Corynebacterium glutaraicum gltA gene (Microbiology, 140, pp. 1817-1828 (1994)), and chromosomal DNA of Brevibacterium lactof-

ermertum ATCC13869 as a template to obtain a *gltA* gene fragment of about 3 kb. This fragment was inserted into a plasmid pHSG399 (purchased from Takara Shuzo Co., Ltd.) digested with *Sma*I to obtain a plasmid pHSGCB (FIG. 9). Subsequently, pHSGCB was digested with *Hind*III, and the excised *gltA* gene fragment of about 3 kb was inserted into a plasmid pSTV29 (purchased from Takara Shuzo Co., Ltd.) digested with *Hind*III to obtain a plasmid pSTVCB (FIG. 9). It was confirmed that the obtained plasmid pSTVCB expressed the *gltA* gene by measuring the enzyme activity in the [Enterobacter agglomerans] *Pantoea ananatis* AJ13355 strain.

(iii) Introduction of RSFCPG and pSTVCB into SC17sucA Strain

The [Enterobacter agglomerans] *Pantoea ananatis* SC17sucA strain was transformed with RSFCPG by electroporation to obtain a transformant SC17sucA/RSFCPG strain showing tetracycline resistance. Further, the SC17sucA/RSFCPG strain was transformed with pSTVCB by electroporation to obtain a transformant SC17sucA/RSFCPG+pSTVCB strain showing chloramphenicol resistance.

<5> Acquisition of Strain with Improved Resistance to L-glutamic Acid in Low pH Environment

A strain with improved resistance to L-glutamic acid at a high concentration in a low pH environment (hereafter, also referred to as "strain with high-concentration Glu-resistance at low pH") was isolated from the [Enterobacter agglomerans] *Pantoea ananatis* SC17sucA/RSFCPG+pSTVCB strain.

The SC17sucA/RSFCPG+pSTVCB strain was cultured overnight at 30° C. in LBG medium (10 g/L of trypton, 5 g/L of yeast extract, 10 g/L of NaCl, 5 g/L of glucose), and the cells washed with saline was appropriately diluted and plated on an M9-E medium (4 g/L of glucose, 17 g/L of Na₂HPO₄·12H₂O, 3 g/L of KH₂PO₄, 0.5 g/L of NaCl, 1 g/L of NH₄Cl, 10 MM of MgSO₄, 10 μM of CaCl₂, 50 mg/L of L-lysine, 50 mg/L of L-methionine, 50 mg/L of DL-diaminopimelic acid, 25 mg/L of tetracycline, 25 mg/L of chloramphenicol, 30 g/L of L-glutamic acid, adjusted to pH 4.5 with aqueous ammonia) plate. A colony emerged after culture at 32° C. for 2 days was obtained as a strain with high-concentration Glu-resistance at low pH.

For the obtained strain, growth level in M9-E liquid medium was measured and L-glutamic acid-producing ability was tested in a 50-ml volume large test tube containing 5 ml of L-glutamic acid production test medium (40 g/L of glucose, 20 g/L of ammonium sulfate, 0.5 g/L of magnesium sulfate heptahydrate, 2 g/L of potassium dihydrogenphosphate, 0.5 g/L of sodium chloride, 0.25 g/L of calcium chloride dihydrate, 0.02 g/L of ferrous sulfate heptahydrate, 0.02 g/L of manganese sulfate tetrahydrate, 0.72 mg/L of zinc sulfate dihydrate, 0.64 mg/L of copper sulfate pentahydrate, 0.72 mg/L of cobalt chloride hexahydrate, 0.4 mg/L of boric acid, 1.2 mg/L of sodium molybdate dihydrate, 2 g/L of yeast extract, 200 mg/L of L-lysine hydrochloride, 200 mg/L of L-methionine, 200 mg/L of DL-α,ε-diaminopimelic acid, 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol). A strain that exhibited the best growth level and the same L-glutamic acid-producing ability as that of its parent strain, the SC17/RSFCPG+pSTVCB strain, was designated as [Enterobacter Agglomerans] *Pantoea ananatis* AJ13601. The AJ13601 strain was deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Ministry of International Trade and Industry (now, International Patent Organism Depository,

National Institute of Advanced Industrial Science and Technology; Central 6, Higashi 1-1-1, Tsukuba-shi, Ibaraki 305-8566, Japan) on Aug. 18, 1999 and received an accession number of FERM P-17516. It was then transferred to an international deposition under the provisions of Budapest Treaty on Jul. 6, 2000 and received an accession number of FERM BP-7207.

Example 1

The [Enterobacter Agglomerans] *Pantoea ananatis* AJ13601 strain was cultured on LBG agar medium (10 g/L of trypton, 5 g/L of yeast extract, 10 g/L of NaCl and 15 g/L of agar) containing 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol at 30° C. for 14 hours, and the cells in one plate (diameter 8.5 cm) were collected and inoculated into 300 mL of a culture medium containing 50 g/L of glucose, 4 g/L of ammonium sulfate, 0.4 g/L of magnesium sulfate heptahydrate, 2 g/L of monopotassium dihydrogenphosphate, 10 mg/L of ferrous sulfate heptahydrate, 10 mg/L of manganese sulfate pentahydrate, 4 g/L of yeast extract, 400 mg/L of L-lysine hydrochloride, 400 mg/L of DL-methionine, 400 mg/L of DL-α,ε-diaminopimelic acid, 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol, in a 1 L-volume jar fermenter to perform culture at 34° C. at pH 6.0. Culture pH was controlled by adding ammonia gas. The culture was terminated about 16 hours after beginning of the culture, at a phase in which the glucose in the culture medium was depleted.

60 ml of the culture broth cultured as described above was inoculated into 300 mL of a main culture medium containing 50 g/L of glucose, 5 g/L of ammonium sulfate, 0.4 g/L of magnesium sulfate heptahydrate, 5 g/L of monopotassium dihydrogenphosphate, 20 mg/L of ferrous sulfate heptahydrate, 20 mg/L of manganese sulfate pentahydrate, 6 g/L of yeast extract, 800 mg/L of L-lysine hydrochloride, 600 mg/L of DL-methionine, 600 mg/L of DL-α,ε-diaminopimelic acid, 1.5 g/L of sodium chloride, 0.75 g/L of calcium chloride dihydrate, 25 mg/L of tetracycline hydrochloride and 25 mg/L of chloramphenicol, in a 1 L-volume jar to perform culture at 34° C. at pH 6.0. As L-glutamic acid accumulated, pH spontaneously decreased to reach pH 4.5 about one or two hours after the beginning of the culture. Thereafter, the culture pH was adjusted to pH 4.5 by adding ammonia gas. After the initially-added glucose was depleted, 700 g/L of an aqueous solution of glucose was continuously added at a rate of 6 ml/hr.

When the fermentative production of L-glutamic acid was continued as described above, the concentration of L-glutamic acid accumulated in the culture broth about 14 hours after the beginning of the culture was about 45 to 55 g/L.

At this time, L-glutamic acid crystals of α-form or β-form were added to the culture broth as dry crystals from the upper part of the jar fermenter, in an amount ranging from 0.008 to 1.0 g (0.022 to 2.778 g/L with respect to the medium amount at the beginning of the culture), and the culture was continued and terminated 36 hours after the beginning of the culture. A considerable amount of L-glutamic acid crystals precipitated in the jar fermenter. The crystals were separated by decantation, and dried overnight at room temperature. The crystal form (crystal lattice form) of the resultant dried crystals was investigated by X-ray powder diffraction method. The results are shown in Table 1. The result of a control experiment in which no crystal was added during the culture is also shown in Table 1.

Experiment No.	Condition of addition of crystals				Result of culture		
	Crystal form	Added weight (g)	Added concentration (g/L)	Glu accumulation at addition (g/L)	Crystal form of precipitated crystals	Total Glu accumulation (g/L)	Of all, precipitated crystals (g/L)
1	α -crystal	1.000	2.778	44.5	α -crystal	145.0	98.0
2	α -crystal	1.000	2.778	47.0	α -crystal	118.0	72.0
3	α -crystal	1.000	2.778	46.5	α -crystal	132.0	73.0
4	α -crystal	1.000	2.778	47.0	α -crystal	131.0	86.0
5	α -crystal	1.000	2.778	46.0	α -crystal	104.0	58.0
6	α -crystal	1.000	2.778	45.5	α -crystal	142.0	90.0
7	α -crystal	0.400	1.111	41.5	α -crystal	102.0	60.0
8	α -crystal	0.400	1.111	30.5	α -crystal	92.0	50.0
9	α -crystal	0.400	1.111	55.5	α -crystal	94.0	50.0
10	α -crystal	0.400	1.111	59.5	α -crystal	93.0	56.0
11	α -crystal	0.400	1.111	61.0	α -crystal	110.0	67.0
12	α -crystal	0.080	0.222	44.5	α -crystal	118.0	77.5
13	α -crystal	0.080	0.222	46.5	α -crystal	132.0	91.0
14	α -crystal	0.080	0.222	46.0	α -crystal	104.0	63.0
15	α -crystal	0.080	0.222	41.5	α -crystal	142.0	101.0
16	α -crystal	0.080	0.222	55.5	α -crystal	102.0	62.0
17	α -crystal	0.040	0.111	50.0	α -crystal	129.0	52.0
18	α -crystal	0.040	0.111	55.0	α -crystal	122.0	81.0
19	α -crystal	0.040	0.111	47.5	α -crystal	123.0	81.0
20	α -crystal	0.040	0.111	49.5	α -crystal	122.0	88.0
21	α -crystal	0.040	0.111	49.0	β -crystal	101.0	67.0
22	α -crystal	0.040	0.111	48.0	β -crystal	100.0	65.0
23	α -crystal	0.008	0.022	52.5	β -crystal	113.0	72.0
24	α -crystal	0.008	0.022	54.0	β -crystal	91.0	50.0
25	β -crystal	1.000	2.778	44.5	β -crystal	113.0	71.0
Control	None	—	—	—	β -crystal	82.0	39.5

From the results shown Table 1, it has been found that the accumulation of L-glutamic acid at the time of termination of the culture when crystals are added was increased compared with when no crystal is added at all.

Also, it has been found that if L-glutamic acid crystals of α -form are added, the proportion of precipitation of

L-glutamic acid crystals of α -form increases. In particular, it has been found that if the crystals are added in an amount of 0.2 g/L or more, L-glutamic acid crystals of α -form can be precipitated in the medium with good reproducibility.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 8

<210> SEQ ID NO 1

<211> LENGTH: 935

<212> TYPE: PRT

<213> ORGANISM: *Pantoea ananatis*

<400> SEQUENCE: 1

Met Gln Asn Ser Ala Met Lys Pro Trp Leu Asp Ser Ser Trp Leu Ala
1 5 10 15

Gly Ala Asn Gln Ser Tyr Ile Glu Gln Leu Tyr Glu Asp Phe Leu Thr
20 25 30

Asp Pro Asp Ser Val Asp Ala Val Trp Arg Ser Met Phe Gln Gln Leu
35 40 45

Pro Gly Thr Gly Val Lys Pro Glu Gln Phe His Ser Ala Thr Arg Glu
50 55 60

Tyr Phe Arg Arg Leu Ala Lys Asp Ala Ser Arg Tyr Thr Ser Ser Val
65 70 75 80

Thr Asp Pro Ala Thr Asn Ser Lys Gln Val Lys Val Leu Gln Leu Ile
85 90 95

Asn Ala Phe Arg Phe Arg Gly His Gln Glu Ala Asn Leu Asp Pro Leu
100 105 110

-continued

Gly	Leu	Trp	Lys	Gln	Asp	Arg	Val	Ala	Asp	Leu	Asp	Pro	Ala	Phe	His
		115					120					125			
Asp	Leu	Thr	Asp	Ala	Asp	Phe	Gln	Glu	Ser	Phe	Asn	Val	Gly	Ser	Phe
	130					135					140				
Ala	Ile	Gly	Lys	Glu	Thr	Met	Lys	Leu	Ala	Asp	Leu	Phe	Asp	Ala	Leu
145					150					155					160
Lys	Gln	Thr	Tyr	Cys	Gly	Ser	Ile	Gly	Ala	Glu	Tyr	Met	His	Ile	Asn
				165					170					175	
Asn	Thr	Glu	Glu	Lys	Arg	Trp	Ile	Gln	Gln	Arg	Ile	Glu	Ser	Gly	Ala
			180					185					190		
Ser	Gln	Thr	Ser	Phe	Ser	Gly	Glu	Glu	Lys	Lys	Gly	Phe	Leu	Lys	Glu
		195					200					205			
Leu	Thr	Ala	Ala	Glu	Gly	Leu	Glu	Lys	Tyr	Leu	Gly	Ala	Lys	Phe	Pro
	210					215					220				
Gly	Ala	Lys	Arg	Phe	Ser	Leu	Glu	Gly	Gly	Asp	Ala	Leu	Val	Pro	Met
225					230					235					240
Leu	Arg	Glu	Met	Ile	Arg	His	Ala	Gly	Lys	Ser	Gly	Thr	Arg	Glu	Val
			245						250					255	
Val	Leu	Gly	Met	Ala	His	Arg	Gly	Arg	Leu	Asn	Val	Leu	Ile	Asn	Val
			260					265					270		
Leu	Gly	Lys	Lys	Pro	Gln	Asp	Leu	Phe	Asp	Glu	Phe	Ser	Gly	Lys	His
		275					280					285			
Lys	Glu	His	Leu	Gly	Thr	Gly	Asp	Val	Lys	Tyr	His	Met	Gly	Phe	Ser
	290					295					300				
Ser	Asp	Ile	Glu	Thr	Glu	Gly	Gly	Leu	Val	His	Leu	Ala	Leu	Ala	Phe
305					310					315					320
Asn	Pro	Ser	His	Leu	Glu	Ile	Val	Ser	Pro	Val	Val	Met	Gly	Ser	Val
				325					330					335	
Arg	Ala	Arg	Leu	Asp	Arg	Leu	Ala	Glu	Pro	Val	Ser	Asn	Lys	Val	Leu
			340					345					350		
Pro	Ile	Thr	Ile	His	Gly	Asp	Ala	Ala	Val	Ile	Gly	Gln	Gly	Val	Val
		355					360					365			
Gln	Glu	Thr	Leu	Asn	Met	Ser	Gln	Ala	Arg	Gly	Tyr	Glu	Val	Gly	Gly
	370					375					380				
Thr	Val	Arg	Ile	Val	Ile	Asn	Asn	Gln	Val	Gly	Phe	Thr	Thr	Ser	Asn
385					390					395					400
Pro	Lys	Asp	Ala	Arg	Ser	Thr	Pro	Tyr	Cys	Thr	Asp	Ile	Gly	Lys	Met
				405					410					415	
Val	Leu	Ala	Pro	Ile	Phe	His	Val	Asn	Ala	Asp	Asp	Pro	Glu	Ala	Val
			420					425					430		
Ala	Phe	Val	Thr	Arg	Leu	Ala	Leu	Asp	Tyr	Arg	Asn	Thr	Phe	Lys	Arg
		435					440					445			
Asp	Val	Phe	Ile	Asp	Leu	Val	Cys	Tyr	Arg	Arg	His	Gly	His	Asn	Glu
	450					455					460				
Ala	Asp	Glu	Pro	Ser	Ala	Thr	Gln	Pro	Leu	Met	Tyr	Gln	Lys	Ile	Lys
465					470					475					480
Lys	His	Pro	Thr	Pro	Arg	Lys	Ile	Tyr	Ala	Asp	Arg	Leu	Glu	Gly	Glu
				485					490					495	
Gly	Val	Ala	Ser	Gln	Glu	Asp	Ala	Thr	Glu	Met	Val	Asn	Leu	Tyr	Arg
			500					505					510		
Asp	Ala	Leu	Asp	Ala	Gly	Glu	Cys	Val	Val	Pro	Glu	Trp	Arg	Pro	Met
		515					520					525			
Ser	Leu	His	Ser	Phe	Thr	Trp	Ser	Pro	Tyr	Leu	Asn	His	Glu	Trp	Asp

-continued

530			535			540										
Glu 545	Pro	Tyr	Pro	Ala	Gln	Val	Asp	Met	Lys	Arg	Leu	Lys	Glu	Leu	Ala	560
Leu	Arg	Ile	Ser	Gln	Val	Pro	Glu	Gln	Ile	Glu	Val	Gln	Ser	Arg	Val	575
Ala	Lys	Ile	Tyr	Asn	Asp	Arg	Lys	Leu	Met	Ala	Glu	Gly	Glu	Lys	Ala	590
Phe	Asp	Trp	Gly	Gly	Ala	Glu	Asn	Leu	Ala	Tyr	Ala	Thr	Leu	Val	Asp	605
Glu	Gly	Ile	Pro	Val	Arg	Leu	Ser	Gly	Glu	Asp	Ser	Gly	Arg	Gly	Thr	620
Phe	Phe	His	Arg	His	Ala	Val	Val	His	Asn	Gln	Ala	Asn	Gly	Ser	Thr	640
Tyr	Thr	Pro	Leu	His	His	Ile	His	Asn	Ser	Gln	Gly	Glu	Phe	Lys	Val	655
Trp	Asp	Ser	Val	Leu	Ser	Glu	Glu	Ala	Val	Leu	Ala	Phe	Glu	Tyr	Gly	670
Tyr	Ala	Thr	Ala	Glu	Pro	Arg	Val	Leu	Thr	Ile	Trp	Glu	Ala	Gln	Phe	685
Gly	Asp	Phe	Ala	Asn	Gly	Ala	Gln	Val	Val	Ile	Asp	Gln	Phe	Ile	Ser	700
Ser	Gly	Glu	Gln	Lys	Trp	Gly	Arg	Met	Cys	Gly	Leu	Val	Met	Leu	Leu	720
Pro	His	Gly	Tyr	Glu	Gly	Gln	Gly	Pro	Glu	His	Ser	Ser	Ala	Arg	Leu	735
Glu	Arg	Tyr	Leu	Gln	Leu	Cys	Ala	Glu	Gln	Asn	Met	Gln	Val	Cys	Val	750
Pro	Ser	Thr	Pro	Ala	Gln	Val	Tyr	His	Met	Leu	Arg	Arg	Gln	Ala	Leu	765
Arg	Gly	Met	Arg	Arg	Pro	Leu	Val	Val	Met	Ser	Pro	Lys	Ser	Leu	Leu	780
Arg	His	Pro	Leu	Ala	Ile	Ser	Ser	Leu	Asp	Glu	Leu	Ala	Asn	Gly	Ser	800
Phe	Gln	Pro	Ala	Ile	Gly	Glu	Ile	Asp	Asp	Leu	Asp	Pro	Gln	Gly	Val	815
Lys	Arg	Val	Val	Leu	Cys	Ser	Gly	Lys	Val	Tyr	Tyr	Asp	Leu	Leu	Glu	830
Gln	Arg	Arg	Lys	Asp	Glu	Lys	Thr	Asp	Val	Ala	Ile	Val	Arg	Ile	Glu	845
Gln	Leu	Tyr	Pro	Phe	Pro	His	Gln	Ala	Val	Gln	Glu	Ala	Leu	Lys	Ala	860
Tyr	Ser	His	Val	Gln	Asp	Phe	Val	Trp	Cys	Gln	Glu	Glu	Pro	Leu	Asn	880
Gln	Gly	Ala	Trp	Tyr	Cys	Ser	Gln	His	His	Phe	Arg	Asp	Val	Val	Pro	895
Phe	Gly	Ala	Thr	Leu	Arg	Tyr	Ala	Gly	Arg	Pro	Ala	Ser	Ala	Ser	Pro	910
Ala	Val	Gly	Tyr	Met	Ser	Val	His	Gln	Gln	Gln	Gln	Gln	Asp	Leu	Val	925
Asn	Asp	Ala	Leu	Asn	Val	Asn										935

-continued

```

<211> LENGTH: 407
<212> TYPE: PRT
<213> ORGANISM: Pantoea ananatis

<400> SEQUENCE: 2

Met Ser Ser Val Asp Ile Leu Val Pro Asp Leu Pro Glu Ser Val Ala
1          5          10          15
Asp Ala Thr Val Ala Thr Trp His Lys Lys Pro Gly Asp Ala Val Ser
20          25          30
Arg Asp Glu Val Ile Val Glu Ile Glu Thr Asp Lys Val Val Leu Glu
35          40          45
Val Pro Ala Ser Ala Asp Gly Val Leu Glu Ala Val Leu Glu Asp Glu
50          55          60
Gly Ala Thr Val Thr Ser Arg Gln Ile Leu Gly Arg Leu Lys Glu Gly
65          70          75          80
Asn Ser Ala Gly Lys Glu Ser Ser Ala Lys Ala Glu Ser Asn Asp Thr
85          90          95
Thr Pro Ala Gln Arg Gln Thr Ala Ser Leu Glu Glu Glu Ser Ser Asp
100         105         110
Ala Leu Ser Pro Ala Ile Arg Arg Leu Ile Ala Glu His Asn Leu Asp
115         120         125
Ala Ala Gln Ile Lys Gly Thr Gly Val Gly Gly Arg Leu Thr Arg Glu
130         135         140
Asp Val Glu Lys His Leu Ala Asn Lys Pro Gln Ala Glu Lys Ala Ala
145         150         155         160
Ala Pro Ala Ala Gly Ala Ala Thr Ala Gln Gln Pro Val Ala Asn Arg
165         170         175
Ser Glu Lys Arg Val Pro Met Thr Arg Leu Arg Lys Arg Val Ala Glu
180         185         190
Arg Leu Leu Glu Ala Lys Asn Ser Thr Ala Met Leu Thr Thr Phe Asn
195         200         205
Glu Ile Asn Met Lys Pro Ile Met Asp Leu Arg Lys Gln Tyr Gly Asp
210         215         220
Ala Phe Glu Lys Arg His Gly Val Arg Leu Gly Phe Met Ser Phe Tyr
225         230         235         240
Ile Lys Ala Val Val Glu Ala Leu Lys Arg Tyr Pro Glu Val Asn Ala
245         250         255
Ser Ile Asp Gly Glu Asp Val Val Tyr His Asn Tyr Phe Asp Val Ser
260         265         270
Ile Ala Val Ser Thr Pro Arg Gly Leu Val Thr Pro Val Leu Arg Asp
275         280         285
Val Asp Ala Leu Ser Met Ala Asp Ile Glu Lys Lys Ile Lys Glu Leu
290         295         300
Ala Val Lys Gly Arg Asp Gly Lys Leu Thr Val Asp Asp Leu Thr Gly
305         310         315         320
Gly Asn Phe Thr Ile Thr Asn Gly Gly Val Phe Gly Ser Leu Met Ser
325         330         335
Thr Pro Ile Ile Asn Pro Pro Gln Ser Ala Ile Leu Gly Met His Ala
340         345         350
Ile Lys Asp Arg Pro Met Ala Val Asn Gly Gln Val Val Ile Leu Pro
355         360         365
Met Met Tyr Leu Ala Leu Ser Tyr Asp His Arg Leu Ile Asp Gly Arg
370         375         380
Glu Ser Val Gly Tyr Leu Val Ala Val Lys Glu Met Leu Glu Asp Pro

```

-continued

385 390 395 400
 Ala Arg Leu Leu Leu Asp Val
 405

 <210> SEQ ID NO 3
 <211> LENGTH: 41
 <212> TYPE: PRT
 <213> ORGANISM: *Pantoea ananatis*

 <400> SEQUENCE: 3

 Met Asn Leu His Glu Tyr Gln Ala Lys Gln Leu Phe Ala Arg Tyr Gly
 1 5 10 15
 Met Pro Ala Pro Thr Gly Tyr Ala Cys Thr Thr Pro Arg Glu Ala Glu
 20 25 30
 Glu Ala Ala Ser Lys Ile Gly Ala Gly
 35 40

 <210> SEQ ID NO 4
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: *Pantoea ananatis*

 <400> SEQUENCE: 4

 Ala Phe Ser Val Phe Arg Cys His Ser Ile Met Asn Cys Val Ser Val
 1 5 10 15
 Cys Pro Lys Gly Leu Asn Pro Thr Arg Ala Ile Gly His Ile Lys Ser
 20 25 30
 Met Leu Leu Gln Arg Ser Ala
 35

 <210> SEQ ID NO 5
 <211> LENGTH: 933
 <212> TYPE: PRT
 <213> ORGANISM: *Escherichia coli*

 <400> SEQUENCE: 5

 Met Gln Asn Ser Ala Leu Lys Ala Trp Leu Asp Ser Ser Tyr Leu Ser
 1 5 10 15
 Gly Ala Asn Gln Ser Trp Ile Glu Gln Leu Tyr Glu Asp Phe Leu Thr
 20 25 30
 Asp Pro Asp Ser Val Asp Ala Asn Trp Arg Ser Thr Phe Gln Gln Leu
 35 40 45
 Pro Gly Thr Gly Val Lys Pro Asp Gln Phe His Ser Gln Thr Arg Glu
 50 55 60
 Tyr Phe Arg Arg Leu Ala Lys Asp Ala Ser Arg Tyr Ser Ser Thr Ile
 65 70 75 80
 Ser Asp Pro Asp Thr Asn Val Lys Gln Val Lys Val Leu Gln Leu Ile
 85 90 95
 Asn Ala Tyr Arg Phe Arg Gly His Gln His Ala Asn Leu Asp Pro Leu
 100 105 110
 Gly Leu Trp Gln Gln Asp Lys Val Ala Asp Leu Asp Pro Ser Phe His
 115 120 125
 Asp Leu Thr Glu Ala Asp Phe Gln Glu Thr Phe Asn Val Gly Ser Phe
 130 135 140
 Ala Ser Gly Lys Glu Thr Met Lys Leu Gly Glu Leu Leu Glu Ala Leu
 145 150 155 160
 Lys Gln Thr Tyr Cys Gly Pro Ile Gly Ala Glu Tyr Met His Ile Thr
 165 170 175

-continued

Ser Thr Glu Glu Lys Arg Trp Ile Gln Gln Arg Ile Glu Ser Gly Arg
 180 185 190
 Ala Thr Phe Asn Ser Glu Glu Lys Lys Arg Phe Leu Ser Glu Leu Thr
 195 200 205
 Ala Ala Glu Gly Leu Glu Arg Tyr Leu Gly Ala Lys Phe Pro Gly Ala
 210 215 220
 Lys Arg Phe Ser Leu Glu Gly Gly Asp Ala Leu Ile Pro Met Leu Lys
 225 230 235 240
 Glu Met Ile Arg His Ala Gly Asn Ser Gly Thr Arg Glu Val Val Leu
 245 250 255
 Gly Met Ala His Arg Gly Arg Leu Asn Val Leu Val Asn Val Leu Gly
 260 265 270
 Lys Lys Pro Gln Asp Leu Phe Asp Glu Phe Ala Gly Lys His Lys Glu
 275 280 285
 His Leu Gly Thr Gly Asp Val Lys Tyr His Met Gly Phe Ser Ser Asp
 290 295 300
 Phe Gln Thr Asp Gly Gly Leu Val His Leu Ala Leu Ala Phe Asn Pro
 305 310 315 320
 Ser His Leu Glu Ile Val Ser Pro Val Val Ile Gly Ser Val Arg Ala
 325 330 335
 Arg Leu Asp Arg Leu Asp Glu Pro Ser Ser Asn Lys Val Leu Pro Ile
 340 345 350
 Thr Ile His Gly Asp Ala Ala Val Thr Gly Gln Gly Val Val Gln Glu
 355 360 365
 Thr Leu Asn Met Ser Lys Ala Arg Gly Tyr Glu Val Gly Gly Thr Val
 370 375 380
 Arg Ile Val Ile Asn Asn Gln Val Gly Phe Thr Thr Ser Asn Pro Leu
 385 390 395 400
 Asp Ala Arg Ser Thr Pro Tyr Cys Thr Asp Ile Gly Lys Met Val Gln
 405 410 415
 Ala Pro Ile Phe His Val Asn Ala Asp Asp Pro Glu Ala Val Ala Phe
 420 425 430
 Val Thr Arg Leu Ala Leu Asp Phe Arg Asn Thr Phe Lys Arg Asp Val
 435 440 445
 Phe Ile Asp Leu Val Ser Tyr Arg Arg His Gly His Asn Glu Ala Asp
 450 455 460
 Glu Pro Ser Ala Thr Gln Pro Leu Met Tyr Gln Lys Ile Lys Lys His
 465 470 475 480
 Pro Thr Pro Arg Lys Ile Tyr Ala Asp Lys Leu Glu Gln Glu Lys Val
 485 490 495
 Ala Thr Leu Glu Asp Ala Thr Glu Met Val Asn Leu Tyr Arg Asp Ala
 500 505 510
 Leu Asp Ala Gly Asp Cys Val Val Ala Glu Trp Arg Pro Met Asn Met
 515 520 525
 His Ser Phe Thr Trp Ser Pro Tyr Leu Asn His Glu Trp Asp Glu Glu
 530 535 540
 Tyr Pro Asn Lys Val Glu Met Lys Arg Leu Gln Glu Leu Ala Lys Arg
 545 550 555 560
 Ile Ser Thr Val Pro Glu Ala Val Glu Met Gln Ser Arg Val Ala Lys
 565 570 575
 Ile Tyr Gly Asp Arg Gln Ala Met Ala Ala Gly Glu Lys Leu Phe Asp
 580 585 590

-continued

Trp Gly Gly Ala Glu Asn Leu Ala Tyr Ala Thr Leu Val Asp Glu Gly
 595 600 605
 Ile Pro Val Arg Leu Ser Gly Glu Asp Ser Gly Arg Gly Thr Phe Phe
 610 615 620
 His Arg His Ala Val Ile His Asn Gln Ser Asn Gly Ser Thr Tyr Thr
 625 630 635 640
 Pro Leu Gln His Ile His Asn Gly Gln Gly Ala Phe Arg Val Trp Asp
 645 650 655
 Ser Val Leu Ser Glu Glu Ala Val Leu Ala Phe Glu Tyr Gly Tyr Ala
 660 665 670
 Thr Ala Glu Pro Arg Thr Leu Thr Ile Trp Glu Ala Gln Phe Gly Asp
 675 680 685
 Phe Ala Asn Gly Ala Gln Val Val Ile Asp Gln Phe Ile Ser Ser Gly
 690 695 700
 Glu Gln Lys Trp Gly Arg Met Cys Gly Leu Val Met Leu Leu Pro His
 705 710 715 720
 Gly Tyr Glu Gly Gln Gly Pro Glu His Ser Ser Ala Arg Leu Glu Arg
 725 730 735
 Tyr Leu Gln Leu Cys Ala Glu Gln Asn Met Gln Val Cys Val Pro Ser
 740 745 750
 Thr Pro Ala Gln Val Tyr His Met Leu Arg Arg Gln Ala Leu Arg Gly
 755 760 765
 Met Arg Arg Pro Leu Val Val Met Ser Pro Lys Ser Leu Leu Arg His
 770 775 780
 Pro Leu Ala Val Ser Ser Leu Glu Glu Leu Ala Asn Gly Thr Phe Leu
 785 790 795 800
 Pro Ala Ile Gly Glu Ile Asp Glu Leu Asp Pro Lys Gly Val Lys Arg
 805 810 815
 Val Val Met Cys Ser Gly Lys Val Tyr Tyr Asp Leu Leu Glu Gln Arg
 820 825 830
 Arg Lys Asn Asn Gln His Asp Val Ala Ile Val Arg Ile Glu Gln Leu
 835 840 845
 Tyr Pro Phe Pro His Lys Ala Met Gln Glu Val Leu Gln Gln Phe Ala
 850 855 860
 His Val Lys Asp Phe Val Trp Cys Gln Glu Glu Pro Leu Asn Gln Gly
 865 870 875 880
 Ala Trp Tyr Cys Ser Gln His His Phe Arg Glu Val Ile Pro Phe Gly
 885 890 895
 Ala Ser Leu Arg Tyr Ala Gly Arg Pro Ala Ser Ala Ser Pro Ala Val
 900 905 910
 Gly Tyr Met Ser Val His Gln Lys Gln Gln Gln Asp Leu Val Asn Asp
 915 920 925
 Ala Leu Asn Val Glu
 930

<210> SEQ ID NO 6
 <211> LENGTH: 405
 <212> TYPE: PRT
 <213> ORGANISM: Escherichia coli

<400> SEQUENCE: 6

Met Ser Ser Val Asp Ile Leu Val Pro Asp Leu Pro Glu Ser Val Ala
 1 5 10 15
 Asp Ala Thr Val Ala Thr Trp His Lys Lys Pro Gly Asp Ala Val Val
 20 25 30

-continued

Arg Asp Glu Val Leu Val Glu Ile Glu Thr Asp Lys Val Val Leu Glu
 35 40 45
 Val Pro Ala Ser Ala Asp Gly Ile Leu Asp Ala Val Leu Glu Asp Glu
 50 55 60
 Gly Thr Thr Val Thr Ser Arg Gln Ile Leu Gly Arg Leu Arg Glu Gly
 65 70 75 80
 Asn Ser Ala Gly Lys Glu Thr Ser Ala Lys Ser Glu Glu Lys Ala Ser
 85 90 95
 Thr Pro Ala Gln Arg Gln Gln Ala Ser Leu Glu Glu Gln Asn Asn Asp
 100 105 110
 Ala Leu Ser Pro Ala Ile Arg Arg Leu Leu Ala Glu His Asn Leu Asp
 115 120 125
 Ala Ser Ala Ile Lys Gly Thr Gly Val Gly Gly Arg Leu Thr Arg Glu
 130 135 140
 Asp Val Glu Lys His Leu Ala Lys Ala Pro Ala Lys Glu Ser Ala Pro
 145 150 155 160
 Ala Ala Ala Ala Pro Ala Ala Gln Pro Ala Leu Ala Ala Arg Ser Glu
 165 170 175
 Lys Arg Val Pro Met Thr Arg Leu Arg Lys Arg Val Ala Glu Arg Leu
 180 185 190
 Leu Glu Ala Lys Asn Ser Thr Ala Met Leu Thr Thr Phe Asn Glu Val
 195 200 205
 Asn Met Lys Pro Ile Met Asp Leu Arg Lys Gln Tyr Gly Glu Ala Phe
 210 215 220
 Glu Lys Arg His Gly Ile Arg Leu Gly Phe Met Ser Phe Tyr Val Lys
 225 230 235 240
 Ala Val Val Glu Ala Leu Lys Arg Tyr Pro Glu Val Asn Ala Ser Ile
 245 250 255
 Asp Gly Asp Asp Val Val Tyr His Asn Tyr Phe Asp Val Ser Met Ala
 260 265 270
 Val Ser Thr Pro Arg Gly Leu Val Thr Pro Val Leu Arg Asp Val Asp
 275 280 285
 Thr Leu Gly Met Ala Asp Ile Glu Lys Lys Ile Lys Glu Leu Ala Val
 290 295 300
 Lys Gly Arg Asp Gly Lys Leu Thr Val Glu Asp Leu Thr Gly Gly Asn
 305 310 315 320
 Phe Thr Ile Thr Asn Gly Gly Val Phe Gly Ser Leu Met Ser Thr Pro
 325 330 335
 Ile Ile Asn Pro Pro Gln Ser Ala Ile Leu Gly Met His Ala Ile Lys
 340 345 350
 Asp Arg Pro Met Ala Val Asn Gly Gln Val Glu Ile Leu Pro Met Met
 355 360 365
 Tyr Leu Ala Leu Ser Tyr Asp His Arg Leu Ile Asp Gly Arg Glu Ser
 370 375 380
 Val Gly Phe Leu Val Thr Ile Lys Glu Leu Leu Glu Asp Pro Thr Arg
 385 390 395 400
 Leu Leu Leu Asp Val
 405

<210> SEQ ID NO 7

<211> LENGTH: 60

<212> TYPE: PRT

<213> ORGANISM: Escherichia coli

-continued

<400> SEQUENCE: 7

```

Met Asn Leu His Glu Tyr Gln Ala Lys Gln Leu Phe Ala Arg Tyr Gly
1           5           10           15
Leu Pro Ala Pro Val Gly Tyr Ala Cys Thr Thr Pro Arg Glu Ala Glu
          20           25           30
Glu Ala Ala Ser Lys Ile Gly Ala Gly Pro Trp Val Val Lys Cys Gln
          35           40           45
Val His Ala Gly Gly Arg Gly Lys Ala Gly Gly Val
          50           55           60

```

<210> SEQ ID NO 8

<211> LENGTH: 58

<212> TYPE: PRT

<213> ORGANISM: Escherichia coli

<400> SEQUENCE: 8

```

Phe Leu Ile Asp Ser Arg Asp Thr Glu Thr Asp Ser Arg Leu Asp Gly
1           5           10           15
Leu Ser Asp Ala Phe Ser Val Phe Arg Cys His Ser Ile Met Asn Cys
          20           25           30
Val Ser Val Cys Pro Lys Gly Leu Asn Pro Thr Arg Ala Ile Gly His
          35           40           45
Ile Lys Ser Met Leu Leu Gln Arg Asn Ala
          50           55

```

What is claimed is:

1. A method for producing L-glutamic acid by fermentation, which comprises culturing a microorganism having L-glutamic acid-producing ability in a liquid medium of which pH is adjusted within the range of 3 to 5 to allow L-glutamic acid to be produced and accumulated with precipitation of L-glutamic acid accompanied, wherein an operation causing existence of L-glutamic acid crystals in the medium is performed after the fermentation begins and when a concentration of L-glutamic acid in the medium is lower than the concentration at which spontaneous crystallization occurs,

wherein the operation causing existence of L-glutamic acid crystals in the medium is addition of L-glutamic acid crystals, and

wherein L-glutamic acid crystals are not added to the medium before the fermentation.

2. The method according to claim 1, wherein the crystals are α -crystals.

3. The method according to claim 2, wherein an amount of the crystals added to the medium is 0.2 g/L or more.

4. The method according to claim 1, wherein the microorganism belongs to the genus *[Enterobacter] Pantoea*.

5. The method according to claim 2, wherein the microorganism belongs to the genus *[Enterobacter] Pantoea*.

6. The method according to claim 3, wherein the microorganism belongs to the genus *[Enterobacter] Pantoea*.

7. The method according to claim 4, wherein the microorganism is *[Enterobacter agglomerans] Pantoea ananatis*.

8. The method according to claim 5, wherein the microorganism is *[Enterobacter agglomerans] Pantoea ananatis*.

9. The method according to claim 6, wherein the microorganism is *[Enterobacter agglomerans] Pantoea ananatis*.

10. The method according to claim 1, wherein the microorganism can metabolize a carbon source in a liquid medium contacting L-glutamic acid at a saturation concentration and the carbon source, the liquid medium having a pH within the

range of 3 to 5 and has an ability to accumulate L glutamic acid in an amount exceeding the saturation concentration of L-glutamic acid in the liquid medium at the pH.

11. The method according to claim 2, wherein the microorganism can metabolize a carbon source in a liquid medium containing L-glutamic acid at a saturation concentration and the carbon source, the liquid medium having a pH within the range of 3 to 5 and has an ability to accumulate L-glutamic acid in an amount exceeding the saturation concentration of L-glutamic acid in the liquid medium at the pH.

12. The method according to claim 3, wherein the microorganism can metabolize a carbon source in a liquid medium containing L-glutamic acid at a saturation concentration and the carbon source, the liquid medium having a pH within the range of 3 to 5 and has an ability to accumulate L glutamic acid in an amount exceeding the saturation concentration of L-glutamic acid in the liquid medium at the pH.

13. The method according to claim 4, wherein the microorganism can metabolize a carbon source in a liquid medium containing L-glutamic acid at a saturation concentration and the carbon source, the liquid medium having a pH within the range of 3 to 5 and has an ability to accumulate L-glutamic acid in an amount exceeding the saturation concentration of L-glutamic acid in the liquid medium at the pH.

14. The method according to claim 5, wherein the microorganism can metabolize a carbon source in a liquid medium containing L-glutamic acid at a saturation concentration and the carbon source, the liquid medium having a pH within the range of 3 to 5 and has an ability to accumulate L-glutamic acid in an amount exceeding the saturation concentration of L-glutamic acid in the liquid medium at the pH.

15. The method according to claim 7, wherein said microorganism is *Pantoea ananatis* AJ13355 strain.

16. The method according to claim 8, wherein said microorganism is *Pantoea ananatis* AJ13355 strain.

17. The method according to claim 9, wherein said microorganism is *Pantoea ananatis* AJ13355 strain.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : RE41,800 E
APPLICATION NO. : 12/485550
DATED : October 5, 2010
INVENTOR(S) : Hiroshi Ueda et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the title page, Item (30), the Foreign Application Priority Data has been omitted.
Item (30) should read:

-- (30) **Foreign Application Priority Data**

Feb. 20, 2001 (JP)2001-044136 --

Signed and Sealed this
Sixteenth Day of August, 2011



David J. Kappos
Director of the United States Patent and Trademark Office