



US00RE44266E

(19) **United States**
 (12) **Reissued Patent**
 Mayfield

(10) **Patent Number:** **US RE44,266 E**
 (45) **Date of Reissued Patent:** ***Jun. 4, 2013**

(54) **EXPRESSION OF EUKARYOTIC
 POLYPEPTIDES IN CHLOROPLASTS**

(75) Inventor: **Stephen P. Mayfield**, Cardiff, CA (US)

(73) Assignee: **The Scripps Research Institute**, La Jolla, CA (US)

(*) Notice: This patent is subject to a terminal disclaimer.

(21) Appl. No.: **11/197,730**

(22) Filed: **Aug. 3, 2005**

Related U.S. Patent Documents

Reissue of:

(64) Patent No.: **6,156,517**
 Issued: **Dec. 5, 2000**
 Appl. No.: **09/341,550**
 PCT Filed: **Jan. 16, 1998**
 PCT No.: **PCT/US98/00840**
 § 371 (c)(1),
 (2), (4) Date: **Jul. 13, 1999**
 PCT Pub. No.: **WO98/31823**
 PCT Pub. Date: **Jul. 23, 1998**

U.S. Applications:

(63) Continuation of application No. 10/310,587, filed on Dec. 4, 2002, now Pat. No. Re. 39,350.

(60) Provisional application No. 60/035,955, filed on Jan. 17, 1997, provisional application No. 60/069,400, filed on Dec. 12, 1997.

(51) **Int. Cl.**
C12P 21/02 (2006.01)

(52) **U.S. Cl.**
 USPC **435/69.1; 435/320.1; 435/419; 435/375;**
435/468; 536/23.1

(58) **Field of Classification Search**
 None
 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,451,513	A *	9/1995	Maliga et al.	800/278
5,661,017	A	8/1997	Dunahay et al.	
5,693,507	A *	12/1997	Daniell et al.	435/470
6,156,517	A	12/2000	Mayfield	435/6
6,271,444	B1	8/2001	McBride et al.	800/300

FOREIGN PATENT DOCUMENTS

WO	WO 97/41228	A2	11/1997
WO	WO 01/64929	A1	2/2001

OTHER PUBLICATIONS

Blowers et al., *The Plant Cell*, vol. 1,(1989) pp. 123-132.*
 Tavladoraki et al, *Nature*, vol. 366 (1993) pp. 469-472.*
 Sidorov et al., "Stable chloroplast transformation in potato: use of green fluorescent protein as a plastid marker." *The Plant Journal*, vol. 19 (2) pp. 209-216 (1999).
 Daniell et al., "Transient Foreign Gene Expression in Chloroplasts of Cultured Tobacco Cells After Biolistic Delivery of Chloroplast Vectors", *Proc. Natl. Acad. Sci.*, 87:88-92 (1990).

Danon and Mayfield, "Light Regulated Translational Activators: Identification of Chloroplast Gene Specific mRNA Binding Proteins" *The EMBO Journal*, 10(13):3993-4001 (1991).

Danon and Mayfield, "ADP-Dependent Phosphorylation Regulates RNA-Binding in vitro: Implications in Light-Modulated Translation", *The EMBO Journal*, 13(9):2227-2235 (1994).

Hauser et al., "Translational Regulation of Chloroplast Genes", *The Journal of Biological Chemistry*, 271(3):1486-1497 (1996).

Kim and Mayfield, "Protein Disulfide Isomerase as a Regulator of Chloroplast Translational Activation", *Science*, 278:1954-1957 (1997).

Kin-Ying et al., "Introduction and Expression of Foreign DNA in Isolated Spinach Chloroplasts by Electroporation" *The Plant Journal*, 10(4):737-743 (1996).

Le et al., "Triticum Aestivum Poly(A)-Binding Protein (wheatpab) mRNA, Complete Cds", *Database EBI (online)*, p. 1-2 (1996).

Pihlajaniemi et al., "Molecular Cloning of the β -Subunit of Human Prolyl 4-Hydroxylase. This Subunit and Protein Disulphide Isomerase are Products of the Same Gene" *The EMBO Journal*, 6(3):643-649 (1987).

Yohn et al., "A Poly(A) Binding Protein Functions in the Chloroplast as a Message-Specific Translation Factor", *Proc. Natl. Acad. Sci.*, 95:2238-2243 (1998).

Yohn et al., "Translation of the Chloroplast *psbA* mRNA Requires the Nuclear-Encoded Poly(A)-Binding Protein, RB47", *The Journal of Cell Biology*, 142:435-442 (1998).

Ohtani et al., "Location and nucleotide sequence of a tobacco chloroplast DNA segment capable of replication in yeast", *Mol. Gen. Genet.* 195:1-4, 1984.

Rochaix et al., "Construction and Characterization of Autonomously Replicating Plasmids in the Green Unicellular *Alga Chlamydomonas reinhardtii*", *Cell*, 36:925-931, 1984.

Uchimiya et al., "Molecular Cloning of Tobacco Chromosomal and Chloroplast DNA Segments Capable of Replication in Yeast", *Mol. Gen. Genet.* 192:1-4, 1983.

Danon et al., *EMBO J.*, vol. 10, 1991, pp. 3993-4001.

Danon et al., *EMBO J.*, vol. 13, 1994, pp. 2227-2235.

* cited by examiner

Primary Examiner — Jim Ketter

(74) *Attorney, Agent, or Firm* — DLA Piper LLP (US)

(57) **ABSTRACT**

The present invention relates to a gene expression system in eukaryotic and prokaryotic cells, preferably plant cells and intact plants. In particular, the invention relates to an expression system having a RB47 binding site upstream of a translation initiation site for regulation of translation mediated by binding of RB47 protein, a member of the poly(A) binding protein family. Regulation is further effected by RB60, a protein disulfide isomerase. The expression system is capable of functioning in the nuclear/cytoplasm of cells and in the chloroplast of plants. Translation regulation of a desired molecule is enhanced approximately 100 fold over that obtained without RB47 binding site activation.

23 Claims, 17 Drawing Sheets

1 GAATTCGGCGCCGCTCCGTTGGTCCCTC ATG GTG TCT TTT TGA AGAGGACCTGAGCCTTTCACCCAAATATA 74
 1 M V S F * 5
 75 TCAAAAACCCGGCAACCGCCAAAATAATGCAAAAGCCCTCTCGTAGGCACAAAAGACCTATTCTAGCCATCAACTTT 154
 155 GTATCCGACGCTGCCGTTTAGCTGCGGCTTGAAGTCAAGC ATG GCG ACT ACT GAG TCC TCG GCC CCG 223
 1 M A T T E S S A P 9
 224 GCG GCC ACC ACC CAG CCG GCC AGC ACC CCG CTG GCG AAC TCG TCG CTG TAC GTC GGT GAC 283
 10 A A T T Q P A S T P L A N S S L Y V G D 29
 284 CTG GAG AAG GAT GTC ACC GAG GCC CAG CTG TTC GAG CTC TTC TCC TCG GTT GGC CCT GTG 343
 30 L E K D V T E A Q L F E L F S S V G P V 49
 344 GCC TCC ATT CGC GTG TGC CGC GAT GCC GTC ACG CGC TCG CTG GGC TAC GCC TAC GTC 403
 50 A S I R V C R D A V T R R S L G Y A Y V 69
 404 AAC TAC AAC AGC GCT CTG GAC CCC CAG GCT GCT GAC CGC GCC ATG GAG ACC CTG AAC TAC 463
 70 N Y N S A L D P Q A A D R A M E T L N Y 89
 464 CAT GTC GTG AAC GGC AAG CCT ATG CGC ATC ATG TGG TCG CAC CGC GAC CCT TCG GCC CGC 523
 90 H V V N G K P M R I M W S H R D P S A R 109
 524 AAG TCG GGC GTC AAC ATC TTC ATC AAG AAC CTG GAC AAG ACC ATC GAC GCC AAG GCC 583
 110 K S G V G N I F I K N L D K T I D A K A 129
 584 CTG CAC GAC ACC TTC TCG GCC TTC GGC AAG ATT CTG TCC TGC AAG GTT GCC ACT GAC GCC 643
 130 L H D T F S A F G K I L S C K V A T D A 149
 644 AAC GGC GTG TCG AAG GGC TAC GGC TTC GAG GAC CAG GCC GCT GCC GAT CGC 703
 150 N G V S K G Y G F V H F E D Q A A D R 169
 704 GCC ATT CAG ACC GTC AAC CAG AAG AAG ATT GAG GCC AAG ATC GTG TAC GTG GCC CCC TTC 763
 170 A I Q T V N Q K K I E G K I V Y V A P F 189

FIG. 1A

764 CAG AAG CGC GCT GAC CGC CCC AGG GCA AGG ACG TTG TAC ACC AAC GTG TTC GTC AAG AAC 823
 190 Q K R A D I G D E L G K M A T N V F V K N 209
 824 TTG CCG GCC GAC ATC GGC GAC GAG CTG GGC AAG ATG GCC ACC GAG CAC GGC GAG ATC 883
 210 L P A D I G D E L G K M A T N V F V K N 229
 884 ACC AGC GCG GTG GTC ATG AAG GAC GAC AAG GGC AGC AAG GGC TTC GGC TTC ATC AAC 943
 230 T S A V V M K D D K G G S K G F G F I N 249
 944 TTC AAG GAC GCC GAG TCG GCG GCC AAG TGC GTG GAG TAC CTG AAC GAG CGC GAG ATG AGC 1003
 250 F K D A E S A A K C V E Y L N E R E M S 269
 1004 GGC AAG ACC CTG TAC GCC GGC CGC CAG AAG AAG ACC GAG CGC GAG GCG ATG CTG CGC 1063
 270 G K T L Y A G R A Q K K T E R E A M L R 289
 1064 CAG AAG GCC GAG AGC AAG CAG GAG CGT TAC CTG AAG TAC CAG AGC ATG AAC CTG TAC 1123
 290 Q K A E S K Q E R Y L K Y Q S M N L Y 309
 1124 GTC AAG AAC CTG TCC GAC GAG GTC GAC GAC GGC CGT CTG GGC GAG CTG TTC GCC AAC 1183
 310 V K N L S D E V D D D A L R E L F A N 329
 1184 TCT GGC ACC ATC ACC TCG TGC AAG GTC ATG AAG GAC GGC AGC GGC AAG TCC AAG GGC TTC 1243
 330 S G T I T S C K V M K D G S G K S K G F 349
 1244 GGC TTC GTG TGC TTC ACC AGC CAC GAC GAG GCC ACC CCG CCC GTG ACC GAG ATG AAC 1303
 350 G F V C F T S H D E A T R P V T E M N 369
 1304 GGC AAG ATG GTC AAG GGC AAG CCC CTG TAC GTG GCC CTG GCG CAG CGC AAG GAC GTG CGC 1363
 370 G K M V K G K P L Y V A L A Q R K D V R 389
 1364 CGT GCC ACC CAG CTG GAG GCC AAC ATG CAG GCG CGC ATG GGC ATG GGC ATG AGC CGC 1423
 390 R A T Q L E A N M Q A R M G A M S R 409

FIG. 1B

1424 CCG CCG AAC CCG ATG GCC GGC ATG AGC CCC TAC CCC GGC GCC ATG CCG TTC TTC GCT CCC 1483
 410 P P N P M A G M S P Y P G A M P F F A P 429
 1484 GGC CCC GGC ATG GCT GCT GGC CCG CGC GGT GGC ATG TAC CCG CCC ATG ATG ATG 1543
 430 G P G G M A A G P R A P G M Y P P M M 449
 1544 CCG CCG GGC ATG CCT GGC CCC GGC CGC GGC CCC ATG ATG CCG CCC CAG 1603
 450 P P R G M P G P G R G P P M M P P Q 469
 1604 ATG ATG GGT GGC CCC ATG ATG GGC CCG CCC GGC CGT GGC GGC CGC CGC 1663
 470 M M G G P M M G P M G P G R G R G G R 489
 1664 GGC CCC TCC GGC CGC GGC CAG GGC AAC AAC GCC CCT GCC CAG CAG CCC AAG CCC 1723
 490 G P S G R G Q G R G N A P A Q Q P K P 509
 1724 GCC GCT GAG CCG GCC GGC CCC GCC GGC CCC GCT GCC GCG GCG CCT GCC GCC 1783
 510 A A E P A A A P A A A A A A A A A P A A 529
 1784 GCG GCG GAG CCG GAG GCC CCC GCC CAG CAG CCG CTG ACC GCC TCC GCG CTG GCC GCC 1843
 530 A A E P E A A P A A Q Q CAG CAG CCG CTG ACC GCC TCC GCG CTG GCC GCC 549
 1844 GCC GCG CCG GAG CAG AAG ATG ATG ATC GGC GAG CGC CTG TAC CCG CAG GTG GCG GAG 1903
 550 A A P E Q Q K M I G E R L Y P Q V A E 569
 1904 CTG CAG CCC GAC CTG GCT GGC AAG ATC ACC GCC ATG CTG GAG ATG GAC AAC GCC GAG 1963
 570 L Q P D L A G K I T G M L L E M D N A E 589
 1964 CTT CTG ATG CTT CTG GAG TCG CAC GAG GCG CTG GTG TCC AAG GTG GAC GAG GCC ATC GCT 2023
 590 L L M L L E S H E A L V S K V D E A I A 609
 2024 GTG CTC AAG CAG CAC AAC GTG ATT GCC GAG GAG AAC AAG GCT TAA AGCGCCTGCACGCTGTGCG 2088
 610 V L K Q H N V I A E E N K A * 624

FIG. 1C

2089 GGCTGGTGGCCGCGCCGCGCGCTGGCGCCGCGCAGC ATG GGC GCG GAC GCG GTG TGG TGG 2159
 1 M G A A D A V W 8
 2160 GAG CAG TGC TTG CTG CTT CTG GCC GTG AAG CCG CGC ACT GGG GCG GAC GCG AGG 2219
 9 E Q C L L L A A V K P R R T G A D G R 28
 2220 CTG GCG TTG ACG CCG GCG CAC AAC ACA AAG TTG GTG GCG TGA AAGTCTCTGGCGTCTCCG 2284
 29 L A L T P A R H N T K L V A * 43
 2285 GACGGTTGTAAGGTTTAAGAAGTGGCTTTTGGCCGGTTCGCCCCAAAGCGGAACGGCGTCTTTTCAGGCCAATCA 2364
 2365 CATCCGGCTGGAATAATTCITACCAAGCCACCCCTGCACCCAAATAATTCGGGTTCGGAAGAACAACCTCCCTTTT 2444
 2445 CCGGCAACGGTTCCTTTCAAGGCCAATCACTTTCCGGTGGGAAGAA ATG TTA CCC GGA AAA GGC GGG AAG 2516
 1 M L P G K G G K 8
 2517 CCC CCT GCA CCC GGA CAA GTT ATT CGG GGT TTC GCC GGG AAT GAG CAA GCG TTC GGG CTG 2576
 9 P A P G Q V I R G F A G N E Q A F G L 28
 2577 TTG GCC GTA TCG CGA ACG CTG TCG GGG TGT CAG GCC CCA GAA GGA AGG ATG ACG TTT TGG 2636
 29 L A V S R T L S G C Q A P E G R M T F W 48
 2637 TGA AGGGTGCAAACTGAGCACACGAGTTTTCGCAATAGACGTGGAGAAAGTCCAGTCCGGGTGAGCGGATAGCGGA 2715
 49 * 49
 2716 ATCAAGCGTGGCGGTCCCTGGCGAGACCGCTTCTGTGTTTGTGTTGCTGAGCCCTTTG ATG GCA CAA TCG CAC 2790
 1 M A Q S H 5
 2791 TGT TTT GAG CAG GCG ACT GTA AAG TGC CCG ACG CTA AAA AAG CCG CCG CGA ATT CC 2846
 6 C F E Q A T V K C P T L K K R P R I 23

FIG. 1D

MNRWNLALTLGLLLVAAPFTKHQFAHASDEYEDDEDDAPAAP
 KDDDDVDVTVTKNWDVTVKSKSFALVEFYAPWCGHCKTLKPEYAKAATALKAAAPDA
 LIAKVDAQTQEEslaQKFGVQGYPTLKWFDVJGELASDYNQPRDADGIVGWVKKKTGPPA
 VTVEDADKLSLEADAEVVVVGYFKALEGEIYDTFKSYAAKTEDVVFVQTTSADVAKA
 AGLDAVDTVSVVKNFAGEDRATAVLAIDLIDTDSLTAfVKSekMPPTIEFNQNSDKIF
 NSGINKQLILWTTADDLKADAEIMTVFREASKKFKGQLVFVTVNNEGDGADPVTNFFFG
 LKGATSPVLLGFFMEKNKKFRMEGEFTADNVAKFAESVVDGTAQAVLKSEAIPEDPYE
 DGVYKIVGKTVESVVLDETKDVLLEVYAPWCGHCKKLEPIYKKLAKRfKkVDSVIIAK
 MDGTENEHPEIEVKGFPPTILFYPAGSDRTPIVFEGGDRSLKSLTKfIKTNAKIPYELP
 KKGSDGDEGTSDDKPKASDKDEL

1 gagtacgttt acgccatgaa ccgttggaac cttcttgccc ttaccctggg sctgctgctg
 61 gtggcagcgc cttcaccaa gcaccagttt gctcatgctt ccgatgagta tgaggacgac
 121 gaggaggacg atgccccgc cgccttaag gacgacgacg tcgacgttac tgggtgacc
 181 gtcaagaact gggatgagac cgtcaagaag tccaagtctg cgttgtgga gttctacgct
 241 ccttggcgcg gccactgcaa gacctcaag cctgagtacg ctaaggctgc caccgccctg
 301 aaggctgctg ctccgatgc cttatcgcc aaggctgacg ccaccagga ggagtccttg
 361 gcccagaagt tcggcgtgca gggctaccc accctcaagt ggttcgctga tggcagctg
 421 gcttctgact acaacggccc cgcgacgct gatggcattg ttggctgggt gaagaagaag
 481 actggcccc cgcctgtgac cgttgaggac gccgacaagc tgaagtcctt ggaggcggac
 541 gctgaggctcg ttgtcgtcgg ctactcaag gccctggagg gcgagatcta cgacaccttc
 601 aagtcctacg ccgccaaagc cgaggacgctg gtgtctgctg agaccaccag cgccgacgctc

FIG. 2A

661 gccaaaggccg ccggccctgga cccggctggac accgtgtccg tggccaagaa cttcggccggc
721 gaggaccgcb ccaccgccgt cctggccacg gacatcgaca ctgactccct gaccgcttc
781 gtcraagtcgg agaagatgcc cccaccatt gagttcaacc agaagaactc tgacaagatc
841 tccaacagcg gacgcggaga tcatgactgt gttccgctgag gccagcaaga agttcaaggg ccagctggctg
901 ttcgtgaccg tcaacaacga ggctgacggc ggccgacccc ggcttcctggc agaaagaacaa gaagtccg
1021 aagggcgcca cctcgcctgt agttcacggc tgacaacgtg gctaaagtcc cggagagcgt ggtggacggc
1081 atggaggcgg agttcacggc cctcgcctgt gttcacggc gctaaagtcc cggagagcgt ggtggacggc
1141 accgcgcagg ccgtgctcaa gtcggaggcc atcccggagg accctatga agaatggctc
1201 tacaagattg tggcaagac cgtggagctt gttggttctgg agagaccaa agagctgctg
1261 ctggaggctg accgcccctg gttggatctc gtaagaagg tggagcccac ctacaagaaag
1321 ctggccaagc gctttaagaa gttggatctc gttacatctcg ccaagatgga cctgttctta tccgcccggc
1381 aacgagcacc ccgagatcga gttcaaggcc gctgaccgct ggctcaagtc cgtgacccaa
1441 agcgaccgca ccccactct gttcgaaggc gctgaccgct gaggctgccc agaaaggctc cgaaggcggc
1501 ttcatacaga ccaacgcaa gatcccgtac gaggctgccc ggtcccgaca aggacggct gtaagcggct
1561 gaggccacct cggacgaca ggaagctctg ggtcctgaca cttgctgcat gcttggagca
1621 atctgaacta cccaggctt taaggaggag acggagcag caacggcggg gctggcggc
1681 gbatgggagt gtagtcctc gtagcagc gtagcagcag tgcctgcccggg cggtagggg
1741 ccggcagcgc gctgtctgct gtagcagcag ctagcagcag gtagcagcag gctggcggc
1801 gctggcggc gctgtctgct gtagcagcag ctagcagcag gtagcagcag gctggcggc
1861 agagatgaga gctgtacggg ctagcagcag ctagcagcag gtagcagcag gctggcggc
1921 cttgctagga gacgcaaggc tttggcacaag agggacgctg agggacgccc tccacagccc tggactgga
1981 agtttttttag gccctgctgt gtagtggctg gtagtggctg gtagtggctg gtagtggctg gtagtggctg
2041 cgtttctctc aagacggagc tactagtatg ctagcagcag ctagcagcag ctagcagcag ctagcagcag
2101 gtcgcccgac catgaggagt gctgtgttgc gctgtgttgc gctgtgttgc gctgtgttgc gctgtgttgc
2161 gtttccgaa gctggagtca tctgttggagg gctgttggagg gctgttggagg gctgttggagg gctgttggagg
2221 ggcgcgtga tgtccggatg gggatctgca gggatctgca gggatctgca gggatctgca gggatctgca
2281 cagggatcg agctagcga gggatctgca gggatctgca gggatctgca gggatctgca gggatctgca
2341 gtagccaagg cggagtgcac gggagtgcac gggagtgcac gggagtgcac gggagtgcac gggagtgcac
2401 cgccttgcgc aaa

FIG. 2B


```

      .250
CTT CTT TAC GGT AAC AAC ATC ATT ACA GGT GCT GTA ATC CCA ACT TCT AAC GCA ATC GGT
Leu Leu Tyr Gly Asn Asn Ile Ile Thr Gly Ala Val Ile Pro Thr Ser Asn Ala Ile Gly 90
      Ala
      Ile
      Ser
      .300
CTT CAC TTC TAC CCA ATT TGG GAA GCT GCT TCT CTA GAC GAG TGG TTA TAC AAC GGT GGT
Leu His Phe Tyr Pro Ile Trp Glu Ala Ser Leu Asp Glu Trp Leu Trp Asn Gly Gly 110
      Val
      .350
CCT TAC CAA CTT ATC GTT TGT CAC TTC CTT CTA GGT GTA TAC TGC TAC ATG GGT [CGT GAG
Pro Tyr Gln Leu Ile Val Cys His Phe Leu Leu Gly Val Tyr Cys Tyr Met Gly] Arg Glu 130
      Glu
      Leu
      Ala
      .400
TGG GAA TTA TCT TTC CGT TTA GGT ATG CGT CCA TGG ATC GCT GTA GCT TAC TCA GCT CCA
Trp Glu Leu Ser Phe Arg Leu Gly Met Arg Pro Trp Ile Ala Val Ala Tyr Ser Ala Pro 150
      .450
GTA GCT GCA GCT TCA GCT GTA TTC TTA GTT TAC CCT ATC GGC CAA GGT TCA TTC TCT GAC
Val Ala Ala Ala Ser Ala Val Phe Leu Val Tyr Pro Ile Gly Gln Gly Ser Phe Ser Asp 170
      Thr
      Ile
      .500
GGT ATG CCT TTA GGT [ATC TCT GGT ACT TTC AAC TTC ATG ATC GTA TTC CAA GCA GAA CAC
Gly Met Pro Leu Gly] Ile Ser Gly Thr Phe Asn Phe Met Ile Val Phe Gln Ala Glu His 190
      .550
AAC ATC CTT ATG CAC CCA TTC CAC ATG TTA GGT GTT GCT GGT GTA TTC GGT TCA TTA
Asn Ile Leu Met His Pro Phe His Met Leu Gly Val Ala Gly Val Phe Gly Ser Leu 210
      .600
TTC TCA GCT ATG CAC GGT TCT TTA GTT ACT TCA TCT TTA ATC CGT GAA ACA ACT GAA AAC
Phe Ser Ala Met His Gly Ser Leu Val Thr Ser Ser Leu Ile Arg Glu Thr Thr Glu Asn 230
      850

```

FIG. 3B

```

.700
GAA TCA GCT AAC GAA GGT TAC CGT TTC GGT CAA GAA GAA ACT TAC AAC ATT GTA GCT .750
Glu Ser Ala Asn Glu Gly Tyr Arg Phe Gly Gln Glu Glu Thr Tyr Asn Ile Val Ala 250

GCT CAT [GGT TAC TTT GGT CGT CTA ATC TTC CAA TAC GCT TCT TTC AAC AAC TCT CGT TCA
Ala His] [Gly Tyr Phe Gly Arg Leu Ile Phe Gln Tyr Ala Ser Phe Asn Asn Ser Arg Ser 270
.800

TTA CAC TTC TTC TTA GCT GCT TGG CCG GTA ATC GGT ATT TGG TTC ACT GCT TTA GGT TTA
Leu His Phe Leu Ala Ala Trp Pro Val Ile Gly Ile Trp Phe Thr Ala Leu Gly Leu 290
Val
.850


TCA ACT ATG GCA TTC AAC TTA AAC GGT TTC AAC TTC AAC CAA TCA GTA GAC TCA CAA
Ser Thr Met Ala Phe Asn Leu Asn Gly Phe Asn Phe Asn Gln Ser Val Val Asp Ser Gln 310
.900

GGT CGT GTA CTA AAC ACT TGG GCA GAC ATC AAC CGT GCT AAC TTA GGT ATG GAA GTA
Gly Arg Val Leu Asn Thr Trp Ala Asp Ile Ile Asn Arg Ala Asn Leu Gly Met Glu Val 330
Ile
.950

ATG CAC GAG CGT AAC GCT CAC AAC TTC CCT CTA GAC TTA GCT TCA ACT AAC TCT AGC TCA
Met His Glu Arg Asn Ala His Asn Phe Pro Leu Asp Leu Ala Ser Thr Asn Ser Ser Ser 350
Ala Ile Glu Ala Pro
.1000
AAC AAC TAA TTT TTTTAAACTAAATAAATCTGGTTAACCATACCTAGTTTATTATTAGTTTATACACTTTT
Asn Asn *Oc
Thr Gly *Oc
.1100
CATATATATACTTAATAGCTACCATAGGCAGTTGGCAGGACGTCCC
.1150

```

FIG. 3C



TS = transcription start and transcription stop

FIG. 4

661 ACG TTG TAC ACC AAC GTG TTC GTC AAC TTG CCG GCC GAC ATC GGC GAC GAG GAG CTG 720
 221 T L Y T N V F V K N L P A D I G D D E L 240
 721 GGC AAG ATG GCC ACC GAG CAC GGC GAG ATC ACC AGC GCG GTG GTC ATG AAG GAC GAC AAG 780
 241 G K M A T E H G E I T S A V V M K D D K 260
 781 GGC GGC AGC AAG GGC TTC GGC TTC AAC TTC AAG GAC GCC GAG TCG GCG GCC AAG TGC 840
 261 G G S K G F G F I N F K D A E S A A K C 280
 841 GTG GAG TAC CTG AAC GAG CGC GAG ATG AGC GGC AAG ACC CTG TAC GCC GGC CGC GCC CAG 900
 281 V E Y L N E R E M S G K T L Y A G R A Q 300
 901 AAG AAG ACC GAG CGC GAG GCG ATG CTG CGC CAG AAG GCC GAG AGC AAG CAG GAG CGT 960
 301 K K T E R E A M L R Q K A E S K Q E R 320
 961 TAC CTG AAG TAC CAG AGC ATG AAC CTG TAC GTC AAC AAC CTG TCC GAC GAG GAG GTC GAC 1020
 321 Y L K Y Q S M N L Y V K N L S D E E V D 340
 1021 GAC GAC GCC CTG CGT GAG CTG TTC GCC AAC TCT GGC ACC ATC ACC TCG TGC AAG GTC ATG 1080
 341 D D A L R E L F A N S G T I T S C K V M 360
 1081 AAG GAC GGC AGC GGC AAG TCC AAG GGC TTC GGC TTC GTG TGC ACC AGC CAC GAC GAG 1140
 361 K D G S G K S K G F G F V C F T S H D E 380
 1141 GCC ACC CCG CCG CCC GTG ACC GAG ATG AAC GGC AAG ATG GTC AAG GGC AAG CCC CTG TAC 1200
 381 A T R P P V T E M N G K M V K G K P L Y 400
 1201 GTG GCC CTG GCG CAG CGC AAG GAC GTG CGC CGT GCC ACC CAG CTG GAG GCC AAC ATG CAG 1260
 401 V A L A Q R K D V R R A T Q L E A N M Q 420
 1261 GCG CGC ATG TAA GGATCC 1278
 421 A R M * 424

FIG. 5B




FIG. 6




FIG. 7





FIG. 8



Bacterial luciferase A and B proteins expressed from a single mRNA containing the psbA 5' UTR with translational activator element.

FIG. 9



FIG. 10




FIG. 11

EXPRESSION OF EUKARYOTIC POLYPEPTIDES IN CHLOROPLASTS

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue.

[This is a stage application filed under 35 USC 371, of PCT/US98/00840, filed Jan. 16, 1998. This application claims benefit of provisional No. 60/035,955 filed Jan. 17, 1997 and provisional appln No. 60/069,400 filed Dec. 12, 1997.] *Notice: More than one reissue application has been filed for the reissue of U.S. Pat. No. 6,156,517. The reissue applications are U.S. application Ser. No. 11/197,730 (the present application) and Ser. No. 10/310,587, issued on Oct. 17, 2006 as U.S. Pat. No. Re. 39,350, all of which are continuation reissues of U.S. Pat. No. 6,156,517. This application is a continuation reissue application of U.S. application Ser. No. 10/310,587, filed Dec. 4, 2002 now U.S. Pat. No. Re. 39,350, which is a reissue application of U.S. Pat. No. 6,156,517, issued Dec. 5, 2000, which is a national stage application filed under 35 U.S.C. §371, of PCT/US98/00840, filed Jan. 16, 1998. This application claims benefit of priority to U.S. Provisional Application Ser. No. 60/035,955, filed Jan. 17, 1997 and U.S. Provisional Patent Application Ser. No. 60/069,400, filed Dec. 12, 1997.*

This invention was made with government support under Contract No. GM 54659 by the National Institutes of Health and Contract No. DO-FG03-93ER20116 by the U.S. Department of Energy. The government has certain rights in the invention.

TECHNICAL FIELD

The invention relates to expression systems and methods for expression of desired genes and gene products in cells. Particularly, the invention relates to a gene encoding a RNA binding protein useful for regulating gene expression in cells, the protein binding site, a gene encoding a regulating protein disulfide isomerase and methods and systems for gene expression of recombinant molecules.

BACKGROUND

Expression systems for expression of exogenous foreign genes in eukaryotic and prokaryotic cells are basic components of recombinant DNA technology. Despite the abundance of expression systems and their wide-spread use, they all have characteristic disadvantages. For example, while expression in *E. coli* is probably the most popular as it is easy to grow and is well understood, eukaryotic proteins expressed therein are not properly modified. Moreover, those proteins tend to precipitate into insoluble aggregates and are difficult to obtain in large amounts. Mammalian expression systems, while practical on small-scale protein production, are more difficult, time-consuming and expensive than in *E. coli*.

A number of plant expression systems exist as well as summarized in U.S. Pat. No. 5,234,834, the disclosures of which are hereby incorporated by reference. One advantage of plants or algae in an expression system is that they can be used to produce pharmacologically important proteins and enzymes on a large scale and in relatively pure form. In addition, micro-algae have several unique characteristics that make them ideal organisms for the production of proteins on

a large scale. First, unlike most systems presently used to produce transgenic proteins, algae can be grown in minimal media (inorganic salts) using sunlight as the energy source. These algae can be grown in contained fermentation vessels or on large scale in monitored ponds. Ponds of up to several acres are routinely used for the production of micro-algae. Second, plants and algae have two distinct compartments, the cytoplasm and the chloroplast, in which proteins can be expressed. The cytoplasm of algae is similar to that of other eukaryotic organisms used for protein expression, like yeast and insect cell cultures. The chloroplast is unique to plants and algae and proteins expressed in this environment are likely to have properties different from those of cytoplasmically expressed proteins.

The present invention describes an expression system in which exogenous molecules are readily expressed in either prokaryotic or eukaryotic hosts and in either the cytoplasm or chloroplast. These beneficial attributes are based on the discovery and cloning of components of translation regulation in plants as described in the present invention.

Protein translation plays a key role in the regulation of gene expression across the spectrum of organisms (Kozak, *Ann. Rev. Cell Biol.*, 8:197-225 (1992) and de Smit and Van Duin, *Prog. Nucleic Acid Res. Mol. Biol.*, 38:1-35 (1990)). The majority of regulatory schemes characterized to date involve translational repression often involving proteins binding to mRNA to limit ribosome association (Winter et al., *Proc. Natl. Acad. Sci., USA*, 84:7822-7826 (1987) and Tang and Draper, *Biochem.*, 29:4434-4439 (1990)). Translational activation has also been observed (Wulczyn and Kahmann, *Cell*, 65:259-269 (1991)), but few of the underlying molecular mechanisms for this type of regulation have been identified. In plants, light activates the expression of many genes. Light has been shown to activate expression of specific chloroplast encoded mRNAs by increasing translation initiation (Mayfield et al., *Ann. Rev. Plant Physiol. Plant Mol. Biol.*, 46:147-166 (1995) and Yohn et al., *Mol. Cell Biol.*, 16:3560-3566 (1996)). Genetic evidence in higher plants and algae has shown that nuclear encoded factors are required for translational activation of specific chloroplast encoded mRNAs (Rochaix et al., *Embo J.*, 8:1013-1021 (1989), Kuchka et al., *Cell*, 58:869-876 (1989), Girard-Bascou et al., *Embo J.*, 13:3170-3181 (1994), Kim et al., *Plant Mol. Biol.*, 127:1537-1545 (1994).

In the green algae *Chlamydomonas reinhardtii*, a number of nuclear mutants have been identified that affect translation of single specific mRNAs in the chloroplast, often acting at translation initiation (Yohn et al., *supra*, (1996)). Mutational analysis of chloroplast mRNAs has identified sequence elements within the 5' untranslated region (UTR) of mRNAs that are required for translational activation (Mayfield et al., *supra*, (1995), Mayfield et al., *J. Cell Biol.*, 127:1537-1545 (1994) and Rochaix, *Ann. Rev. Cell Biol.*, 8:1-28 (1992)), and the 5' UTR of a chloroplast mRNA can confer a specific translation phenotype on a reporter gene in vivo (Zerges and Rochaix, *Mol. Cell Biol.*, 14:5268-5277 (1994) and Staub and Maliga, *Embo J.*, 12:601-606 (1993).

Putative translational activator proteins were identified by purifying a complex of four proteins that binds with high affinity and specificity to the 5' UTR of the chloroplast encoded *psbA* mRNA [encoding the D1 protein, a major component of Photosystem II (PS II)] (Danon and Mayfield, *Embo J.*, 10:3993-4001 (1991)). Binding of these proteins to the 5' UTR of *psbA* mRNA correlates with translation of this mRNA under a variety of physiological (Danon and Mayfield, *id.*, (1991)) and biochemical conditions (Danon and Mayfield, *Science*, 266:1717-1719 (1994) and Danon and

Mayfield, *Embo J.*, 13:2227-2235 (1994)), and in different genetic backgrounds (Yohn et al., *supra*, (1996)). The binding of this complex to the *psbA* mRNA can be regulated in vitro in response to both redox potential (Danon and Mayfield, *Science*, 266:1717-1719 (1994)) and phosphorylation (Danon and Mayfield, *Embo J.*, 13:2227-2235 (1994)), both of which are thought to transduce the light signal to activate translation of *psbA* mRNA. The 47 kDa member of the *psbA* RNA binding complex (RB47) is in close contact with the RNA, and antisera specific to this protein inhibits binding to the *psbA* mRNA in vitro (Danon and Mayfield, *supra*, (1991)).

Although the translational control of *psbA* mRNA by RB47 has been reported, the protein has not been extensively characterized and the gene encoding RB47 has not been identified, cloned and sequenced. In addition, the regulatory control of the activation of RNA binding activity to the binding site by nuclear-encoded trans-acting factors, such as RB60, have not been fully understood. The present invention now describes the cloning and sequencing of both RB47 and RB60. Based on the translation regulation mechanisms of RB47 and RB60 with the RB47 binding site, the present invention also describes a translation regulated expression system for use in both prokaryotes and eukaryotes.

BRIEF DESCRIPTION OF THE INVENTION

The RB47 gene encoding the RB47 activator protein has now been cloned and sequenced, and the target binding site for RB47 on messenger RNA (mRNA) has now been identified. In addition, a regulatory protein disulfide isomerase, a 60 kilodalton protein referred to as RB60, has also been cloned, sequenced and characterized. Thus, the present invention is directed to gene expression systems in eukaryotic and prokaryotic cells based on translational regulation by RB47 protein, its binding site and the RB60 regulation of RB47 binding site activation.

More particularly, the present invention describes the use of the RB47 binding site, i.e., a 5' untranslated region (UTR) of the chloroplast *psbA* gene, in the context of an expression system for regulating the expression of genes encoding a desired recombinant molecule. Protein translation is effected by the combination of the RB47 binding site and the RB47 binding protein in the presence of protein translation components. Regulation can be further imposed with the use of the RB60 regulatory protein disulfide isomerase. Therefore, the present invention describes reagents and expression cassettes for controlling gene expression by affecting translation of a coding nucleic acid sequence in a cell expression system.

Thus, in one embodiment, the invention contemplates a RB47 binding site sequence, i.e., a mRNA sequence, typically a mRNA leader sequence, which contains the RB47 binding site. A preferred RB47 binding site is *psbA* mRNA. For use in expressing recombinant molecules, the RB47 binding site is typically inserted 5' to the coding region of the preselected molecule to be expressed. In a preferred embodiment, the RB47 binding site is inserted into the 5' untranslated region along with an upstream *psbA* promoter to drive the expression of a preselected nucleic acid encoding a desired molecule. In alternative embodiments, the RB47 binding site is inserted into the regulatory region downstream of any suitable promoter present in a eukaryotic or prokaryotic expression vector. Preferably, the RB47 binding site is positioned within 100 nucleotides of the translation initiation site. In a further aspect, 3' to the coding region is a 3' untranslated region (3' UTR) necessary for transcription termination and RNA processing.

Thus, in a preferred embodiment, the invention contemplates an expression cassette or vector that contains a transcription unit constructed for expression of a preselected nucleic acid or gene such that upon transcription, the resulting mRNA contains the RB47 binding site for regulation of the translation of the preselected gene transcript through the binding of the activating RB47 protein. The RB47 protein is provided endogenously in a recipient cell and/or is a recombinant protein expressed in that cell.

Thus, the invention also contemplates a nucleic acid molecule containing the sequence of the RB47 gene. The nucleic acid molecule is preferably in an expression vector capable of expressing the gene in a cell for use in interacting with a RB47 binding site. The invention therefore contemplates an expressed recombinant RB47 protein. In one embodiment, the RB47 binding site and RB47 encoding nucleotide sequences are provided on the same genetic element. In alternative embodiments, the RB47 binding site and RB47 encoding nucleotide sequences are provided separately.

The invention further contemplates a nucleic acid molecule containing the sequence encoding the 69 kilodalton precursor to RB47. In alternative embodiments, the RB47 nucleic acid sequence contains a sequence of nucleotides to encode a histidine tag. Thus, the invention relates to the use of recombinant RB47, precursor RB47, and histidine-modified RB47 for use in enhancing translation of a desired nucleic acid.

The invention further contemplates a nucleic acid molecule containing a nucleotide sequence of a polypeptide which regulates the binding of RB47 to RB47 binding site. A preferred regulatory molecule is the protein disulfide isomerase RB60. The RB60-encoding nucleic acid molecule is preferably in an expression vector capable of expressing the gene in a cell for use in regulating the interaction of RB47 with a RB47 binding site. Thus, the invention also contemplates an expressed recombinant RB60 protein. In one embodiment, the RB47 binding site, RB47 encoding and RB60 encoding nucleotide sequences are provided on the same genetic element. In alternative embodiments, the expression control nucleotide sequences are provided separately. In a further aspect, the RB60 gene and RB47 binding site sequence are provided on the same construct.

The invention can therefore be a cell culture system, an in vitro expression system or a whole tissue, preferably a plant, in which the transcription unit is present that contains the RB47 binding site and further includes a (1) transcription unit capable of expressing RB47 protein or (2) the endogenous RB47 protein itself for the purpose of enhancing translation of the preselected gene having an RB47 binding site in the mRNA. Preferred cell culture systems are eukaryotic and prokaryotic cells. Particularly preferred cell culture systems include plants and more preferably algae.

A further preferred embodiment includes (1) a separate transcription unit capable of expressing a regulatory molecule, preferably RB60 protein, or (2) the endogenous RB60 protein itself for the purpose of regulating translation of the preselected gene having an RB47 binding site in the mRNA. In an alternative preferred embodiment, one transcription unit is capable of expressing both the RB47 and RB60 proteins. In a further aspect, the RB47 binding site sequence and RB60 sequence are provided on the same construct.

In one aspect of the present invention, plant cells endogenously containing RB47 and RB60 proteins are used for the expression of recombinant molecules, such as proteins or polypeptides, through activation of the RB47 binding in an exogenously supplied expression cassette. Alternatively, stable plant cell lines containing endogenous RB47 and RB60 are first generated in which RB47 and/or RB60 proteins are

overexpressed. Overexpression is obtained preferably through the stable transformation of the plant cell with one or more expression cassettes for encoding recombinant RB47 and RB60. In a further embodiment, stable cell lines, such as mammalian or bacterial cell lines, lacking endogenous RB47 and/or RB60 proteins are created that express exogenous RB47 and/or RB60.

Plants for use with the present invention can be a transgenic plant, or a plant in which the genetic elements of the invention have been introduced. Based on the property of controlled translation provided by the combined use of the RB47 protein and the RB47 binding site, translation can be regulated for any gene product, and the system can be introduced into any plant species. Similarly, the invention is useful for any prokaryotic or eukaryotic cell system.

Methods for the preparation of expression vectors is well known in the recombinant DNA arts, and for expression in plants is well known in the transgenic plant arts. These particulars are not essential to the practice of the invention, and therefore will not be considered as limiting.

The invention allows for high level of protein synthesis in plant chloroplasts and in the cytoplasm of both prokaryotic and eukaryotic cells. Because the chloroplast is such a productive plant organ, synthesis in chloroplasts is a preferred site of translation by virtue of the large amounts of protein that can be produced. This aspect provides for great advantages in agricultural production of mass quantities of a pre-selected protein product.

The invention further provides for the ability to screen for agonists or antagonists of the binding of RB47 to the RB47 binding site using the expression systems as described herein. Antagonists of the binding are useful in the prevention of plant propagation.

Also contemplated by the present invention is a screening assay for agonists or antagonists of RB60 in a manner analogous to that described above for RB47. Such agonists or antagonists would be useful in general to modify expression of RB60 as a way to regulate cellular processes in a redox manner.

Kits containing expression cassettes and expression systems, along with packaging materials comprising a label with instructions for use, as described in the claimed embodiments are also contemplated for use in practicing the methods of this invention.

Other uses will be apparent to one skilled in the art in light of the present disclosures.

BRIEF DESCRIPTION OF DRAWINGS

In the figures forming a portion of this disclosure:

FIGS. 1A-1D show the complete protein amino acid residue sequence of RB47 is shown from residues 1-623, together with the corresponding nucleic acid sequence encoding the RB47 sequence, from base 1 to base 2732. The nucleotide coding region is shown from base 197-2065, the precursor form. The mature form is from nucleotide position 197-1402. Also shown is the mRNA leader, bases 1-196, and poly A tail of the mRNA, bases 2066-2732. Both the nucleotide and amino acid sequence are listed in SEQ ID NO 5.

FIGS. 2A-2B show the complete protein amino acid residue sequence of RB60 is shown from residues 1-488, together with the corresponding nucleic acid sequence from base 1 to base 2413, of which bases 16-1614 encode the RB60 sequence. Both the nucleotide and amino acid sequence are listed in SEQ ID NO 10.

FIGS. 3A-3C show the complete sequence of the psbA mRNA, showing both encoded psbA protein amino acid resi-

due sequence (residues 1-352) and the nucleic acid sequence as further described in Example 3 is illustrated. Both the nucleotide and amino acid sequence are listed in SEQ ID NO 13.

FIG. 4 is a schematic diagram of an expression cassette containing on one transcription unit from 5' to 3', a promoter region derived from the psbA gene for encoding the D1 protein from *C. reinhardtii* further containing a transcription initiation site (TS), the RB47 binding site, a region for insertion of a foreign or heterologous coding region, a RB47 coding region, a RB60 coding region, and the 3' flanking region containing transcription termination site (TS), flanked by an origin of replication and selection marker. Restriction endonuclease sites for facilitating insertion of the independent genetic elements are indicated and further described in Example 4A.

FIGS. 5A-5B show the nucleotide and amino acid sequence of the RB47 molecule containing a histidine tag, the sequences of which are also listed in SEQ ID NO 14.

FIG. 6 is a schematic diagram of an expression cassette containing on one transcription unit from 5' to 3', a promoter region derived from the psbA gene for encoding the D1 protein from *C. reinhardtii* further containing a transcription initiation site (TS), the RB47 binding site, a region for RB47 is also shown in FIGS. 1A-1D (SEQ ID NO 5). As described in Section 2 above, the predicted protein sequence from the cloned cDNA contained both the derived peptide sequences of RB47 and is highly homologous to poly(A) binding proteins (PABP) from a variety of eukaryotic organisms.

FIG. 7 diagrams a construct is essentially pD1/Nde including a heterologous coding sequence having a 3' XbaI restriction site for ligation with the 3' psbA gene.

FIG. 8 shows two of the transformants that contained the single chain chimeric gene produced single chain antibodies at approximately 1% of total protein levels.

FIG. 9 shows a construct, the bacterial LuxAB coding region was ligated between the psbA 5' UTR and the psbA 3' end in an *E. coli* plasmid.

FIG. 10 shows luciferase activity accumulated with the chloroplast.

FIG. 11 shows a construct engineered so that the psbA promoter and 5' UTR are used to drive the synthesis of the light chain and heavy chains of an antibody, and the J chain normally associated with IgA molecules.

2. Cloning of RB60

To clone the cDNA encoding the 60 kDa psbA mRNA binding protein (RB60), the psbA-specific RNA binding proteins were purified from light-grown *C. reinhardtii* cells using heparin-agarose chromatography followed by psbA RNA affinity chromatography (RAC). RAC-purified proteins were separated by two-dimensional polyacrylamide gel electrophoresis. The region corresponding to RB60 was isolated from the PVDF membrane. RB60 protein was then digested with trypsin. Unambiguous amino acid sequences were obtained from two peptide tryptic fragments (WFVDGELASDYNGPR (SEQ ID NO 6) and (QLILWTTAD-DLKADAEIMTVFR (SEQ ID NO 7)) as described above for RB47. The calculated molecular weights of the two tryptic peptides used for further analysis precisely matched with the molecular weights determine by mass spectrometry. The DNA sequence corresponding to one peptide of 22 amino acid residues was amplified by PCR using degenerate oligonucleotides, the forward primer 5'CGCGGATCCGAYGCB-GAGATYATGAC3' (SEQ ID NO 8) and the reverse primer 5'CGCGAATTCGTCATRATCTCVGCRTC3' (SEQ ID NO

9), where R can be A or G (the other IUPAC nucleotides have been previously defined above). The amplified sequence was then used to screen a λ -gt10 cDNA library from *C. reinhardtii*. Three clones were identified with the largest being 2.2 kb. Selection and sequencing was performed as described for RB47 cDNA.

The resulting RB60 cDNA sequence is available via GenBank (Accession Number AF027727). The nucleotide and encoded amino acid sequence of RB60 is also shown in FIGS. 2A-2B (SEQ ID NO 10). The protein coding sequence of 488 amino acid residues corresponds to nucleotide positions 16-1614 of the 2413 base pair sequence. The predicted amino acid sequence of the cloned cDNA contained the complete amino acid sequences of the two tryptic peptides. The amino acid sequence of the encoded protein revealed that it has high sequence homology to both plant and mammalian protein disulfide isomerase (PDI), and contains the highly conserved thioredoxin-like domains with —CysGlyHisCys— (—CGHC—) (SEQ ID NO 11) catalytic sites in both the N-terminal and C-terminal regions and the —LysAspGluLeu— (—KDEL—) (SEQ ID NO 12) endoplasmic reticulum (ER) retention signal at the C-terminus found in all PDIs. PDI is a multifunctional protein possessing enzymatic activities for the formation, reduction, and isomerization of disulfide bonds during protein folding, and is typically found in the ER. The first 30 amino acid residues of RB60 were found to lack sequence homology with the N-terminal signal sequence of PDI from plants or mammalian cells. However, this region has characteristics of chloroplast transit peptides of *C. reinhardtii*, which have similarities with both mitochondrial and higher plant chloroplast presequences. A transit peptide sequence should override the function of the —KDEL— ER retention signal and target the protein to the chloroplast since the —KDEL— signal acts only to retain the transported protein in the ER.

3. Preparation of psbA Promoter Sequence and RB47 Binding Site Nucleotide Sequence

The chloroplast psbA gene from the green unicellular alga *C. reinhardtii* was cloned and sequenced as described by Erickson et al., Embo J., 3:2753-2762 (1984), the disclosure of which is hereby incorporated by reference. The DNA sequence of the coding regions and the 5' and 3' untranslated (UTR) flanking sequences of the *C. reinhardtii* psbA gene is shown in FIGS. 3A-3C. The psbA gene sequence is also available through GenBank as further discussed in Example 4. The nucleotide sequence is also listed as SEQ ID NO 13. The deduced amino acid sequence (also listed in SEQ ID NO 13) of the coding region is shown below each codon beginning with the first methionine in the open reading frame. Indicated in the 5' non-coding sequence are a putative Shine-Dalgarno sequence in the dotted box, two putative transcription initiation sites determined by S1 mapping (S1) and the Pribnow-10 sequence in the closed box. Inverted repeats of eight or more base pairs are marked with arrows and labeled A-D. A direct repeat of 31 base pairs with only two mismatches is marked with arrows labeled 31. Indicated in the 3' non-coding sequence is a large inverted repeat marked by a forward arrow and the SI cleavage site marking the 3' end of the mRNA. Both the 5' and 3' untranslated regions are used in preparing one of the expression cassettes of this invention as further described below.

The 5' UTR as previously discussed contains both the psbA promoter and the RB47 binding site. The nucleotide sequence defining the psbA promoter contains the region of the psbA DNA involved in binding of RNA polymerase to initiate

transcription. The -10 sequence component of the psbA promoter is indicated by the boxed nucleotide sequence upstream of the first S1 while the -35 sequence is located approximately 35 bases before the putative initiation site. As shown in FIGS. 3A-3C, the -10 sequence is boxed, above which is the nucleotide position (-100) from the first translated codon. The -35 sequence is determined accordingly. A psbA promoter for use in an expression cassette of this invention ends at the first indicated S1 site (nucleotide position -92 as counting from the first ATG) in FIGS. 3A-3C and extends to the 5' end (nucleotide position -251 as shown in FIGS. 3A-3C). Thus, the promoter region is 160 bases in length. A more preferred promoter region extends at least 100 nucleotides to the 5' end from the S1 site. A most preferred region contains nucleotide sequence ending at the s1 site and extending 5' to include the -35 sequence, i.e., from -92 to -130 as counted from the first encoded amino acid residue (39 bases).

The psbA RB47 binding site region begins at the first S1 site as shown in FIGS. 3A-3C and extends to the first adenine base of the first encoded methionine residue. Thus, a psbA RB47 binding site in the psbA gene corresponds to the nucleotide positions from -91 to -1 as shown in FIG. 3A-3C.

The above-identified regions are used to prepare expression constructs as described below. The promoter and RB47 binding site regions can be used separately; for example, the RB47 binding site sequence can be isolated and used in a eukaryotic or prokaryotic plasmid with a non-psbA promoter. Alternatively, the entire psbA 5' UTR having 251 nucleotides as shown in FIGS. 3A-3C is used for the regulatory region in an expression cassette containing both the psbA promoter and RB47 binding site sequence as described below.

4. Preparation of Expression Vectors and Expression of Coding Sequences

A. Constructs Containing an psbA Promoter, an RB47 Binding Site Nucleotide Sequence, a Desired Heterologous Coding Sequence, an RB47-Encoding Sequence and an RB60-Encoding Sequence

Plasmid expression vector constructs, alternatively called plasmids, vectors, constructs and the like, are constructed containing various combinations of elements of the present invention as described in the following examples. Variations of the positioning and operably linking of the genetic elements described in the present invention and in the examples below are contemplated for use in practicing the methods of this invention. Methods for manipulating DNA elements into operable expression cassettes are well known in the art of molecular biology. Accordingly, variations of control elements, such as constitutive or inducible promoters, with respect to prokaryotic or eukaryotic expression systems as described in Section C. are contemplated herein although not enumerated. Moreover, the expression the various elements is not limited to one transcript producing one mRNA; the invention contemplates protein expression from more than one transcript if desired.

As such, while the examples below recite one or two types of expression cassettes, the genetic elements of RB47 binding site, any desired coding sequence, in combination with RB47 and RB60 coding sequences along with a promoter are readily combined in a number of operably linked permutations depending on the requirements of the cell system selected for the expression. For example, for expression in a chloroplast, endogenous RB47 protein is present therefore an expression cassette having an RB47 binding site and a desired coding sequence is minimally required along with an operative promoter sequence. Overexpression of RB47 may be preferable to enhance the translation of the coding sequence; in that case, the chloroplast is further transformed with an expression

cassette containing an RB47-encoding sequence. Although the examples herein and below utilize primarily the sequence encoding the precursor form of RB47, any of the RB47-encoding sequences described in the present invention, i.e., RB47 precursor, mature RB47 and histidine-modified RB47 are contemplated for use in any expression cassette and system as described herein. To regulate the activation of translation, an RB60-encoding element is provided to the expression system to provide the ability to regulate redox potential in the cell as taught in Section B. These examples herein and below represent a few of the possible permutations of genetic elements for expression in the methods of this invention.

In one embodiment, a plasmid is constructed containing an RB47 binding site directly upstream of an inserted coding region for a heterologous protein of interest, and the RB47 and RB60 coding regions. Heterologous refers to the nature of the coding region being dissimilar and not from the same gene as the regulatory molecules in the plasmid, such as RB47 and RB60. Thus, all the genetic elements of the present invention are produced in one transcript from the IPTG-inducible *psbA* promoter. Alternative promoters are similarly acceptable.

The final construct described herein for use in a prokaryotic expression system makes a single mRNA from which all three proteins are translated. The starting plasmid is any *E. coli* based plasmid containing an origin of replication and selectable marker gene. For this example, the Bluescript plasmid, pBS, commercially available through Stratagene, Inc., La Jolla, Calif., which contains a polylinker-cloning site and an ampicillin resistant marker is selected for the vector.

The wild-type or native *psbA* gene (Erickson et al., *Embo J.*, 3:2753-2762 (1984), also shown in FIGS. 3A-3C, is cloned into pBS at the EcoRI and BamHI sites of the polylinker. The nucleotide sequence of the *psbA* gene is available on GenBank with the 5' UTR and 3' UTR respectively listed in Accession Numbers X01424 and X02350. The EcoRI site of *psbA* is 1.5 kb upstream of the *psbA* initiation codon and the BamHI site is 2 kb downstream of the stop codon. This plasmid is referred to as pDI.

Using site-directed PCR mutagenesis, well known to one of ordinary skill in the art, an NdeI site is placed at the initiation codon of *psbA* in the pDI plasmid so that the ATG of the NdeI restriction site is the ATG initiation codon. This plasmid is referred to as pDI/Nde. An Nde site is then placed at the initiation codon of the gene encoding the heterologous protein of interest and an Xho I site is placed directly downstream (within 10 nucleotides) of the TAA stop codon of the heterologous protein coding sequence. Again using site-directed mutagenesis, an XhoI site is placed within 10 nucleotides of the initiation codon of RB47, the preparation of which is described in Example 2, and an NotI site is placed directly downstream of the stop codon of RB47. The heterologous coding region and the RB47 gene are then ligated into pDI/Nde so that the heterologous protein gene is directly adjacent to the RB47 binding site and the RB47 coding region is downstream of the heterologous coding region, using the Xho I site at the heterologous stop codon and the Not I site of the pDI polylinker.

These genetic manipulations result in a plasmid containing the 5' end of the *psbA* gene including the promoter region and with the RB47 binding site immediately upstream of a heterologous coding region, and the RB47 coding region immediately downstream of the heterologous coding region. The nucleotides between the stop codon of the heterologous coding region and the initiation codon of the RB47 coding region is preferably less than 20 nucleotides and preferably does not

contain any additional stop codons in any reading frame. This plasmid is referred to as pDI/RB47.

Using site-directed mutagenesis, a NotI site is placed immediately (within 10 nucleotides) upstream of the initiation codon of RB60, the preparation of which is described in Example 2, and an Xba I site is placed downstream of the RB60 stop codon. This DNA fragment is then ligated to the 3' end of the *psbA* gene using the Xba I site found in the 3' end of the *psbA* gene so that the *psbA* 3' end is downstream of the RB60 coding region. This fragment is then ligated into the pDI/RB47 plasmid using the NotI and BamHI sites so that the RB60 coding region directly follows the RB47 coding region. The resulting plasmid is designated pDI/RB47/RB60. Preferably there is less than 20 nucleotides between the RB47 and RB60 coding regions and preferably there are no stop codons in any reading frame in that region. The final plasmid thus contains the following genetic elements operably linked in the 5' to 3' direction: the 5' end of the *psbA* gene with a promoter capable of directing transcription in chloroplasts, an RB47 binding site, a desired heterologous coding region, the RB47 coding region, the RB60 coding region, and the 3' end of the *psbA* gene which contains a transcription termination and mRNA processing site, and an *E. coli* origin of replication and ampicillin resistance gene. A diagram of this plasmid with the restriction sites is shown in FIG. 4.

Expression of pDI/RB47/RB60 in *E. coli* to produce recombinant RB47, RB60 and the recombinant heterologous protein is performed as described in Example 4B. The heterologous protein is then purified as further described.

Expression cassettes in which the sequences encoding RB47 and RB60 are similarly operably linked to a heterologous coding sequence having the *psbA* RB47 binding site as described in Example 3 are prepared with a different promoter for use in eukaryotic, such as mammalian expression systems. In this aspect, the cassette is similarly prepared as described above with the exception that restriction cloning sites are dependent upon the available multiple cloning sites in the recipient vector. Thus, the RB47 binding site prepared in Example 3 is prepared for directed ligation into a selected expression vector downstream of the promoter in that vector. The RB47 and RB60 coding sequences are obtained from the pDI/RB47/RB60 plasmid by digestion with XhoI and XbaI and inserted into a similarly digested vector if the sites are present. Alternatively, site-directed mutagenesis is utilized to create appropriate linkers. A desired heterologous coding sequence is similarly ligated into the vector for expression.

B. Constructs Containing RB47 Nucleotide Sequence

1) Purified Recombinant RB47 Protein

In one approach to obtain purified recombinant RB47 protein, the full length RB47 cDNA prepared above was cloned into the *E. coli* expression vector pET3A (Studier et al., *Methods Enzymol.*, 185:60-89 (1990)), also commercially available by Novagen, Inc., Madison, Wis. and transformed into BL21 *E. coli* cells. The cells were grown to a density of 0.4 (OD₆₀₀), then induced with 0.5 mM IPTG. Cells were then allowed to grow for an additional 4 hours, at which point they were pelleted and frozen.

Confirmation of the identity of the cloned cDNA as encoding the authentic RB47 protein was accomplished by examining protein expressed from the cDNA by immunoblot analysis and by RNA binding activity assay. The recombinant RB47 protein produced when the RB47 cDNA was expressed was recognized by antisera raised against the *C. reinhardtii* RB47 protein. The *E. coli* expressed protein migrated at 80 kDa on SDS-PAGE, but the protein was actually 69 kDa, as determined by mass spectrometry of the *E. coli* expressed protein. This mass agrees with the mass predicted from the

cDNA sequence. A 60 kDa product was also produced in *E. coli*, and recognized by the antisera against the *C. reinhardtii* protein, which is most likely a degradation or early termination product of the RB47 cDNA. The recombinant RB47 protein expressed from the RB47 cDNA is recognized by the antisera raised against the *C. reinhardtii* protein at levels similar to the recognition of the authentic *C. reinhardtii* RB47 protein, demonstrating that the cloned cDNA produces a protein product that is immunologically related to the naturally produced RB47 protein. In order to generate a recombinant equivalent of the endogenous native RB47, the location of the 47 kDa polypeptide was mapped on the full-length recombinant protein by comparing mass spectrometric data of tryptic digests of the *C. reinhardtii* 47 kDa protein and the full-length recombinant protein. Thus, peptide mapping by mass spectrometry has shown that the endogenous RB47 protein corresponds primarily to the RNA binding domains contained within the N-terminal region of the predicted precursor protein, suggesting that a cleavage event is necessary to produce the mature 47 kDa protein. Thus, full-length recombinant RB47 is 69 kDa and contains a carboxy domain that is cleaved in vivo to generate the endogenous mature form of RB47 that is 47 kDa.

To determine if the heterologously expressed RB47 protein was capable of binding the *psbA* RNA, the *E. coli* expressed protein was purified by heparin agarose chromatography. The recombinant RB47 protein expressed in *E. coli* was purified using a protocol similar to that used previously for purification of RB47 from *C. reinhardtii*. Approximately 5 g of *E. coli* cells grown as described above were resuspended in low salt extraction buffer (10 mM Tris [pH 7.5], 10 mM NaCl, 10 mM MgCl₂, 5 mM β-mercaptoethanol) and disrupted by sonication. The soluble cell extract was applied to a 5 mL Econo-Pac heparin cartridge (Bio-Rad) which was washed prior to elution of the RB47 protein (Danon and Mayfield, *Embo J.*, 10:3993-4001 (1991)).

The *E. coli* expressed protein that bound to the heparin agarose matrix was eluted from the column at the same salt concentration as used to elute the authentic *C. reinhardtii* RB47 protein. This protein fraction was used in *in vitro* binding assays with the *psbA* 5' UTR. Both the 69 and 60 kDa *E. coli* expressed proteins crosslinked to the radiolabeled *psbA* 5' UTR at levels similar to crosslinking of the endogenous RB47 protein, when the RNA/protein complex is subjected to UV irradiation.

Heparin agarose purified proteins, both from the *E. coli* expressed RB47 cDNA and from *C. reinhardtii* cells, were used in an RNA gel mobility shift assay to determine the relative affinity and specificity of these proteins for the 5' UTR of the *psbA* mRNA. The *E. coli* expressed proteins bound to the *psbA* 5' UTR *in vitro* with properties that are similar to those of the endogenous RB47 protein purified from *C. reinhardtii*. RNA binding to both the *E. coli* expressed and the endogenous RB47 protein was competed using either 200 fold excess of unlabeled *psbA* RNA or 200 fold excess of poly(A) RNA. RNA binding to either of these proteins was poorly competed using 200 fold excess of total RNA or 200 fold excess of the 5' UTR of the *psbD* or *psbC* RNAs. Different forms of the RB47 protein (47 kDa endogenous protein vs. the 69 kDa *E. coli* expressed protein) may account for the slight differences in mobility observed when comparing the binding profiles of purified *C. reinhardtii* protein to heterologously expressed RB47.

The mature form of RB47 is also produced in recombinant form by the insertion by PCR of an artificial stop codon in the RB47 cDNA at nucleotide positions 1403-1405 with a stop codon resulting in a mature RB47 recombinant protein hav-

ing 402 amino acids as shown in FIGS. 1A-1D. An example of this is shown in FIGS. 5A-5B for the production of a recombinant histidine-modified RB47 mature protein as described below. The complete RB47 cDNA is inserted into an expression vector, such as pET3A as described above, for expression of the mature 47 kDa form of the RB47 protein. In the absence of the inserted stop codon, the transcript reads through to nucleotide position 2066-2068 at the TAA stop codon to produce the precursor RB47 having the above-described molecular weight characteristics and 623 amino acid residues.

Recombinant RB47 is also expressed and purified in plant cells. For this aspect, *C. reinhardtii* strains were grown in complete media (Tris-acetate-phosphate [TAP] (Harris, *The Chlamydonas Sourcebook*, San Diego, Calif., Academic Press (1989)) to a density of 5×10^6 cells/mL under constant light. Cells were harvested by centrifugation at 4° C. for 5 minutes at 4,000 g. Cells were either used immediately or frozen in liquid N₂ for storage at -70° C.

Recombinant RB47 protein was also produced as a modified RB47 protein with a histidine tag at the amino-terminus according to well known expression methods using pET19-D vectors available from Novagen, Inc., Madison, Wis. The nucleotide and amino acid sequence of a recombinant histidine-modified RB47 of the mature 47 kDa form is shown in FIGS. 5A-5B with the nucleotide and amino acid sequence also listed in SEQ ID NO 14. Thus the nucleotide sequence of a histidine-modified RB47 is 1269 bases in length. The precursor form of the RB47 protein is similarly obtained in the expression system, both of which are modified by the presence of a histidine tag that allows for purification by metal affinity chromatography.

The recombinant histidine-modified RB47 purified through addition of a poly-histidine tag followed by Ni⁺² column chromatography showed similar binding characteristics as that described for recombinant precursor RB47 described above.

C. Constructs Containing RB60 Nucleotide Sequence

In one approach to obtain purified recombinant RB60 protein, the full-length RB60 cDNA prepared above was cloned into the *E. coli* expression vector pET3A (Studier et al., *Methods Enzymol.*, 185:60-89 (1990)), also commercially available by Novagen, Inc., Madison, Wis. and transformed into BL21 *E. coli* cells. The cells were grown to a density of 0.4 (OD₆₀₀), then induced with 0.5 mM IPTG. Cells were then allowed to grow for an additional 4 hours, at which point they were pelleted and frozen.

Recombinant histidine-modified RB60 was also expressed with a pET19-D vector as described above for RB47 that was similarly modified. Purification of the recombinant RB60 proteins was performed as described for RB47 thereby producing recombinant RB60 proteins for use in the present invention.

The RB60 coding sequence is also mutagenized for directional ligation into an selected vector for expression in alternative systems, such as mammalian expression systems.

D. Constructs Containing an RB47-Encoding Sequence and an RB60-Encoding Sequence

To prepare an expression cassette for encoding both RB47 and RB60, one approach is to digest plasmid pD1/RB47/RB60 prepared above with XhoI and XbaI to isolate the fragment for both encoding sequences. The fragment is then inserted into a similarly digested expression vector if available or is further mutagenized to prepare appropriate restriction sites.

Alternatively, the nucleotide sequences of RB47 and RB60, as described in Example 2, are separately prepared for directional ligation into a selected vector.

An additional embodiment of the present invention is to prepare an expression cassette containing the RB47 binding site along with the coding sequences for RB47 and RB60, the plasmid pD1/RB47/RB60 prepared above is digested with NdeI and XhoI to prepare an expression cassette in which any desired coding sequence having similarly restriction sites is directionally ligated. Expression vectors containing both the RB47 and RB60 encoding sequences in which the RB47 binding site sequence is utilized with a different promoter are also prepared as described in Example 4A.

E. Constructs Containing an RB47 Binding Site Nucleotide Sequence, Insertion Sites for a Desired Heterologous Coding Sequence, and an RB47-Encoding Sequence

In another permutation, a plasmid or expression cassette is constructed containing an RB47 binding site directly upstream of an inserted coding region for a heterologous protein of interest, and the RB47 coding region. The final construct described herein for use in a prokaryotic expression system makes a single mRNA from which both proteins are translated.

The plasmid referred to as pD1/RB47 is prepared as described above in Example 4A. A diagram of this plasmid with the restriction sites is shown in FIG. 6.

Expression of pD1/RB47 in *E. coli* to produce recombinant RB47 and the recombinant heterologous protein is performed as described in above. The heterologous protein is then purified as further described.

To produce an expression cassette that allows for insertion of an alternative desired coding sequence, the plasmid pD1/RB47 is digested with NdeI and XhoI resulting in a vector having restriction endonuclease sites for insertion of a desired coding sequence operably linked to a RB47 binding site and RB47 coding sequence on one transcriptional unit.

F. Constructs Containing an RB47 Binding Site Nucleotide Sequence, Insertion Sites for a Desired Heterologous Coding Sequence, and an RB47-Encoding Sequence

In another permutation, a plasmid or expression cassette is constructed containing an RB47 binding site directly upstream of an inserted coding region for a heterologous protein of interest, and the RB60 coding region. The final construct described herein for use in a prokaryotic expression system makes a single mRNA from which both proteins are translated. In this embodiment, a separate construct encoding recombinant RB47 as described in Example 4B is co-transformed into the *E. coli* host cell for expression.

The plasmid referred to as pD1/RB60 is prepared as described above for pD1/RB47 in Example 4A with the exception that XhoI and XbaI sites are created on RB60 rather than RB47.

Expression of pD1/RB60 in *E. coli* to produce recombinant RB60 and the recombinant heterologous protein is performed as described in above with the combined expression of RB47 from a separate expression cassette. The heterologous protein is then purified as further described.

To produce an expression cassette that allows for insertion of an alternative desired coding sequence, the plasmid pD1/RB60 is digested with NdeI and XhoI resulting in a vector having restriction endonuclease sites for insertion of a desired coding sequence operably linked to a RB47 binding site and RB60 coding sequence on one transcriptional unit.

G. Constructs Containing RB47 Binding Site Nucleotide Sequence and Heterologous Coding Sequences

1) Expression of Recombinant Tetanus Toxin Single Chain Antibody

The examples herein describe constructs that are variations of those described above. The constructs described below contain an RB47 binding site sequence and a heterologous coding sequence. The activating protein RB47 was endogenously provided in the chloroplast and or plant cell. In other aspects however as taught by the methods of the present invention, the chloroplast is further transformed with an RB47-expression construct as described above for overexpression of RB47 to enhance translation capacities.

A strain of the green algae *Chlamydomonas reinhardtii* was designed to allow expression of a single chain antibody gene in the chloroplast. The transgenically expressed antibody was produced from a chimeric gene containing the promoter and 5' untranslated region (UTR) of the chloroplast *psbA* gene prepared as described above, followed by the coding region of a single chain antibody (encoding a tetanus toxin binding antibody), and then the 3' UTR of the *psbA* gene also prepared as described above to provide for transcription termination and RNA processing signals. This construct is essentially pD1/Nde including a heterologous coding sequence having a 3' XbaI restriction site for ligation with the 3' *psbA* gene and is diagramed in FIG. 7.

The *psbA*-single chain construct was first transformed into *C. reinhardtii* chloroplast and transformants were then screened for single chain gene integration. Transformation of chloroplast was performed via bolistic delivery as described in U.S. Pat. Nos. 5,545,818 and 5,553,878, the disclosures of which are hereby incorporated by reference. Transformation is accomplished by homologous recombination via the 5' and 3' UTR of the *psbA* mRNA.

As shown in FIG. 8, two of the transformants that contained the single chain chimeric gene produced single chain antibodies at approximately 1% of total protein levels. The transgenic antibodies were of the correct size and were completely soluble, as would be expected of a correctly folded protein. Few degradation products were detectable by this Western analysis, suggesting that the proteins were fairly stable within the chloroplast. To identify if the produced antibody retained the binding capacity for tetanus toxin, ELISA assays were performed using a mouse-produced Fab, from the original tetanus toxin antibody, as the control. The chloroplast single chain antibody bound tetanus toxin at levels similar to Fab, indicating that the single chain antibody produced in *C. reinhardtii* is a fully functional antibody. These results clearly demonstrate the ability of the chloroplast to synthesis and accumulate function antibody molecules resulting from the translational activation of an RB47 binding site in an expression cassette by endogenous RB47 protein in the chloroplast.

2) Expression of Bacterial Luciferase Enzyme Having Two Subunits

For the production of molecules that contain more than one subunit, such as dIgA and bacterial luciferase enzyme, several proteins must be produced in stoichiometric quantities within the chloroplast. Chloroplast have an advantage for this type of production over cytoplasmic protein synthesis in that translation of multiple proteins can originate from a single mRNA. For example, a dicistronic mRNA having 5' and 3' NdeI and XbaI restriction sites and containing both the A and B chains of the bacterial luciferase enzyme was inserted downstream of the *psbA* promoter and 5' UTR of the pD1/Nde construct prepared in Example 4A above. In this construct, the bacterial LuxAB coding region was ligated

15

between the psbA 5' UTR and the psbA 3' end in an *E. coli* plasmid that was then transformed into *Chlamydomonas reinhardtii* cells as described above for expression in the chloroplast. A schematic of the construct is shown in FIG. 9. Single transformant colonies were then isolated. A plate containing a single isolate was grown for 10 days on complete media and a drop of the luciferase substrate n-Decyl Aldehyde was placed on the plate and the luciferase visualized by video-photography in a dark chamber. Both proteins were synthesized from this single mRNA and luciferase activity accumulated within the chloroplast as shown in FIG. 10. Some mRNA within plastids contained as many as 5 separate proteins encoded on a single mRNA.

3) Expression of Dimeric IgA

To generate dimeric IgA, the construct shown in FIG. 11 is engineered so that the psbA promoter and 5' UTR are used to drive the synthesis of the light chain and heavy chains of an antibody, and the J chain normally associated with IgA molecules. The nucleic acid sequences for the dimeric IgA are inserted into the RB47 binding site construct prepared in Example 4A. The construct is then transformed into *C. reinhardtii* cells as previously described for expression of the recombinant dIgA.

Production of these three proteins within the plastid allows for the self assembly of a dimeric IgA (dIgA). Production of this complex is monitored in several ways. First, Southern analysis of transgenic algae is used to identify strains con-

16

taining the polycistronic chimeric dIgA gene. Strains positive for integration of the dIgA gene are screened by Northern analysis to ensure that the chimeric mRNA is accumulating. Western blot analysis using denaturing gels is used to monitor the accumulation of the individual light, heavy and J chain proteins, and native gels Western blot analysis will be used to monitor the accumulation of the assembled dIgA molecule.

By using a single polycistronic mRNA in the context of RB47 regulated translation, two of the potential pitfalls in the assembly of multimeric dIgA molecule are overcome. First, this construct ensures approximately stoichiometric synthesis of the subunits, as ribosomes reading through the first protein are likely to continue to read through the second and third proteins as well. Second, all of the subunits are synthesized in close physical proximity to each other, which increases the probability of the proteins self assembling into a multimeric molecule. Following the production of a strain producing dIgA molecules, the production of dIgA on an intermediate scale by growing algae in 300 liter fermentors is then performed. Larger production scales are then performed thereafter.

The foregoing specification, including the specific embodiments and examples, is intended to be illustrative of the present invention and is not to be taken as limiting. Numerous other variations and modifications can be effected without departing from the true spirit and scope of the invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: *Chlamydomonas reinhardtii*

<400> SEQUENCE: 1

Gln Tyr Gly Phe Val His Phe Glu Asp Gln Ala Ala Ala Asp Arg
 1 5 10 15

<210> SEQ ID NO 2
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: *Chlamydomonas reinhardtii*

<400> SEQUENCE: 2

Gly Phe Gly Phe Ile Asn Phe Lys Asp Ala Glu Ser Ala Ala
 1 5 10

<210> SEQ ID NO 3
 <211> LENGTH: 32
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 3

cagtacggyt tcgtbcaytt cgaggaycag gc

32

<210> SEQ ID NO 4
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

-continued

<400> SEQUENCE: 4

ggaattcggg ttcggyttca tyaacttcaa ggaygcbgag 40

<210> SEQ ID NO 5

<211> LENGTH: 2846

<212> TYPE: DNA

<213> ORGANISM: Chlamydomonas reinhardtii

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (197)..(2065)

<400> SEQUENCE: 5

gaattcgcgg cgcctccgtg gttggctctc atggtgtctt tttgaagagg acctgagcct 60

ttcacccaaa tatatcaaaa aaccggggca accggccaaa aaaattgcaa aagcctctcg 120

taggcacaaa agacctattc tagccatcaa ctttgatccc gacgctgccg tttagctgcg 180

cgtcttgaag tcaagc atg gcg act act gag tcc tcg gcc ccg gcg gcc acc 232
Met Ala Thr Thr Glu Ser Ser Ala Pro Ala Ala Thr
1 5 10acc cag ccg gcc agc acc ccg ctg gcg aac tcg tcg ctg tac gtc ggt 280
Thr Gln Pro Ala Ser Thr Pro Leu Ala Asn Ser Ser Leu Tyr Val Gly
15 20 25gac ctg gag aag gat gtc acc gag gcc cag ctg ttc gag ctc ttc tcc 328
Asp Leu Glu Lys Asp Val Thr Glu Ala Gln Leu Phe Glu Leu Phe Ser
30 35 40tcg gtt ggc cct gtg gcc tcc att cgc gtg tgc cgc gat gcc gtc acg 376
Ser Val Gly Pro Val Ala Ser Ile Arg Val Cys Arg Asp Ala Val Thr
45 50 55 60cgc cgc tcg ctg ggc tac gcc tac gtc aac tac aac agc gct ctg gac 424
Arg Arg Ser Leu Gly Tyr Ala Tyr Val Asn Tyr Asn Ser Ala Leu Asp
65 70 75ccc cag gct gct gac cgc gcc atg gag acc ctg aac tac cat gtc gtg 472
Pro Gln Ala Ala Asp Arg Ala Met Glu Thr Leu Asn Tyr His Val Val
80 85 90aac ggc aag cct atg cgc atc atg tgg tcg cac cgc gac cct tcg gcc 520
Asn Gly Lys Pro Met Arg Ile Met Trp Ser His Arg Asp Pro Ser Ala
95 100 105cgc aag tcg ggc gtc ggc aac atc ttc atc aag aac ctg gac aag acc 568
Arg Lys Ser Gly Val Gly Asn Ile Phe Ile Lys Asn Leu Asp Lys Thr
110 115 120atc gac gcc aag gcc ctg cac gac acc ttc tcg gcc ttc ggc aag att 616
Ile Asp Ala Lys Ala Leu His Asp Thr Phe Ser Ala Phe Gly Lys Ile
125 130 135 140ctg tcc tgc aag gtt gcc act gac gcc aac ggc gtg tcg aag ggc tac 664
Leu Ser Cys Lys Val Ala Thr Asp Ala Asn Gly Val Ser Lys Gly Tyr
145 150 155ggc ttc gtg cac ttc gag gac cag gcc gct gcc gat cgc gcc att cag 712
Gly Phe Val His Phe Glu Asp Gln Ala Ala Ala Asp Arg Ala Ile Gln
160 165 170acc gtc aac cag aag aag att gag ggc aag atc gtg tac gtg gcc ccc 760
Thr Val Asn Gln Lys Lys Ile Glu Gly Lys Ile Val Tyr Val Ala Pro
175 180 185ttc cag aag cgc gct gac cgc ccc agg gca agg acg ttg tac acc aac 808
Phe Gln Lys Arg Ala Asp Arg Pro Arg Ala Arg Thr Leu Tyr Thr Asn
190 195 200gtg ttc gtc aag aac ttg ccg gcc gac atc ggc gac gac gag ctg ggc 856
Val Phe Val Lys Asn Leu Pro Ala Asp Ile Gly Asp Asp Glu Leu Gly
205 210 215 220aag atg gcc acc gag cac ggc gag atc acc agc gcg gtg gtc atg aag 904
Lys Met Ala Thr Glu His Gly Glu Ile Thr Ser Ala Val Val Met Lys

-continued

545	550	555	
atg atg atc ggc gag cgc ctg tac	ccg cag gtg gcg gag ctg cag ccc		1912
Met Met Ile Gly Glu Arg Leu Tyr	Pro Gln Val Ala Glu Leu Gln Pro		
560	565	570	
gac ctg gct ggc aag atc acc ggc atg ctg ctg gag atg gac aac gcc			1960
Asp Leu Ala Gly Lys Ile Thr Gly Met Leu Leu Glu Met Asp Asn Ala			
575	580	585	
gag ctt ctg atg ctt ctg gag tgc cac gag gcg ctg gtg tcc aag gtg			2008
Glu Leu Leu Met Leu Leu Glu Ser His Glu Ala Leu Val Ser Lys Val			
590	595	600	
gac gag gcc atc gct gtg ctg aag cag cac aac gtg att gcc gag gag			2056
Asp Glu Ala Ile Ala Val Leu Lys Gln His Asn Val Ile Ala Glu Glu			
605	610	615	620
aac aag gct taaagcgct gcacgcttgt gcgggctggt ggcgcggcg			2105
Asn Lys Ala			
cgcgccggcg ctgcttgggc cgccggcagc atgggctggt cggacgcggt gtgggagcag			2165
tgettgtgcttctgtggccgc cgtgaagccg cgccgaactg gggcggacgg caggttggcg			2225
ttgacgccgg cgcgccacaa cacaaagttg gtggcgtgaa agtctctggg cgtgctccgg			2285
acggttgtaa ggttttaaga actggctttt ggccgggttg ccgcccaaag gcggaacggc			2345
ggtcttttca ggccaatcac atccggctgg aaaaattctt accaaagcca acccctgcac			2405
ccaaaaattt cgggttccga aagaacactc cccttttttc cggcaacgcg ttctttcaag			2465
gccaatcact ttccgggttg gaagaaaatg ttaccgggaa aaggcgggaa gccccctgca			2525
cccgacaag ttattcgggg tttcgccggg aatgagcaag cgttcgggct gttggccgta			2585
tcgcgaacgc tgcgggggtg tcaggcgcca gaaggaagga tgacgttttg gtgaaggggt			2645
gcaaactgag cacacgagtt ttggcaatag acgtggagaa agtccagtgc ggggtgagggc			2705
ggatagcgga atcaagcgtg gcgggtccct ggcgagacga gacgcttctg ttgttttgct			2765
gagcccttg atggcacaat cgcactgttt tgagcaggcg actgtaaagt gcccgacgct			2825
aaaaaagcgg ccgcgaattc c			2846

<210> SEQ ID NO 6
 <211> LENGTH: 623
 <212> TYPE: PRT
 <213> ORGANISM: Chlamydomonas reinhardtii

<400> SEQUENCE: 6

Met	Ala	Thr	Thr	Glu	Ser	Ser	Ala	Pro	Ala	Ala	Thr	Thr	Gln	Pro	Ala
1				5				10						15	
Ser	Thr	Pro	Leu	Ala	Asn	Ser	Ser	Leu	Tyr	Val	Gly	Asp	Leu	Glu	Lys
			20					25					30		
Asp	Val	Thr	Glu	Ala	Gln	Leu	Phe	Glu	Leu	Phe	Ser	Ser	Val	Gly	Pro
			35				40					45			
Val	Ala	Ser	Ile	Arg	Val	Cys	Arg	Asp	Ala	Val	Thr	Arg	Arg	Ser	Leu
	50					55					60				
Gly	Tyr	Ala	Tyr	Val	Asn	Tyr	Asn	Ser	Ala	Leu	Asp	Pro	Gln	Ala	Ala
65					70					75					80
Asp	Arg	Ala	Met	Glu	Thr	Leu	Asn	Tyr	His	Val	Val	Asn	Gly	Lys	Pro
				85					90					95	
Met	Arg	Ile	Met	Trp	Ser	His	Arg	Asp	Pro	Ser	Ala	Arg	Lys	Ser	Gly
			100					105					110		
Val	Gly	Asn	Ile	Phe	Ile	Lys	Asn	Leu	Asp	Lys	Thr	Ile	Asp	Ala	Lys
			115				120					125			
Ala	Leu	His	Asp	Thr	Phe	Ser	Ala	Phe	Gly	Lys	Ile	Leu	Ser	Cys	Lys

-continued

130				135				140							
Val	Ala	Thr	Asp	Ala	Asn	Gly	Val	Ser	Lys	Gly	Tyr	Gly	Phe	Val	His
145				150						155					160
Phe	Glu	Asp	Gln	Ala	Ala	Ala	Asp	Arg	Ala	Ile	Gln	Thr	Val	Asn	Gln
				165				170						175	
Lys	Lys	Ile	Glu	Gly	Lys	Ile	Val	Tyr	Val	Ala	Pro	Phe	Gln	Lys	Arg
			180					185					190		
Ala	Asp	Arg	Pro	Arg	Ala	Arg	Thr	Leu	Tyr	Thr	Asn	Val	Phe	Val	Lys
		195					200					205			
Asn	Leu	Pro	Ala	Asp	Ile	Gly	Asp	Asp	Glu	Leu	Gly	Lys	Met	Ala	Thr
	210					215					220				
Glu	His	Gly	Glu	Ile	Thr	Ser	Ala	Val	Val	Met	Lys	Asp	Asp	Lys	Gly
225					230					235					240
Gly	Ser	Lys	Gly	Phe	Gly	Phe	Ile	Asn	Phe	Lys	Asp	Ala	Glu	Ser	Ala
				245					250					255	
Ala	Lys	Cys	Val	Glu	Tyr	Leu	Asn	Glu	Arg	Glu	Met	Ser	Gly	Lys	Thr
			260					265					270		
Leu	Tyr	Ala	Gly	Arg	Ala	Gln	Lys	Lys	Thr	Glu	Arg	Glu	Ala	Met	Leu
		275					280					285			
Arg	Gln	Lys	Ala	Glu	Glu	Ser	Lys	Gln	Glu	Arg	Tyr	Leu	Lys	Tyr	Gln
	290						295				300				
Ser	Met	Asn	Leu	Tyr	Val	Lys	Asn	Leu	Ser	Asp	Glu	Glu	Val	Asp	Asp
305					310					315					320
Asp	Ala	Leu	Arg	Glu	Leu	Phe	Ala	Asn	Ser	Gly	Thr	Ile	Thr	Ser	Cys
				325					330					335	
Lys	Val	Met	Lys	Asp	Gly	Ser	Gly	Lys	Ser	Lys	Gly	Phe	Gly	Phe	Val
			340					345					350		
Cys	Phe	Thr	Ser	His	Asp	Glu	Ala	Thr	Arg	Pro	Pro	Val	Thr	Glu	Met
		355					360					365			
Asn	Gly	Lys	Met	Val	Lys	Gly	Lys	Pro	Leu	Tyr	Val	Ala	Leu	Ala	Gln
	370					375					380				
Arg	Lys	Asp	Val	Arg	Arg	Ala	Thr	Gln	Leu	Glu	Ala	Asn	Met	Gln	Ala
385					390					395					400
Arg	Met	Gly	Met	Gly	Ala	Met	Ser	Arg	Pro	Pro	Asn	Pro	Met	Ala	Gly
				405					410					415	
Met	Ser	Pro	Tyr	Pro	Gly	Ala	Met	Pro	Phe	Phe	Ala	Pro	Gly	Pro	Gly
			420					425					430		
Gly	Met	Ala	Ala	Gly	Pro	Arg	Ala	Pro	Gly	Met	Met	Tyr	Pro	Pro	Met
		435					440					445			
Met	Pro	Pro	Arg	Gly	Met	Pro	Gly	Pro	Gly	Arg	Gly	Pro	Arg	Gly	Pro
	450					455					460				
Met	Met	Pro	Pro	Gln	Met	Met	Gly	Gly	Pro	Met	Met	Gly	Pro	Pro	Met
465					470					475					480
Gly	Pro	Gly	Arg	Gly	Arg	Gly	Gly	Arg	Gly	Pro	Ser	Gly	Arg	Gly	Gln
				485					490					495	
Gly	Arg	Gly	Asn	Asn	Ala	Pro	Ala	Gln	Gln	Pro	Lys	Pro	Ala	Ala	Glu
			500					505					510		
Pro	Ala	Ala	Ala	Pro	Ala	Ala	Ala	Ala	Pro	Ala	Ala	Ala	Ala	Pro	Ala
		515					520					525			
Ala	Ala	Ala	Glu	Pro	Glu	Ala	Pro	Ala	Ala	Gln	Gln	Pro	Leu	Thr	Ala
	530					535					540				
Ser	Ala	Leu	Ala	Ala	Ala	Ala	Pro	Glu	Gln	Gln	Lys	Met	Met	Ile	Gly
545					550					555					560

-continued

Glu Arg Leu Tyr Pro Gln Val Ala Glu Leu Gln Pro Asp Leu Ala Gly
 565 570 575

Lys Ile Thr Gly Met Leu Leu Glu Met Asp Asn Ala Glu Leu Leu Met
 580 585 590

Leu Leu Glu Ser His Glu Ala Leu Val Ser Lys Val Asp Glu Ala Ile
 595 600 605

Ala Val Leu Lys Gln His Asn Val Ile Ala Glu Glu Asn Lys Ala
 610 615 620

<210> SEQ ID NO 7
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Chlamydomonas reinhardtii

<400> SEQUENCE: 7

Trp Phe Val Asp Gly Glu Leu Ala Ser Asp Tyr Asn Gly Pro Arg
 1 5 10 15

<210> SEQ ID NO 8
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: Chlamydomonas reinhardtii

<400> SEQUENCE: 8

Gln Leu Ile Leu Trp Thr Thr Ala Asp Asp Leu Lys Ala Asp Ala Glu
 1 5 10 15

Ile Met Thr Val Phe Arg
 20

<210> SEQ ID NO 9
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 9

cgcggatccg aygcbgagat yatgac 26

<210> SEQ ID NO 10
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 10

cgcgaattcg tcatratctc vgcrtc 26

<210> SEQ ID NO 11
 <211> LENGTH: 2413
 <212> TYPE: DNA
 <213> ORGANISM: Chlamydomonas reinhardtii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (16)..(1614)

<400> SEQUENCE: 11

gagtacgttt acgcc atg aac cgt tgg aac ctt ctt gcc ctt acc ctg ggg 51
 Met Asn Arg Trp Asn Leu Leu Ala Leu Thr Leu Gly
 1 5 10

ctg ctg ctg gtg gca gcg ccc ttc acc aag cac cag ttt gct cat gct 99
 Leu Leu Leu Val Ala Ala Pro Phe Thr Lys His Gln Phe Ala His Ala
 15 20 25

-continued

tcc gat gag tat gag gac gac gag gag gac gat gcc ccc gcc gcc cct	147
Ser Asp Glu Tyr Glu Asp Asp Glu Glu Asp Asp Ala Pro Ala Ala Pro	
30 35 40	
aag gac gac gac gtc gac gtt act gtg gtg acc gtc aag aac tgg gat	195
Lys Asp Asp Asp Val Asp Val Thr Val Val Thr Val Lys Asn Trp Asp	
45 50 55 60	
gag acc gtc aag aag tcc aag ttc gcg ctt gtg gag ttc tac gct cct	243
Glu Thr Val Lys Lys Ser Lys Phe Ala Leu Val Glu Phe Tyr Ala Pro	
65 70 75	
tgg tgc ggc cac tgc aag acc ctc aag cct gag tac gct aag gct gcc	291
Trp Cys Gly His Cys Lys Thr Leu Lys Pro Glu Tyr Ala Lys Ala Ala	
80 85 90	
acc gcc ctg aag gct gct gct ccc gat gcc ctt atc gcc aag gtc gac	339
Thr Ala Leu Lys Ala Ala Ala Pro Asp Ala Leu Ile Ala Lys Val Asp	
95 100 105	
gcc acc cag gag gag tcc ctg gcc cag aag ttc ggc gtg cag ggc tac	387
Ala Thr Gln Glu Glu Ser Leu Ala Gln Lys Phe Gly Val Gln Gly Tyr	
110 115 120	
ccc acc ctc aag tgg ttc gtt gat ggc gag ctg gct tct gac tac aac	435
Pro Thr Leu Lys Trp Phe Val Asp Gly Glu Leu Ala Ser Asp Tyr Asn	
125 130 135 140	
ggc ccc cgc gac gct gat ggc att gtt ggc tgg gtg aag aag aag act	483
Gly Pro Arg Asp Ala Asp Gly Ile Val Gly Trp Val Lys Lys Lys Thr	
145 150 155	
ggc ccc ccc gcc gtg acc gtt gag gac gcc gac aag ctg aag tcc ctg	531
Gly Pro Pro Ala Val Thr Val Glu Asp Ala Asp Lys Leu Lys Ser Leu	
160 165 170	
gag gcg gac gct gag gtc gtt gtc gtc ggc tac ttc aag gcc ctg gag	579
Glu Ala Asp Ala Glu Val Val Val Val Gly Tyr Phe Lys Ala Leu Glu	
175 180 185	
ggc gag atc tac gac acc ttc aag tcc tac gcc gcc aag acc gag gac	627
Gly Glu Ile Tyr Asp Thr Phe Lys Ser Tyr Ala Ala Lys Thr Glu Asp	
190 195 200	
gtg gtg ttc gtg cag acc acc agc gcc gac gtc gcc aag gcc gcc ggc	675
Val Val Phe Val Gln Thr Thr Ser Ala Asp Val Ala Lys Ala Ala Gly	
205 210 215 220	
ctg gac gcc gtg gac acc gtg tcc gtg gtc aag aac ttc gcc ggt gag	723
Leu Asp Ala Val Asp Thr Val Ser Val Val Lys Asn Phe Ala Gly Glu	
225 230 235	
gac cgc gcc acc gcc gtc ctg gcc acg gac atc gac act gac tcc ctg	771
Asp Arg Ala Thr Ala Val Leu Ala Thr Asp Ile Asp Thr Asp Ser Leu	
240 245 250	
acc gcg ttc gtc aag tcg gag aag atg ccc ccc acc att gag ttc aac	819
Thr Ala Phe Val Lys Ser Glu Lys Met Pro Pro Thr Ile Glu Phe Asn	
255 260 265	
cag aag aac tct gac aag atc ttc aac agc ggc atc aac aag cag ctg	867
Gln Lys Asn Ser Asp Lys Ile Phe Asn Ser Gly Ile Asn Lys Gln Leu	
270 275 280	
att ctg tgg acc acc gcc gac gac ctg aag gcc gac gcc gag atc atg	915
Ile Leu Trp Thr Thr Ala Asp Asp Leu Lys Ala Asp Ala Glu Ile Met	
285 290 295 300	
act gtg ttc cgc gag gcc agc aag aag ttc aag ggc cag ctg gtg ttc	963
Thr Val Phe Arg Glu Ala Ser Lys Lys Phe Lys Gly Gln Leu Val Phe	
305 310 315	
gtg acc gtc aac aac gag ggc gac ggc gcc gac ccc gtc acc aac ttc	1011
Val Thr Val Asn Asn Glu Gly Asp Gly Ala Asp Pro Val Thr Asn Phe	
320 325 330	
ttc ggc ctc aag ggc gcc acc tcg cct gtg ctg ctg ggc ttc ttc atg	1059
Phe Gly Leu Lys Gly Ala Thr Ser Pro Val Leu Leu Gly Phe Phe Met	
335 340 345	

-continued

gag aag aac aag aag ttc cgc atg gag ggc gag ttc acg gct gac aac	1107
Glu Lys Asn Lys Lys Phe Arg Met Glu Gly Glu Phe Thr Ala Asp Asn	
350 355 360	
gtg gct aag ttc gcc gag agc gtg gtg gac ggc acc gcg cag gcc gtg	1155
Val Ala Lys Phe Ala Glu Ser Val Val Asp Gly Thr Ala Gln Ala Val	
365 370 375 380	
ctc aag tcg gag gcc atc ccc gag gac ccc tat gag gat ggc gtc tac	1203
Leu Lys Ser Glu Ala Ile Pro Glu Asp Pro Tyr Glu Asp Gly Val Tyr	
385 390 395	
aag att gtg ggc aag acc gtg gag tct gtg gtt ctg gac gag acc aag	1251
Lys Ile Val Gly Lys Thr Val Glu Ser Val Val Leu Asp Glu Thr Lys	
400 405 410	
gac gtg ctg ctg gag gtg tac gcc ccc tgg tgc ggc cac tgc aag aag	1299
Asp Val Leu Leu Glu Val Tyr Ala Pro Trp Cys Gly His Cys Lys Lys	
415 420 425	
ctg gag ccc atc tac aag aag ctg gcc aag cgc ttt aag aag gtg gat	1347
Leu Glu Pro Ile Tyr Lys Lys Leu Ala Lys Arg Phe Lys Lys Val Asp	
430 435 440	
tcc gtc atc atc gcc aag atg gat ggc act gag aac gag cac ccc gag	1395
Ser Val Ile Ile Ala Lys Met Asp Gly Thr Glu Asn Glu His Pro Glu	
445 450 455 460	
atc gag gtc aag ggc ttc cct acc atc ctg ttc tat ccc gcc ggc agc	1443
Ile Glu Val Lys Gly Phe Pro Thr Ile Leu Phe Tyr Pro Ala Gly Ser	
465 470 475	
gac cgc acc ccc atc gtg ttc gag ggc ggc gac cgc tcg ctc aag tcc	1491
Asp Arg Thr Pro Ile Val Phe Glu Gly Gly Asp Arg Ser Leu Lys Ser	
480 485 490	
ctg acc aag ttc atc aag acc aac gcc aag atc ccg tac gag ctg ccc	1539
Leu Thr Lys Phe Ile Lys Thr Asn Ala Lys Ile Pro Tyr Glu Leu Pro	
495 500 505	
aag aag ggc tcc gac ggc gac gag ggc acc tcg gac gac aag gac aag	1587
Lys Lys Gly Ser Asp Gly Asp Glu Gly Thr Ser Asp Asp Lys Asp Lys	
510 515 520	
ccc gcg tcc gac aag gac gag ctg taa gcggctatct gaactacccc	1634
Pro Ala Ser Asp Lys Asp Glu Leu	
525 530	
aggtttgag cgtctgctg cgcgcttgcg cttgcacact gtgcatggat gggagttaag	1694
gaggagacgg agcacggagg ctgcgctcgg ttggtggctt ggagcaccgg cagcgcgtga	1754
tccgtcctgg cagcagcaac ggcggagcgg gcgcatattg gcgagagctg gcgagcggct	1814
gttctgggag aggatatgct gccggggcgg aggaagggct aggggcagag atgagagcgt	1874
tacgggctgg catgcgggcg cccgtgcctc tcctgcggt gcagtccttg ctaggagacg	1934
cacggttttg ccaaagagg acgctgtcca cagccctgcg actggaagt ttttaggccc	1994
tgcggtggta gtggtgttg tacggttggtg tgcataagat gaacaacgtt tctctcaaga	2054
cgagactact agtatgctga cgggtgtgt atgtggtgga tggattgtgc cccgaccatg	2114
aagagtgctg tgttgccctg gcgcttctgt cgccttgat gtgctgtggt ccgaacgctg	2174
gagtcactctg ttgaggagcg aggggtgtgt cgggtccgcc cggcacggcc gcgtgatgct	2234
cggatgggga ttgcgagcga gggcaaccgc agcgcagata gcgccgcagc ggatcgagct	2294
agcgcaggat gatgagagcc gggccttcgc ggcgtgggat cagggaggag ccaaggcggg	2354
gtgcatgcga ggaaaacagt gtgcggaaca gaacgggctg caagaacgcc ttgcgcaaa	2413

<210> SEQ ID NO 12

<211> LENGTH: 532

<212> TYPE: PRT

<213> ORGANISM: Chlamydomonas reinhardtii

-continued

<400> SEQUENCE: 12

Met Asn Arg Trp Asn Leu Leu Ala Leu Thr Leu Gly Leu Leu Leu Val
 1 5 10 15
 Ala Ala Pro Phe Thr Lys His Gln Phe Ala His Ala Ser Asp Glu Tyr
 20 25 30
 Glu Asp Asp Glu Glu Asp Asp Ala Pro Ala Ala Pro Lys Asp Asp Asp
 35 40 45
 Val Asp Val Thr Val Val Thr Val Lys Asn Trp Asp Glu Thr Val Lys
 50 55 60
 Lys Ser Lys Phe Ala Leu Val Glu Phe Tyr Ala Pro Trp Cys Gly His
 65 70 75 80
 Cys Lys Thr Leu Lys Pro Glu Tyr Ala Lys Ala Ala Thr Ala Leu Lys
 85 90 95
 Ala Ala Ala Pro Asp Ala Leu Ile Ala Lys Val Asp Ala Thr Gln Glu
 100 105 110
 Glu Ser Leu Ala Gln Lys Phe Gly Val Gln Gly Tyr Pro Thr Leu Lys
 115 120 125
 Trp Phe Val Asp Gly Glu Leu Ala Ser Asp Tyr Asn Gly Pro Arg Asp
 130 135 140
 Ala Asp Gly Ile Val Gly Trp Val Lys Lys Lys Thr Gly Pro Pro Ala
 145 150 155 160
 Val Thr Val Glu Asp Ala Asp Lys Leu Lys Ser Leu Glu Ala Asp Ala
 165 170 175
 Glu Val Val Val Val Gly Tyr Phe Lys Ala Leu Glu Gly Glu Ile Tyr
 180 185 190
 Asp Thr Phe Lys Ser Tyr Ala Ala Lys Thr Glu Asp Val Val Phe Val
 195 200 205
 Gln Thr Thr Ser Ala Asp Val Ala Lys Ala Ala Gly Leu Asp Ala Val
 210 215 220
 Asp Thr Val Ser Val Val Lys Asn Phe Ala Gly Glu Asp Arg Ala Thr
 225 230 235 240
 Ala Val Leu Ala Thr Asp Ile Asp Thr Asp Ser Leu Thr Ala Phe Val
 245 250 255
 Lys Ser Glu Lys Met Pro Pro Thr Ile Glu Phe Asn Gln Lys Asn Ser
 260 265 270
 Asp Lys Ile Phe Asn Ser Gly Ile Asn Lys Gln Leu Ile Leu Trp Thr
 275 280 285
 Thr Ala Asp Asp Leu Lys Ala Asp Ala Glu Ile Met Thr Val Phe Arg
 290 295 300
 Glu Ala Ser Lys Lys Phe Lys Gly Gln Leu Val Phe Val Thr Val Asn
 305 310 315 320
 Asn Glu Gly Asp Gly Ala Asp Pro Val Thr Asn Phe Phe Gly Leu Lys
 325 330 335
 Gly Ala Thr Ser Pro Val Leu Leu Gly Phe Phe Met Glu Lys Asn Lys
 340 345 350
 Lys Phe Arg Met Glu Gly Glu Phe Thr Ala Asp Asn Val Ala Lys Phe
 355 360 365
 Ala Glu Ser Val Val Asp Gly Thr Ala Gln Ala Val Leu Lys Ser Glu
 370 375 380
 Ala Ile Pro Glu Asp Pro Tyr Glu Asp Gly Val Tyr Lys Ile Val Gly
 385 390 395 400
 Lys Thr Val Glu Ser Val Val Leu Asp Glu Thr Lys Asp Val Leu Leu
 405 410 415

-continued

Glu Val Tyr Ala Pro Trp Cys Gly His Cys Lys Lys Leu Glu Pro Ile
 420 425 430

Tyr Lys Lys Leu Ala Lys Arg Phe Lys Lys Val Asp Ser Val Ile Ile
 435 440 445

Ala Lys Met Asp Gly Thr Glu Asn Glu His Pro Glu Ile Glu Val Lys
 450 455 460

Gly Phe Pro Thr Ile Leu Phe Tyr Pro Ala Gly Ser Asp Arg Thr Pro
 465 470 475 480

Ile Val Phe Glu Gly Gly Asp Arg Ser Leu Lys Ser Leu Thr Lys Phe
 485 490 495

Ile Lys Thr Asn Ala Lys Ile Pro Tyr Glu Leu Pro Lys Lys Gly Ser
 500 505 510

Asp Gly Asp Glu Gly Thr Ser Asp Asp Lys Asp Lys Pro Ala Ser Asp
 515 520 525

Lys Asp Glu Leu
 530

<210> SEQ ID NO 13
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Chlamydomonas reinhardtii

<400> SEQUENCE: 13

Cys Gly His Cys
 1

<210> SEQ ID NO 14
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Chlamydomonas reinhardtii

<400> SEQUENCE: 14

Lys Asp Glu Leu
 1

<210> SEQ ID NO 15
 <211> LENGTH: 1424
 <212> TYPE: DNA
 <213> ORGANISM: Chlamydomonas reinhardtii

<220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (252)..(1310)

<220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (279)..(279)
 <223> OTHER INFORMATION: Codon also can encode Ser

<220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (282)..(282)
 <223> OTHER INFORMATION: Codon also can encode Glu

<220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (294)..(294)
 <223> OTHER INFORMATION: Codon also can encode Gly

<220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (306)..(306)
 <223> OTHER INFORMATION: Codon also can encode Asn

<220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (357)..(357)
 <223> OTHER INFORMATION: Codon also can encode Leu

<220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (369)..(369)
 <223> OTHER INFORMATION: Codon also can encode Thr

<220> FEATURE:
 <221> NAME/KEY: misc_feature

-continued

```

<222> LOCATION: (486)..(486)
<223> OTHER INFORMATION: Codon also can encode Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (495)..(495)
<223> OTHER INFORMATION: Codon also can encode Ile
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (510)..(510)
<223> OTHER INFORMATION: Codon also can encode Ala
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (555)..(555)
<223> OTHER INFORMATION: Codon also can encode Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (588)..(588)
<223> OTHER INFORMATION: Codon also can encode Glu
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (600)..(600)
<223> OTHER INFORMATION: Codon also can encode Leu
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (621)..(621)
<223> OTHER INFORMATION: Codon also can encode Ala
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (714)..(714)
<223> OTHER INFORMATION: Codon also can encode Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (729)..(729)
<223> OTHER INFORMATION: Codon also can encode Ile
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1092)..(1092)
<223> OTHER INFORMATION: Codon also can encode Val
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1191)..(1191)
<223> OTHER INFORMATION: Codon also can encode Ile
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1284)..(1284)
<223> OTHER INFORMATION: Codon also can encode Ala
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1287)..(1287)
<223> OTHER INFORMATION: Codon also can encode Ile
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1290)..(1290)
<223> OTHER INFORMATION: Codon also can encode Glu
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1293)..(1293)
<223> OTHER INFORMATION: Codon also can encode Ala
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1296)..(1296)
<223> OTHER INFORMATION: Codon also can encode Pro
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1302)..(1302)
<223> OTHER INFORMATION: Codon also can encode Thr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1308)..(1308)
<223> OTHER INFORMATION: Codon also can encode Gly

```

```

<400> SEQUENCE: 15

```

```

cgctctatatt taatactccg aaggaggcag ttggcaggca actgccactg acgtcccgta      60
agggtaaggg gacgtccact ggcgtcccgt aaggggaagg ggacgtaggt acataaatgt      120
gctaggtaac taacgtttga tttttgtgg tataatatat gtaccatgct tttaatagaa      180
gcttgaattt ataaattaa atatttttac aatattttac ggagaaatta aaactttaa      240

```


-continued

aaaattaaca t atg aca gca att tta gaa cgt cgt gaa aat tct agc cta	290
Met Thr Ala Ile Leu Glu Arg Arg Glu Asn Ser Ser Leu	
1 5 10	
tgg gct cgt ttt tgt gag tgg atc act tca act gaa aac cgt tta tac	338
Trp Ala Arg Phe Cys Glu Trp Ile Thr Ser Thr Glu Asn Arg Leu Tyr	
15 20 25	
atc ggt tgg ttc ggt gta atc atg atc cca tgt ctt ctt act gca aca	386
Ile Gly Trp Phe Gly Val Ile Met Ile Pro Cys Leu Leu Thr Ala Thr	
30 35 40 45	
tca gta ttc atc atc gct ttc atc gct gct ccg cca gta gac atc gat	434
Ser Val Phe Ile Ile Ala Phe Ile Ala Ala Pro Pro Val Asp Ile Asp	
50 55 60	
ggt atc cgt gaa cca gtt tca ggt tct ctt ctt tac ggt aac aac atc	482
Gly Ile Arg Glu Pro Val Ser Gly Ser Leu Leu Tyr Gly Asn Asn Ile	
65 70 75	
att aca ggt gct gta atc cca act tct aac gca atc ggt ctt cac ttc	530
Ile Thr Gly Ala Val Ile Pro Thr Ser Asn Ala Ile Gly Leu His Phe	
80 85 90	
tac cca att tgg gaa gct gct tct cta gac gag tgg tta tac aac ggt	578
Tyr Pro Ile Trp Glu Ala Ala Ser Leu Asp Glu Trp Leu Tyr Asn Gly	
95 100 105	
ggt cct tac caa ctt atc gtt tgt cac ttc ctt cta ggt gta tac tgc	626
Gly Pro Tyr Gln Leu Ile Val Cys His Phe Leu Leu Gly Val Tyr Cys	
110 115 120 125	
tac atg ggt cgt gag tgg gaa tta tct ttc cgt tta ggt atg cgt cca	674
Tyr Met Gly Arg Glu Trp Glu Leu Ser Phe Arg Leu Gly Met Arg Pro	
130 135 140	
tgg atc gct gta gct tac tca gct cca gta gct gca gct tca gct gta	722
Trp Ile Ala Val Ala Tyr Ser Ala Pro Val Ala Ala Ala Ser Ala Val	
145 150 155	
ttc tta gtt tac cct atc ggc caa ggt tca ttc tct gac ggt atg cct	770
Phe Leu Val Tyr Pro Ile Gly Gln Gly Ser Phe Ser Asp Gly Met Pro	
160 165 170	
tta ggt atc tct ggt act ttc aac ttc atg atc gta ttc caa gca gaa	818
Leu Gly Ile Ser Gly Thr Phe Asn Phe Met Ile Val Phe Gln Ala Glu	
175 180 185	
cac aac atc ctt atg cac cca ttc cac atg tta ggt gtt gct ggt gta	866
His Asn Ile Leu Met His Pro Phe His Met Leu Gly Val Ala Gly Val	
190 195 200 205	
ttc ggt ggt tca tta ttc tca gct atg cac ggt tct tta gtt act tca	914
Phe Gly Gly Ser Leu Phe Ser Ala Met His Gly Ser Leu Val Thr Ser	
210 215 220	
tct tta atc cgt gaa aca act gaa aac gaa tca gct aac gaa ggt tac	962
Ser Leu Ile Arg Glu Thr Thr Glu Asn Glu Ser Ala Asn Glu Gly Tyr	
225 230 235	
cgt ttc ggt caa gaa gaa gaa act tac aac att gta gct gct cat ggt	1010
Arg Phe Gly Gln Glu Glu Glu Thr Tyr Asn Ile Val Ala Ala His Gly	
240 245 250	
tac ttt ggt cgt cta atc ttc caa tac gct tct ttc aac aac tct cgt	1058
Tyr Phe Gly Arg Leu Ile Phe Gln Tyr Ala Ser Phe Asn Asn Ser Arg	
255 260 265	
tca tta cac ttc ttc tta gct gct tgg ccg gta atc ggt att tgg ttc	1106
Ser Leu His Phe Phe Leu Ala Ala Trp Pro Val Ile Gly Ile Trp Phe	
270 275 280 285	
act gct tta ggt tta tca act atg gca ttc aac tta aac ggt ttc aac	1154
Thr Ala Leu Gly Leu Ser Thr Met Ala Phe Asn Leu Asn Gly Phe Asn	
290 295 300	
ttc aac caa tca gta gta gac tca caa ggt cgt gta cta aac act tgg	1202
Phe Asn Gln Ser Val Val Asp Ser Gln Gly Arg Val Leu Asn Thr Trp	
305 310 315	

-continued

gca gac atc atc aac cgt gct aac tta ggt atg gaa gta atg cac gag 1250
 Ala Asp Ile Ile Asn Arg Ala Asn Leu Gly Met Glu Val Met His Glu
 320 325 330

cgt aac gct cac aac ttc cct cta gac tta gct tca act aac tct agc 1298
 Arg Asn Ala His Asn Phe Pro Leu Asp Leu Ala Ser Thr Asn Ser Ser
 335 340 345

tca aac aac taa ttttttttta aactaaaata aatctgggta accataccta 1350
 Ser Asn Asn
 350

gtttatttta gtttatacac acttttcata tatatatact taatagctac cataggcagt 1410

tggcaggacg tccc 1424

<210> SEQ ID NO 16
 <211> LENGTH: 352
 <212> TYPE: PRT
 <213> ORGANISM: Chlamydomonas reinhardtii

<400> SEQUENCE: 16

Met Thr Ala Ile Leu Glu Arg Arg Glu Asn Ser Ser Leu Trp Ala Arg
 1 5 10 15

Phe Cys Glu Trp Ile Thr Ser Thr Glu Asn Arg Leu Tyr Ile Gly Trp
 20 25 30

Phe Gly Val Ile Met Ile Pro Cys Leu Leu Thr Ala Thr Ser Val Phe
 35 40 45

Ile Ile Ala Phe Ile Ala Ala Pro Pro Val Asp Ile Asp Gly Ile Arg
 50 55 60

Glu Pro Val Ser Gly Ser Leu Leu Tyr Gly Asn Asn Ile Ile Thr Gly
 65 70 75 80

Ala Val Ile Pro Thr Ser Asn Ala Ile Gly Leu His Phe Tyr Pro Ile
 85 90 95

Trp Glu Ala Ala Ser Leu Asp Glu Trp Leu Tyr Asn Gly Gly Pro Tyr
 100 105 110

Gln Leu Ile Val Cys His Phe Leu Leu Gly Val Tyr Cys Tyr Met Gly
 115 120 125

Arg Glu Trp Glu Leu Ser Phe Arg Leu Gly Met Arg Pro Trp Ile Ala
 130 135 140

Val Ala Tyr Ser Ala Pro Val Ala Ala Ala Ser Ala Val Phe Leu Val
 145 150 155 160

Tyr Pro Ile Gly Gln Gly Ser Phe Ser Asp Gly Met Pro Leu Gly Ile
 165 170 175

Ser Gly Thr Phe Asn Phe Met Ile Val Phe Gln Ala Glu His Asn Ile
 180 185 190

Leu Met His Pro Phe His Met Leu Gly Val Ala Gly Val Phe Gly Gly
 195 200 205

Ser Leu Phe Ser Ala Met His Gly Ser Leu Val Thr Ser Ser Leu Ile
 210 215 220

Arg Glu Thr Thr Glu Asn Glu Ser Ala Asn Glu Gly Tyr Arg Phe Gly
 225 230 235 240

Gln Glu Glu Glu Thr Tyr Asn Ile Val Ala Ala His Gly Tyr Phe Gly
 245 250 255

Arg Leu Ile Phe Gln Tyr Ala Ser Phe Asn Asn Ser Arg Ser Leu His
 260 265 270

Phe Phe Leu Ala Ala Trp Pro Val Ile Gly Ile Trp Phe Thr Ala Leu
 275 280 285

Gly Leu Ser Thr Met Ala Phe Asn Leu Asn Gly Phe Asn Phe Asn Gln

-continued

290	295	300	
Ser Val Val Asp Ser Gln Gly Arg Val Leu Asn Thr Trp Ala Asp Ile			
305	310	315	320
Ile Asn Arg Ala Asn Leu Gly Met Glu Val Met His Glu Arg Asn Ala			
	325	330	335
His Asn Phe Pro Leu Asp Leu Ala Ser Thr Asn Ser Ser Ser Asn Asn			
	340	345	350

<210> SEQ ID NO 17
 <211> LENGTH: 1278
 <212> TYPE: DNA
 <213> ORGANISM: Chlamydomonas reinhardtii
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(1272)

<400> SEQUENCE: 17

atg ggc cat cat cat cat cat cat cat cat cat cac agc agc ggc cat	48
Met Gly His His His His His His His His His His Ser Ser Gly His	
1	5 10 15
atc gaa ggt cgt cat atg gcg act act gag tcc tcg gcc ccg gcg gcc	96
Ile Glu Gly Arg His Met Ala Thr Thr Glu Ser Ser Ala Pro Ala Ala	
	20 25 30
acc acc cag ccg gcc agc acc ccg ctg gcg aac tcg tcg ctg tac gtc	144
Thr Thr Gln Pro Ala Ser Thr Pro Leu Ala Asn Ser Ser Leu Tyr Val	
	35 40 45
ggt gac ctg gag aag gat gtc acc gag gcc cag ctg ttc gag ctc ttc	192
Gly Asp Leu Glu Lys Asp Val Thr Glu Ala Gln Leu Phe Glu Leu Phe	
	50 55 60
tcc tcg gtt ggc cct gtg gcc tcc att cgc gtg tgc cgc gat gcc gtc	240
Ser Ser Val Gly Pro Val Ala Ser Ile Arg Val Cys Arg Asp Ala Val	
	65 70 75 80
acg cgc cgc tcg ctg ggc tac gcc tac gtc aac tac aac agc gct ctg	288
Thr Arg Arg Ser Leu Gly Tyr Ala Tyr Val Asn Tyr Asn Ser Ala Leu	
	85 90 95
gac ccc cag gct gct gac cgc gcc atg gag acc ctg aac tac cat gtc	336
Asp Pro Gln Ala Ala Asp Arg Ala Met Glu Thr Leu Asn Tyr His Val	
	100 105 110
gtg aac ggc aag cct atg cgc atc atg tgg tcg cac cgc gac cct tcg	384
Val Asn Gly Lys Pro Met Arg Ile Met Trp Ser His Arg Asp Pro Ser	
	115 120 125
gcc cgc aag tcg ggc gtc gcc aac atc ttc atc aag aac ctg gac aag	432
Ala Arg Lys Ser Gly Val Gly Asn Ile Phe Ile Lys Asn Leu Asp Lys	
	130 135 140
acc atc gac gcc aag gcc ctg cac gac acc ttc tcg gcc ttc ggc aag	480
Thr Ile Asp Ala Lys Ala Leu His Asp Thr Phe Ser Ala Phe Gly Lys	
	145 150 155 160
att ctg tcc tgc aag gtt gcc act gac gcc aac ggc gtg tcg aag ggc	528
Ile Leu Ser Cys Lys Val Ala Thr Asp Ala Asn Gly Val Ser Lys Gly	
	165 170 175
tac ggc ttc gtg cac ttc gag gac cag gcc gct gcc gat cgc gcc att	576
Tyr Gly Phe Val His Phe Glu Asp Gln Ala Ala Ala Asp Arg Ala Ile	
	180 185 190
cag acc gtc aac cag aag aag att gag gcc aag atc gtg tac gtg gcc	624
Gln Thr Val Asn Gln Lys Lys Ile Glu Gly Lys Ile Val Tyr Val Ala	
	195 200 205
ccc ttc cag aag cgc gct gac cgc ccc agg gca agg acg ttg tac acc	672
Pro Phe Gln Lys Arg Ala Asp Arg Pro Arg Ala Arg Thr Leu Tyr Thr	
	210 215 220
aac gtg ttc gtc aag aac ttg ccg gcc gac atc ggc gac gac gag ctg	720
Asn Val Phe Val Lys Asn Leu Pro Ala Asp Ile Gly Asp Asp Glu Leu	

-continued

115	120	125
Ala Arg Lys Ser Gly Val Gly Asn Ile Phe Ile Lys Asn Leu Asp Lys 130 135 140		
Thr Ile Asp Ala Lys Ala Leu His Asp Thr Phe Ser Ala Phe Gly Lys 145 150 155 160		
Ile Leu Ser Cys Lys Val Ala Thr Asp Ala Asn Gly Val Ser Lys Gly 165 170 175		
Tyr Gly Phe Val His Phe Glu Asp Gln Ala Ala Ala Asp Arg Ala Ile 180 185 190		
Gln Thr Val Asn Gln Lys Lys Ile Glu Gly Lys Ile Val Tyr Val Ala 195 200 205		
Pro Phe Gln Lys Arg Ala Asp Arg Pro Arg Ala Arg Thr Leu Tyr Thr 210 215 220		
Asn Val Phe Val Lys Asn Leu Pro Ala Asp Ile Gly Asp Asp Glu Leu 225 230 235 240		
Gly Lys Met Ala Thr Glu His Gly Glu Ile Thr Ser Ala Val Val Met 245 250 255		
Lys Asp Asp Lys Gly Gly Ser Lys Gly Phe Gly Phe Ile Asn Phe Lys 260 265 270		
Asp Ala Glu Ser Ala Ala Lys Cys Val Glu Tyr Leu Asn Glu Arg Glu 275 280 285		
Met Ser Gly Lys Thr Leu Tyr Ala Gly Arg Ala Gln Lys Lys Thr Glu 290 295 300		
Arg Glu Ala Met Leu Arg Gln Lys Ala Glu Glu Ser Lys Gln Glu Arg 305 310 315 320		
Tyr Leu Lys Tyr Gln Ser Met Asn Leu Tyr Val Lys Asn Leu Ser Asp 325 330 335		
Glu Glu Val Asp Asp Asp Ala Leu Arg Glu Leu Phe Ala Asn Ser Gly 340 345 350		
Thr Ile Thr Ser Cys Lys Val Met Lys Asp Gly Ser Gly Lys Ser Lys 355 360 365		
Gly Phe Gly Phe Val Cys Phe Thr Ser His Asp Glu Ala Thr Arg Pro 370 375 380		
Pro Val Thr Glu Met Asn Gly Lys Met Val Lys Gly Lys Pro Leu Tyr 385 390 395 400		
Val Ala Leu Ala Gln Arg Lys Asp Val Arg Arg Ala Thr Gln Leu Glu 405 410 415		
Ala Asn Met Gln Ala Arg Met 420		

What is claimed is:

[1. An expression cassette for expression of a desired molecule, which cassette comprises:

- a) an RB47 binding site nucleotide sequence upstream of a restriction endonuclease site for insertion of a desired coding sequence to be expressed; and
- b) a nucleotide sequence encoding a polypeptide which binds RB47 binding site.]

[2. The expression cassette of claim 1 further comprising a promoter sequence operably linked to and positioned upstream of the RB47 binding site nucleotide sequence.]

[3. The expression cassette of claim 2 wherein the promoter sequence is derived from a psbA gene.]

[4. The expression cassette of claim 3 wherein the coding sequence is heterologous to the psbA gene.]

[5. The expression cassette of claim 1 wherein the cassette comprises a plasmid or virus.]

[6. The expression cassette of claim 1 further comprising and operably linked thereto a nucleotide sequence encoding RB60.]

[7. The expression cassette of claim 1 wherein the RB47 binding polypeptide is selected from the group consisting of RB47, RB47 precursor and a histidine-modified RB47.]

[8. An expression cassette for expression of a desired molecule, which cassette comprises:

- a) an RB47 binding site nucleotide sequence upstream of a restriction endonuclease site for insertion of a desired coding sequence to be expressed;

and

- b) a nucleotide sequence encoding a polypeptide which regulates the binding of RB47 to the RB47 binding site.]

[9. The expression cassette of claim 8 wherein the regulatory polypeptide is RB60.]

[10. A method of screening for agonists or antagonists of RB47 binding to RB47 binding site, the method comprising the steps:

- a) providing a cell expression system containing:
 - 1) a promoter sequence;
 - 2) a RB47 binding site sequence;
 - 3) a coding sequence for an indicator polypeptide; and
 - 4) a polypeptide which binds to the RB47 binding site sequence;
- b) introducing an antagonist or agonist into the cell; and
- c) detecting the amount of indicator polypeptide expressed in the cell.]

[11. A method of screening for agonists or antagonists of RB60 in regulating RB47 binding to RB47 binding site, the method comprising the steps:

- a) providing an expression system in a cell containing:
 - 1) a promoter sequence;
 - 2) a RB47 binding site sequence;
 - 3) a coding sequence for an indicator polypeptide;
 - 4) a polypeptide which binds to the RB47 binding site sequence; and
 - 5) a RB60 polypeptide;
- b) introducing an agonist or antagonist into the cell; and
- c) detecting the amount of indicator polypeptide expressed in the cell.]

[12. An isolated nucleotide sequence encoding RB47.]

[13. An isolated nucleotide sequence encoding a histidine-modified RB47.]

[14. An isolated nucleotide sequence encoding RB47 precursor.]

[15. The nucleotide sequence of claim 12 from nucleotide position 197 to 1402 in FIGS. 1A-1B and SEQ ID NO 5.]

[16. The nucleotide sequence of claim 13 from nucleotide position 1 to 1269 in FIGS. 5A-5B and SEQ ID NO 14.]

[17. The nucleotide sequence of claim 14 shown in from nucleotide position 197 to 2065 in FIGS. 1A-1C and SEQ ID NO 5.]

[18. An expression cassette comprising the nucleotide sequence of claim 12, 13 or 14.]

[19. An isolated nucleotide sequence encoding RB60.]

[20. The nucleotide sequence of claim 18 from nucleotide position 16 to 1614 in FIGS. 2A-2B and SEQ ID NO 10.]

[21. An expression cassette comprising the nucleotide sequence of claim 19.]

[22. An expression system comprising a cell transformed with the expression cassette of claim 1.]

[23. The expression system of claim 22 wherein the cell is a plant cell.]

[24. The expression system of claim 23 wherein the plant cell endogenously expresses RB47.]

[25. The expression system of claim 23 wherein the plant cell endogenously expresses RB60.]

[26. The expression system of claim 23 wherein the plant cell endogenously expresses RB47 and RB60.]

[27. The expression system of claim 22 wherein the cell is a eukaryotic cell.]

[28. The expression system of claim 22 wherein the cell is a prokaryotic cell.]

[29. The expression system of claim 22 further comprising the expression cassette of claim 21.]

[30. An expression system comprising a cell transformed with the expression cassette of claim 8.]

[31. The expression system of claim 29 further comprising the expression cassette of claim 18.]

[32. A cell stably transformed with the expression cassette of claim 18.]

[33. A cell stably transformed with the expression cassette of claim 21.]

[34. A cell stably transformed with the expression cassette of claims 18 and 21.]

[35. The expression cassette of claim 1 further comprising an inserted desired coding sequence.]

[36. An expression system comprising a cell transformed with the expression cassette of claim 35, wherein the coding sequence is expressed forming the desired molecule upon activation of the RB47 binding site with RB47.]

[37. The expression system of claim 36 wherein the cell is a plant cell endogenously expressing RB47.]

[38. The expression system of claim 36 wherein the cell is stably transformed with the expression cassette of claim 21.]

[39. An expression system comprising a cell transformed with an expression cassette comprising a promoter sequence, a RB47 binding site sequence, a desired coding sequence for a molecule, and a nucleotide sequence for encoding a polypeptide which binds RB47 binding site, wherein all sequences are operably linked.]

[40. A method of preparing a desired recombinant molecule wherein the method comprises cultivating the expression system of claim 36.]

[41. A method of preparing a desired recombinant molecule wherein the method comprises cultivating the expression system of claim 39.]

[42. A method for expressing a desired coding sequence comprising:

- a) forming an expression cassette by operably linking:

- 1) a promoter sequence;
- 2) a RB47 binding site sequence;
- 3) a desired coding sequence; and
- 4) a nucleotide sequence encoding a polypeptide which binds RB47 binding site; and

- b) introducing the expression cassette into a cell.]

[43. The method of claim 42 wherein the cell is a plant cell endogenously expressing RB47.]

[44. The method of claim 42 wherein the cell is a plant cell endogenously expressing RB60.]

[45. The method of claim 42 further comprising inducing expression with a promoter inducer molecule.]

[46. The method of claim 45 wherein the promoter inducer molecule is IPTG.]

[47. The method of claim 42 wherein the cell is transformed with the expression cassette of claim 21.]

[48. A method for expressing a desired coding sequence comprising:

- a) forming an expression cassette by operably linking:

- 1) a promoter sequence;
- 2) a RB47 binding site sequence; and
- 3) a desired coding sequence;

and

- b) introducing the expression cassette into a plant cell endogenously expressing RB47.]

[49. The method of claim 48 wherein the expression cassette further comprises a nucleotide sequence encoding RB60.]

[50. A method for the regulated production of a recombinant molecule from a desired coding sequence in a cell, wherein the cell contains the expression cassette of claim 34, wherein expression of the coding sequence is activated by RB47 binding to the RB47 binding site thereby producing the recombinant molecule.]

[51. A method of forming an expression cassette by operably linking:

- a) a RB47 binding site sequence;
- b) a cloning site for insertion of a desired coding sequence downstream of the RB47 binding site sequence; and
- c) a nucleotide sequence encoding a polypeptide which binds the RB47 binding site.]

[52. The method of claim 51 further comprising a promoter sequence operably linked upstream to the RB47 binding site sequence.]

[53. The method of claim 51 further comprising a desired coding sequence inserted into the insertion site.]

[54. An article of manufacture comprising a packaging material and contained therein in a separate container the expression cassette of claim 1, wherein the expression cassette is useful for expression of a desired coding sequence, and wherein the packaging material comprises a label which indicates that the expression cassette can be used for expressing a desired coding sequence when the RB47 binding site is activated by RB47.]

[55. The article of manufacture of claim 54 further comprising in a separate container the expression cassette of claim 18.]

[56. The article of manufacture of claim 54 further comprising in a separate container the expression cassette of claim 21.]

[57. An article of manufacture comprising a packaging material and contained therein in a separate container the expression system of claim 22, wherein the expression system is useful for expression of a desired coding sequence, and wherein the packaging material comprises a label which indicates that the expression system can be used for expressing a desired coding sequence when the RB47 binding site is activated by RB47.]

[58. An article of manufacture comprising a packaging material and contained therein in a separate container the stably transformed cell of claim 32, wherein the cell is useful as an expression system, and wherein the packaging material comprises a label which indicates that the expression system can be used for expressing a desired coding sequence when the RB47 binding site is activated by RB47.]

[59. An article of manufacture comprising a packaging material and contained therein in a separate container the stably transformed cell of claim 33, wherein the cell is useful as an expression system, and wherein the packaging material comprises a label which indicates that the expression system can be used for expressing a desired coding sequence when the RB47 binding site is activated by RB47 and regulated by RB60.]

[60. An article of manufacture comprising a packaging material and contained therein in a separate container the stably transformed cell of claim 34, wherein the cell is useful as an expression system, and wherein the packaging material comprises a label which indicates that the expression system can be used for expressing a desired coding sequence when the RB47 binding site is activated by RB47 and regulated by RB60.]

[61. An article of manufacture comprising a packaging material and contained therein in a separate container the expression cassette of claim 2, wherein the expression cassette is useful for expression of a RNA transcript, and wherein the packaging material comprises a label which indicates that the expression cassette can be used for producing in vitro a RNA transcript when the RB47 binding site is activated by RB47.]

[62. The article of manufacture of claim 61 wherein the promoter sequence is selected from the group consisting of T3 and T7 promoters.]

[63. The article of manufacture of claim 61 further comprising in separate containers a polymerase, a buffer and each of four ribonucleotides, reagents for in vitro RNA transcription.]

64. A chloroplast expression cassette comprising the following components in the 5' to 3' direction of transcription:

a) a promoter functional in a chloroplast;

b) a 5' leader sequence comprising a 5' untranslated region (UTR), wherein the 5' UTR comprises an RB47 binding site; and

c) a DNA sequence encoding a heterologous protein of interest.

65. The chloroplast expression cassette of claim 64, wherein the DNA sequence encodes a vertebrate polypeptide.

66. The chloroplast cassette of claim 64, wherein the DNA sequence encodes a mammalian polypeptide.

67. The chloroplast expression cassette of claim 64, wherein the polypeptide is an antibody.

68. The chloroplast cassette of claim 67, wherein the polypeptide is a single chain antibody.

69. The chloroplast cassette of claim 64, wherein the chloroplast is a plant chloroplast.

70. The chloroplast cassette of claim 64, wherein the chloroplast is an algal chloroplast.

71. The chloroplast expression cassette of claim 64, wherein the 5' leader sequence is a 5' untranslated region (UTR).

72. A cell containing the chloroplast expression cassette of claim 64.

73. An alga or plant comprising a cell of claim 72.

74. An algal chloroplast comprising the expression cassette of claim 64.

75. A micro-algae containing a chloroplast of claim 74.

76. The micro-algae of claim 75, wherein the algae is Chlamydomonas reinhardtii.

77. The chloroplast expression cassette of claim 64, further comprising a 3' UTR.

78. The expression cassette of claim 64, wherein the DNA sequence encodes a eukaryotic protein.

79. The chloroplast expression cassette of claim 77, wherein the promoter and the 5' leader sequence and the 3' UTR are of a length which allows for replacement of a homologous gene by genetic recombination upon introduction into the chloroplast genome.

80. The chloroplast expression cassette of claim 79, wherein the homologous gene to be replaced is a psbA gene.

81. A method for producing a non-plant, non-plastid protein in a chloroplast, comprising:

a) transforming a chloroplast of a cell with a cassette of claim 64, and

b) growing the cell comprising the transformed chloroplast under conditions wherein the DNA sequence is expressed to produce the protein in the chloroplast.

82. A eukaryotic cell comprising a transformed chloroplast producing a protein according to the method of claim 81.

83. The chloroplast expression cassette of claim 79, wherein the protein is an antibody.

84. A microalgal chloroplast transformed with an expression cassette of claim 65.

85. The microalgal chloroplast of claim 84, wherein said microalga is Chlamydomonas reinhardtii.

86. A method for producing a heterologous eukaryotic protein in a microalgal chloroplast, comprising:

a) transforming a microalgal chloroplast of a cell with a cassette of claim 65, and

b) growing the cell comprising the transformed microalgal chloroplast under conditions wherein the DNA sequence is expressed to produce the protein in the microalgal chloroplast.

* * * * *